

**BASIC CANON IN  $D = 4$ ,  $N = 1$  SUPERFIELD THEORY:  
Five Primer Lectures**

S. JAMES GATES, JR.

*Department of Physics, University of Maryland,  
College Park, MD 20742-4111, USA  
E-mail: [gatess@wam.umd.edu](mailto:gatess@wam.umd.edu)*

The topic of  $4D$ ,  $N = 1$  supersymmetry is introduced for the reader with a prior background in relativistic quantum field theory. The presentation is designed to be a useful primer for those who plan to later engage in serious investigation of the area or as an overview for the generally interested.

## Prologue

At MIT when I was a graduate student looking for a topic on which I would write my doctoral thesis, one thing I did was to make a survey of the research literature. There were many topics that were of current interest; “the Bag Model,” the dilute gas approximation of QCD, unified electroweak models, instanton physics and numbers of other things. However, being a graduate student, I figured that the best thing on which to work must be something completely new. After all if it was as new as possible, then I had a fair chance to compete with everyone else who also did not know anything about this new topic. I very rapidly came to the conclusion that my choice ought to be the (then) newly discovered class of quantum field theories which possessed a property called “supersymmetry.” I could tell it was new because in the first article I read about it, there appeared the notion of a gradient operator which possessed a *spinorial* index! I had never seen anything similar in any of my classes, so this seemed a pretty good indication that this was really *new*.

For a while it seemed as though my approach had worked too well. No one with whom I spoke in the department knew anything about this stuff that involved “superspace” and “superfields.” Worse yet, it seemed as though it was slightly disreputable to work on this fanciful topic especially when there was ‘real physics’ to be done. So there were no obvious faculty members to whom I could be an apprentice. I did, however, find a thesis adviser from whom I had learned a lot about electroweak models and he trusted me enough to let me follow my nose. But he made it clear that I would be totally responsible for teaching myself the topic, although he was available to discuss things whenever I got stuck. Thus, I was off on the adventure of mentally going to superspace and learning about the lay of the land. The fact that I have been invited to give these lectures is a hint that I succeeded in learning something about this place.

In preparing these lectures, I have decided by and large to keep them as simple as possible. In fact, my idea for these discussions is an attempt to try to recall the things that I found confusing when I was first teaching myself. The best way to accomplish this seems for me to imagine that you had met me while I was working on my doctoral thesis and then to tell you about the kinds of things on which I worked. The one major exception to this is the final section of the text. There I discuss a form of supergravity with some unique properties and which was first presented in 1989. I am particularly moved to make this exception because this special form of supergravity, in my opinion,

will ultimately be found to be the 4D,  $N = 1$  supergeometrical limit of heterotic and superstring theories.

So these discussions are designed almost exclusively for the student who has completed courses in relativistic quantum field theory, group theory, linear algebra and a few other useful mathematical topics and who is embarking on the quest of learning about supersymmetry. As Minkowski space is the natural setting for a discussion of relativistic theories, so is superspace the natural setting for a discussion of supersymmetry. Today this does not seem such a radical notion. But when I was learning and creating some of this stuff, there were actually places where graduate students, even those working on supersymmetric topics, were extremely discouraged from learning how to use superspace and superfields.

These lectures are extremely limited in their scope and choice of topics. The student is encouraged to view these discussions solely as a primer. There are by now a plethora of books available on supersymmetry and supersymmetrical field theory. I regard most books on this topic as giving a “beginning” to “intermediate” level treatment. As such they are pretty interchangeable. There are, to my knowledge, only two books that give an “advanced” treatment. I hope that those who read these lectures will easily be able to go on to any real textbook of their choice having benefitted from my 1997 TASI lectures.

I would be remiss if I did not thank the organizers, Jon Bagger and K. T. Mahanthappa, for their kind invitation to make these presentations. Additional thanks are extended to Ms. Betty Krusberg and Ms. Delores Kight for their assistance in the preparation of the manuscript. Of course, any errors are my own responsibility.

## 1 First Lecture: Tyro 4D, N = 1 Superfield Theory

### *Introduction*

In this first lecture, I will endeavor to get you, the reader, to become comfortable with the notion of superspace. The simplest and easiest way to do this is to recall an experience which you most likely have already had if you are reading this. Namely, you experienced a “paradigm shift” when you first encountered special relativity and the four vector formulation of theories consistent with its symmetries. This must have forced a radical change in your notions of space and time which became the Minkowski space-time manifold  $\mathcal{M}_4$ . In a similar way, I wish for you to make another paradigm shift. The best way to gain insight into supersymmetrical theories is to jump right into superspace where the Minkowski space-time manifold is replaced by the Salam-Strathdee superspace manifold  $s\mathcal{M}_{(4,N)}$ . We won’t actually learn very much superfield theory in this lecture. But we will learn a lot about the appropriate setting for these theories.

### *Lecture*

#### *1.1 Minkowski Space: The Boundary of Superspace*

Ordinary 4D relativistic field theory is defined in terms of fields over a four dimensional manifold  $\mathcal{M}_4$ . The objects that make up this manifold are points that may be labeled by four numbers such as  $x^{\underline{m}} = (x^0, x^1, x^2, x^3)$ . The 4D Minkowski manifold is actually an equivalence class. We regard the manifold with coordinates given by  $X^{\underline{m}}$  as an equivalent one if

$$X^{\underline{m}} = x^{\underline{n}} (K_0)_{\underline{n}}^{\underline{m}} + (k_0)^{\underline{m}} \quad , \quad (1)$$

where  $(K_0)_{\underline{n}}^{\underline{m}}$  and  $(k_0)^{\underline{m}}$  are sets of constants such that

$$(dX^{\underline{m}}) \eta_{\underline{m}\underline{n}} (dX^{\underline{n}}) = (dx^{\underline{m}}) \eta_{\underline{m}\underline{m}} (dx^{\underline{n}}) \quad ,$$

$$\eta_{\underline{m}\underline{n}} = \begin{pmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad , \quad (2)$$

which implies the following restrictions on the matrix  $(K_0)_{\underline{n}}^{\underline{m}}$

$$\eta_{\underline{m}\underline{n}} = (K_0)_{\underline{m}}^{\underline{r}} \eta_{\underline{r}\underline{s}} (K_0)_{\underline{n}}^{\underline{s}} \quad . \quad (3)$$

We can also write

$$X^{\underline{m}} = \exp[i\frac{1}{2}(K_0)_{\underline{r}}^{\underline{s}}L_{\underline{s}}^{\underline{r}}] \exp[i(k_0)^{\underline{m}}P_{\underline{m}}] x^{\underline{m}} \quad , \quad (4)$$

where  $P_{\underline{m}}$  and  $L_{\underline{s}}^{\underline{r}}$  are differential operators

$$P_{\underline{m}} \equiv \frac{1}{i} \frac{\partial}{\partial x^{\underline{m}}} = -i \partial_{\underline{m}} \quad , \quad L_{\underline{s}}^{\underline{r}} \equiv \eta^{\underline{r}\underline{t}} L_{\underline{s}\underline{t}} = \eta^{\underline{r}\underline{t}} [ -i ( x_{\underline{s}} \partial_{\underline{t}} - x_{\underline{t}} \partial_{\underline{s}} ) ] \quad . \quad (5)$$

It is easy to prove that

$$\begin{aligned} [P_{\underline{r}} , P_{\underline{s}}] &= 0 \quad , \quad [L_{\underline{r}\underline{s}} , P_{\underline{t}}] = i ( \eta_{\underline{r}\underline{t}} P_{\underline{s}} - \eta_{\underline{s}\underline{t}} P_{\underline{r}} ) \quad , \\ [L_{\underline{r}\underline{s}} , L_{\underline{t}\underline{u}}] &= i ( \eta_{\underline{r}\underline{t}} L_{\underline{s}\underline{u}} - \eta_{\underline{s}\underline{t}} L_{\underline{r}\underline{u}} - \eta_{\underline{r}\underline{u}} L_{\underline{s}\underline{t}} + \eta_{\underline{s}\underline{u}} L_{\underline{r}\underline{t}} ) \quad . \end{aligned} \quad (6)$$

This is the Poincaré Lie algebra. The Poincaré algebra is exponentiated to form the Poincaré group. The operators  $P_{\underline{m}}$  and  $L_{\underline{r}\underline{s}}$  respectively generate “translations” and “Lorentz rotations.” When exponentiated each of these also separately form groups.

All 4D fields are<sup>a</sup> representations of these groups. The simplest such fields are

$$\begin{aligned} \phi(x) \quad , \quad \psi_{(\alpha)}(x) \quad , \quad A_{\underline{a}}(x), \dots \\ \text{scalar} \quad , \quad \text{spinor} \quad , \quad \text{vector} \end{aligned} \quad (7)$$

and this list goes on forever where fields are defined with increasing numbers of four-component spinor ( $\alpha$ ) or 4-vector  $\underline{a}$  indices. These various possibilities arise because there exist constant numerical matrices ( $\tilde{S}_{\underline{a}\underline{b}}$ ) which satisfy the same commutation algebra as that of the  $L$ 's.

$$\begin{aligned} \text{Ex. 1 : } \quad \tilde{S}_{\underline{a}\underline{b}} &\equiv 0 \quad , \\ \text{Ex. 2 : } \quad \tilde{S}_{\underline{a}\underline{b}} &\equiv i\frac{1}{4} (\sigma_{\underline{a}\underline{b}})_{(\alpha)}^{(\beta)} \quad , \\ \text{Ex. 3 : } \quad \tilde{S}_{\underline{a}\underline{b}} &\equiv i ( \eta_{\underline{a}\underline{r}} \delta_{\underline{b}}^{\underline{s}} - \eta_{\underline{b}\underline{r}} \delta_{\underline{a}}^{\underline{s}} ) \quad . \end{aligned} \quad (8)$$

The total angular momentum operator for any field is  $J_{\underline{a}\underline{b}} = L_{\underline{a}\underline{b}} + S_{\underline{a}\underline{b}}$  where the explicit matrix form for  $S_{\underline{a}\underline{b}}$  depends on the spin of the field upon which  $S_{\underline{a}\underline{b}}$  acts. This allows us to define an abstract Lie algebra operator  $\mathcal{M}_{\underline{a}\underline{b}} = -iS_{\underline{a}\underline{b}}$

---

<sup>a</sup>Following Wigner, we may say that the definition of an elementary particle is that it is a representation of Lie algebras.

so that

$$\begin{aligned}
\mathcal{M}_{\underline{a}\underline{b}}\phi(x) &= 0 \quad , \\
\mathcal{M}_{\underline{a}\underline{b}}\psi_{(\alpha)}(x) &= \frac{1}{4}(\sigma_{\underline{a}\underline{b}})_{(\alpha)}^{(\beta)}\psi_{(\beta)}(x) \quad , \\
\mathcal{M}_{\underline{a}\underline{b}}A_{\underline{c}}(x) &= (\eta_{\underline{a}\underline{c}}\delta_{\underline{b}}^{\underline{d}} - \eta_{\underline{b}\underline{c}}\delta_{\underline{a}}^{\underline{d}})A_{\underline{d}}(x) \quad , \\
&\cdot \\
&\cdot \\
&\cdot
\end{aligned} \tag{9}$$

The basic ideas of Salam-Strathdee superspace<sup>1</sup> is that the geometrical notion of points should be replaced by “superpoints” and the 4D Minkowskian spacetime manifold is replaced by “superspace.”

$$x^{\underline{m}} \rightarrow z^{\underline{m}} \equiv \left( \theta^{(\mu)i}, x^{\underline{m}} \right) \quad , \quad i = 1, \dots, N \quad . \tag{10}$$

The quantity  $\theta^{(\mu)i}$  corresponds to  $4N$  Grassmann numbers (or anticommuting classical numbers ACN's) where the  $(\mu)$  index indicates that these carry a spinorial representation of the Lorentz group. If  $i$  takes on a single value, we call this “4D,  $N = 1$  superspace.” We want to discuss field theories in 4D,  $N = 1$  superspace. In a sense, Minkowski space lies on the boundary of superspace. Any physical consequences of superspace are only detectable for us on this boundary.

## 1.2 Grassmann Numerology

As observed above, the extra coordinates of superspace are not ordinary numbers. This distinguishes superspace from similar constructions such as Kaluza-Klein models. By definition a Grassmann number  $\theta$  is a non-trivial solution to the equation

$$\theta^2 = 0 \quad . \tag{11}$$

This should be thought of as an analogy to the fundamental definition of complex numbers

$$x^2 = -1 \quad . \tag{12}$$

Since we possess four of these (i.e.  $\theta^{(\mu)}$ ,  $(\mu) = (1), (2), (3), (4)$ ) we define also

$$\theta^{(\alpha)}\theta^{(\beta)} = -\theta^{(\beta)}\theta^{(\alpha)} \quad , \tag{13}$$

and furthermore may regard these as real quantities for the sake of simplicity

$$\theta^{(\alpha)} = \left[ \theta^{(\alpha)} \right]^* \quad . \tag{14}$$

We will continue the discussion of complex conjugation in superspace later. There is a certain subtlety which occurs and is the source of numbers of (from my viewpoint) ‘strange’ conventions in the literature about superspace and superfields.

### 1.3 A Notational Interlude

In order to calculate most efficiently in 4D,  $N = 1$  superspace, it is convenient to introduce a notation that is especially adapted to this purpose. The two-component notation of the van der Waerden formalism is the best. We begin by rewriting  $x^{\underline{m}}$  as a  $2 \times 2$  hermitian matrix

$$x^{\underline{m}} = \begin{pmatrix} x^0 + x^3 & x^1 - ix^2 \\ x^1 + ix^2 & x^0 - x^3 \end{pmatrix} \equiv x^{\mu\dot{\mu}} . \quad (15)$$

This permits us to regard the index  $\underline{m}$  as either the components of a 4-vector or as a pair of undotted and dotted indices as in the equation above. We will liberally make use of this convention in all subsequent discussions. So the interesting point is instead of thinking of  $x^{\underline{m}}$  as a 4-component vector, we may also regard it as a  $2 \times 2$  hermitian matrix. (Note  $\det(x^{\mu\dot{\mu}}) = -x^{\underline{m}}\eta_{\underline{m}\underline{n}}x^{\underline{n}}$ ).

Usually one first sees spinors expressed in terms of 4-component entities. This is mostly due to our familiarity with the Dirac equation and its use of four component entities to describe spinors. In all of our discussion up to this point, in the back of our minds we had equations like

$$\psi_{(\alpha)}(x) = \begin{pmatrix} \psi_{(1)}(x) \\ \psi_{(2)}(x) \\ \psi_{(3)}(x) \\ \psi_{(4)}(x) \end{pmatrix} , \quad (16)$$

which may be multiplied by  $4 \times 4$  “Dirac gamma” matrices  $(\gamma_{\underline{a}})$ . These latter quantities satisfy the usual algebraic equation

$$\left(\gamma_{\underline{a}}\right)_{(\alpha)}^{(\beta)} \left(\gamma_{\underline{b}}\right)_{(\beta)}^{(\gamma)} + \left(\gamma_{\underline{b}}\right)_{(\alpha)}^{(\beta)} \left(\gamma_{\underline{a}}\right)_{(\beta)}^{(\gamma)} = 2\eta_{\underline{a}\underline{b}} \left(\mathbf{I}\right)_{(\alpha)}^{(\gamma)} , \quad (17)$$

where  $(\mathbf{I})$  denotes the identity matrix.

The “chiral projection matrices”  $P_{\pm}$  may be defined by first noting (suppressing  $(\alpha)$ -type indices)

$$\gamma^5 \equiv i \frac{1}{4} \epsilon^{\underline{a}\underline{b}\underline{c}\underline{d}} \gamma_{\underline{a}} \gamma_{\underline{b}} \gamma_{\underline{c}} \gamma_{\underline{d}} , \quad (18)$$

which implies  $(\gamma^5)^2 = \text{I}$ , so that

$$P_{\pm} \equiv \frac{1}{2} \left( \text{I} \pm \gamma^5 \right) , \quad (19)$$

satisfy the equations

$$\text{I} = P_+ + P_- , \quad P_{\pm} P_{\mp} = 0 , \quad (P_{\pm})^2 = P_{\pm} . \quad (20)$$

All of this implies that there exist a basis where

$$P_+ = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} , \quad P_- = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} . \quad (21)$$

When a four component spinor is multiplied by these we find

$$\left( P_+ \right)_{(\alpha)}^{(\beta)} \psi_{(\beta)}(x) = \begin{pmatrix} \psi_{(1)}(x) \\ \psi_{(2)}(x) \\ 0 \\ 0 \end{pmatrix} \equiv \psi_{\alpha}(x) , \quad (22)$$

$$\left( P_- \right)_{(\alpha)}^{(\beta)} \psi_{(\beta)}(x) = \begin{pmatrix} 0 \\ 0 \\ \psi_{(3)}(x) \\ \psi_{(4)}(x) \end{pmatrix} \equiv \chi_{\alpha}(x) . \quad (23)$$

The spinors  $\psi_{\alpha}(x)$  and  $\chi_{\alpha}(x)$  are independent with each possessing only two components. In our notation, such spinor indices are described by Greek letters without the accompanying parenthesis marks. Note we use “dotted” and “undotted” indices to distinguish the chirality

$$\left( \gamma^5 P_+ \right)_{(\alpha)}^{(\beta)} \psi_{(\beta)}(x) = + \psi_{(\alpha)}(x) , \quad \left( \gamma^5 P_- \right)_{(\alpha)}^{(\beta)} \psi_{(\beta)}(x) = - \psi_{(\alpha)}(x) . \quad (24)$$

So  $\psi_{\alpha}(x)$  is the positive chiral (“right-handed”) part of the four component spinor  $\psi_{(\alpha)}(x)$  and  $\chi_{\alpha}(x)$  is the negative chiral (“left-handed”) part.

Finally we may use  $P_+$  and  $P_-$  in one more way. We define  $2 \times 2$  Pauli matrices  $\sigma_{\underline{a}}$  via

$$\left( P_+ \gamma_{\underline{a}} P_- \right)_{(\alpha)}^{(\beta)} = \begin{pmatrix} 0 & (\sigma_{\underline{a}})_{\alpha}^{\dot{\beta}} \\ 0 & 0 \end{pmatrix} , \quad \left( P_- \gamma_{\underline{a}} P_+ \right)_{(\alpha)}^{(\beta)} = \begin{pmatrix} 0 & 0 \\ (\sigma_{\underline{a}})^{\beta}_{\dot{\alpha}} & 0 \end{pmatrix} , \quad (25)$$



and as well observe  $P_{\pm}\gamma_{\underline{a}}P_{\pm}=0$ . The Pauli matrices permit us to write

$$\begin{aligned} x^0(\sigma_0)_{\alpha}^{\dot{\alpha}} + x^1(\sigma_1)_{\alpha}^{\dot{\alpha}} + x^2(\sigma_2)_{\alpha}^{\dot{\alpha}} + x^3(\sigma_3)_{\alpha}^{\dot{\alpha}} &= x_{\alpha}^{\dot{\alpha}} \quad , \\ (\sigma_{\underline{a}})_{\alpha}^{\dot{\beta}}(\sigma_{\underline{b}})^{\gamma}_{\dot{\beta}} + (\sigma_{\underline{b}})_{\alpha}^{\dot{\beta}}(\sigma_{\underline{a}})^{\gamma}_{\dot{\beta}} &= 2\eta_{\underline{ab}}\left(\mathbf{I}\right)_{\alpha}^{\gamma} \quad , \\ (\sigma_{\underline{a}})^{\beta}_{\dot{\alpha}}(\sigma_{\underline{b}})^{\dot{\gamma}}_{\beta} + (\sigma_{\underline{b}})^{\beta}_{\dot{\alpha}}(\sigma_{\underline{a}})^{\dot{\gamma}}_{\beta} &= 2\eta_{\underline{ab}}\left(\mathbf{I}\right)^{\dot{\gamma}}_{\dot{\alpha}} \quad , \end{aligned} \quad (26)$$

but as we will see these last two need never be used by a clever choice of conventions.

In ordinary 4D spacetime, four vector indices are raised and lowered with  $\eta_{ab}$ . What can we do in Superspace? It turns out that some unexpected results can be obtained via some more definitions.

$$\eta_{\underline{ab}} \equiv C_{\alpha\beta}C_{\dot{\alpha}\dot{\beta}} \quad , \quad \eta^{\underline{ab}} \equiv C^{\alpha\beta}C^{\dot{\alpha}\dot{\beta}} \quad , \quad (27)$$

$$C_{\alpha\beta} = i \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad , \quad C^{\alpha\beta} = -i \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad , \quad (28)$$

$$C_{\dot{\alpha}\dot{\beta}} = i \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad , \quad C^{\dot{\alpha}\dot{\beta}} = -i \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad . \quad (29)$$

All the usual properties of  $\eta_{\underline{ab}}$  can be seen to be true using this representation, (i.e.  $\eta_{\underline{ab}}\eta^{\underline{ab}}=4$ ,  $\eta_{\underline{ab}}=\eta_{\underline{ba}}$ , etc.) The  $C$ 's can also be used to raise and lower spinor indices

$$\theta_{\alpha} = \theta^{\beta}C_{\beta\alpha} \quad , \quad \theta^{\alpha} = C^{\alpha\beta}\theta_{\beta} \quad , \quad \bar{\theta}_{\dot{\alpha}} = \bar{\theta}^{\dot{\beta}}C_{\dot{\beta}\dot{\alpha}} \quad , \quad \bar{\theta}^{\dot{\alpha}} = C^{\dot{\alpha}\dot{\beta}}\bar{\theta}_{\dot{\beta}} \quad . \quad (30)$$

Notice that since  $C_{\alpha\beta}=-C_{\beta\alpha}$  we must exercise some care, i.e.

$$\theta^{\alpha}\theta_{\alpha} \neq \theta_{\alpha}\theta^{\alpha} \quad , \quad \bar{\theta}^{\dot{\alpha}}\bar{\theta}_{\dot{\alpha}} \neq \bar{\theta}_{\dot{\alpha}}\bar{\theta}^{\dot{\alpha}} \quad . \quad (31)$$

As well these are useful to define the ‘‘magnitude’’ of the Grassmann numbers

$$\theta^2 \equiv -\frac{1}{2}\theta^{\alpha}C_{\alpha\beta}\theta^{\beta} \quad , \quad \bar{\theta}^2 \equiv -\frac{1}{2}\bar{\theta}^{\dot{\alpha}}C_{\dot{\alpha}\dot{\beta}}\bar{\theta}^{\dot{\beta}} \quad . \quad (32)$$

There is one additional invariant four tensor that is very useful in  $\mathcal{M}_4$ , the Levi-Civita tensor. This too has a representation in terms of the spinor  $C$ -metrics.

$$\begin{aligned} \epsilon_{\underline{abcd}} &\equiv i\frac{1}{2}\left[C_{\alpha\beta}C_{\gamma\delta}C_{\dot{\alpha}(\gamma}C_{\dot{\delta})\beta} - C_{\dot{\alpha}\dot{\beta}}C_{\dot{\gamma}\dot{\delta}}C_{\alpha(\gamma}C_{\delta)\beta}\right] \quad , \\ \epsilon^{\underline{abcd}} &\equiv i\frac{1}{2}\left[C^{\alpha\beta}C^{\gamma\delta}C^{\dot{\alpha}(\gamma}C^{\dot{\delta})\beta} - C^{\dot{\alpha}\dot{\beta}}C^{\dot{\gamma}\dot{\delta}}C^{\alpha(\gamma}C^{\delta)\beta}\right] \quad . \end{aligned} \quad (33)$$

Once again it is easy to verify familiar properties (i.e.  $\epsilon_{\underline{b}\underline{a}\underline{c}\underline{d}} = -\epsilon_{\underline{a}\underline{b}\underline{c}\underline{d}}$ , etc.  $\eta^{\underline{a}\underline{b}}\epsilon_{\underline{a}\underline{b}\underline{c}\underline{d}} = 0$ ). Above the notation  $(\ )$  acting on a pair of indices is given by

$$\psi_{(\alpha}\bar{\chi}_{\beta)} \equiv \psi_{\alpha}\chi_{\beta} + \psi_{\beta}\bar{\chi}_{\alpha} \quad . \quad (34)$$

We shall also use the notation  $[\ ]$  acting on a pair of indices

$$\begin{aligned} \psi_{[\alpha}\bar{\chi}_{\beta]} &\equiv \psi_{\alpha}\bar{\chi}_{\beta} - \psi_{\beta}\bar{\chi}_{\alpha} = -C_{\alpha\beta}\psi^{\gamma}\bar{\chi}_{\gamma} \quad , \\ \bar{\psi}_{[\dot{\alpha}}\chi_{\dot{\beta}}] &\equiv \bar{\psi}_{\dot{\alpha}}\chi_{\dot{\beta}} - \bar{\psi}_{\dot{\beta}}\chi_{\dot{\alpha}} = -C_{\dot{\alpha}\dot{\beta}}\bar{\psi}^{\dot{\gamma}}\chi_{\dot{\gamma}} \quad . \end{aligned} \quad (35)$$

We end this notational interlude by observing that as we regard the  $2 \times 2$  matrix in (15) as the definition of the ‘bosonic’ component of a superpoint, derivatives of fields may be calculated with respect to this  $2 \times 2$  matrix. In particular, when we write the symbol  $P_{\underline{m}}$ , this may be regarded as an instruction to differentiate with respect to the variable in (15). Adhering to this convention, no Pauli nor Dirac matrices ever appear in our formulation of supersymmetrical theories. As consequences of this we find

$$\begin{aligned} \partial_{\mu\dot{\mu}}\partial^{\nu\dot{\mu}} &\equiv \delta_{\mu}^{\nu}\square \rightarrow \partial_{\mu\dot{\mu}}\partial_{\nu}^{\dot{\mu}} = C_{\mu\nu}\square \quad , \quad \square = \frac{1}{2}\partial^{\underline{m}}\partial_{\underline{m}} \quad , \\ \partial_{\mu\dot{\mu}}\partial^{\mu\dot{\nu}} &\equiv \delta_{\mu}^{\nu}\square \rightarrow \partial_{\mu\dot{\mu}}\partial^{\mu\dot{\nu}} = C_{\mu\nu}\square \quad . \end{aligned} \quad (36)$$

Since no Pauli or Dirac matrices ever appear, it is never necessary to calculate any traces thereof.

This interlude has been rather lengthy and the notation to be used in the subsequent discussion has now been described. It may seem as though this set of conventions is arbitrary and unwieldy. However, I encourage the reader to actually use it. In fact for supersymmetrical theories it is the most efficient notation I have seen.

#### 1.4 Super Translation Generators

Just as Minkowski space is actually an equivalence class, so too is superspace. In the following we are going to learn the form of the generalization of the Poincaré group. We consider our superpoints of 4D,  $N = 1$  superspace to be represented by

$$z^{\underline{M}} \equiv \left( \theta^{\mu}, \bar{\theta}^{\dot{\mu}}, x^{\underline{m}} \right) \quad , \quad (37)$$

using 2-component spinor Grassmann coordinates. It is simplest to first describe the super-translations as differential operators similar to those in (5).

$$\begin{aligned}
P_{\underline{m}} &= -i \partial_{\underline{m}} = -i \frac{\partial}{\partial x^{\mu\mu}} , \\
Q_{\mu} &= i \left[ \frac{\partial}{\partial \theta^{\mu}} - i \frac{1}{2} \bar{\theta}^{\dot{\mu}} \partial_{\underline{m}} \right] = i \left[ \partial_{\mu} - i \frac{1}{2} \bar{\theta}^{\dot{\mu}} \partial_{\underline{m}} \right] , \\
\bar{Q}_{\dot{\mu}} &= \left[ \frac{\partial}{\partial \bar{\theta}^{\dot{\mu}}} - i \frac{1}{2} \theta^{\mu} \partial_{\underline{m}} \right] = i \left[ \bar{\partial}_{\dot{\mu}} - i \frac{1}{2} \theta^{\mu} \partial_{\underline{m}} \right] .
\end{aligned} \tag{38}$$

It is a simple matter to show that the follow algebra is satisfied,

$$\begin{aligned}
[P_{\underline{r}}, Q_{\nu}] &= 0 , \quad [P_{\underline{r}}, \bar{Q}_{\dot{\nu}}] = 0 , \quad [P_{\underline{r}}, P_{\underline{s}}] = 0 , \\
\{Q_{\mu}, Q_{\nu}\} &= 0 , \quad \{\bar{Q}_{\dot{\mu}}, \bar{Q}_{\dot{\nu}}\} = 0 , \quad \{Q_{\mu}, \bar{Q}_{\dot{\mu}}\} = i P_{\underline{m}} .
\end{aligned} \tag{39}$$

Note the anti-commutators, this is a “super” Lie algebra. (Also note absence of gamma matrices, i.e. simplicity.)

The ordinary translation generators “create” motions

$$i(k_0)^{\underline{m}} P_{\underline{m}} x^{\underline{n}} \rightarrow \exp[i(k_0)^{\underline{m}} P_{\underline{m}}] x^{\underline{n}} = x^{\underline{n}} + (k_0)^{\underline{n}} . \tag{40}$$

So it is revealing to study the motions of superpoints generated by the operators in (38).

$$\begin{aligned}
i(k_0)^{\underline{m}} P_{\underline{m}} z^{\underline{N}} &= \left( 0, 0, (k_0)^{\underline{m}} \right) , \\
i \epsilon_0^{\mu} Q_{\mu} z^{\underline{N}} &= \left( -\epsilon_0^{\mu}, 0, i \frac{1}{2} \epsilon_0^{\mu} \bar{\theta}^{\dot{\mu}} \right) , \\
i \bar{\epsilon}_0^{\dot{\mu}} \bar{Q}_{\dot{\mu}} z^{\underline{N}} &= \left( 0, -\bar{\epsilon}_0^{\dot{\mu}}, i \frac{1}{2} \bar{\epsilon}_0^{\dot{\mu}} \theta^{\mu} \right) ,
\end{aligned} \tag{41}$$

where we have introduced Grassmann parameters  $\epsilon_0^{\mu}$  and  $\bar{\epsilon}_0^{\dot{\mu}}$  so that we can form a ‘super-translation’ parameter  $(k_0)^{\underline{M}} = (\epsilon_0^{\mu}, \bar{\epsilon}_0^{\dot{\mu}}, (k_0)^{\underline{m}})$ . Also rather obviously it is possible to combine the translations in a similar manner,  $P_{\underline{M}} = (Q_{\mu}, \bar{Q}_{\dot{\mu}}, P_{\underline{m}})$ . Using a more concise notation, we have  $\delta z^{\underline{N}} = i(k_0)^{\underline{M}} P_{\underline{M}} z^{\underline{N}}$ , so that

$$\delta \theta^{\mu} = -\epsilon_0^{\mu} , \quad \delta \bar{\theta}^{\dot{\mu}} = -\bar{\epsilon}_0^{\dot{\mu}} , \quad \delta x^{\underline{m}} = i \frac{1}{2} ( \epsilon_0^{\mu} \bar{\theta}^{\dot{\mu}} + \bar{\epsilon}_0^{\dot{\mu}} \theta^{\mu} ) . \tag{42}$$

### 1.5 Super Lorentz Generators

It is also possible to generalize the orbital angular momentum generator to a superspace orbital angular momentum generator  $L_{\underline{M}\underline{N}}$ . The only non-vanishing components of this tensor are  $L_{(\underline{\mu}\underline{\nu})}$ ,  $L_{(\dot{\underline{\mu}}\dot{\underline{\nu}})}$  and  $L_{\underline{m}\underline{n}}$ . Since the Grassmann coordinates are relativistic spinors, it is natural that they should appear in the appropriate generalization,

$$\begin{aligned} L_{\underline{m}\underline{n}} &\equiv C_{\underline{\mu}\underline{\nu}} L_{(\dot{\underline{\mu}}\dot{\underline{\nu}})} + C_{\dot{\underline{\mu}}\dot{\underline{\nu}}} L_{(\underline{\mu}\underline{\nu})} \quad , \\ L_{(\underline{\mu}\underline{\nu})} &\equiv -i \frac{1}{2} [ x_{(\underline{\mu}} \dot{\partial}_{\underline{\nu})} + \theta_{(\underline{\mu}} \partial_{\underline{\nu})} ] \quad , \\ L_{(\dot{\underline{\mu}}\dot{\underline{\nu}})} &\equiv -i \frac{1}{2} [ x^{\underline{\nu}}_{(\dot{\underline{\mu}}} \partial_{\dot{\underline{\nu}})} + \bar{\theta}_{(\dot{\underline{\mu}}} \bar{\partial}_{\dot{\underline{\nu}})} ] \quad . \end{aligned} \quad (43)$$

This first equation is consistent since  $\underline{m} = \underline{\mu}\dot{\underline{\mu}}$  and  $\underline{n} = \underline{\nu}\dot{\underline{\nu}}$ . The generator  $L_{\underline{m}\underline{n}}$  does exactly what we expect. Namely it implies that  $x^{\underline{m}}$  transforms as a Lorentz 4-vector and  $\theta^{\underline{\mu}}$ ,  $\bar{\theta}^{\dot{\underline{\mu}}}$  transform as Lorentz spinors.

The set of operators  $\{\mathcal{A}\} = \{P_{\underline{M}}, L_{\underline{m}\underline{n}}\}$  forms a super-Lie algebra. The basic point is that we may define

$$(-1)^P \equiv 1 \quad , \quad (-1)^L \equiv 1 \quad , \quad (-1)^Q \equiv -1 \quad , \quad (-1)^{\bar{Q}} \equiv -1 \quad , \quad (44)$$

and a generalized or graded commutator through

$$[ \mathcal{A}_1 , \mathcal{A}_2 ] \equiv \mathcal{A}_1 \mathcal{A}_2 - (-1)^{\mathcal{A}_1 \mathcal{A}_2} \mathcal{A}_2 \mathcal{A}_1 \quad . \quad (45)$$

Under the operation of this generalized commutator  $\{\mathcal{A}\}$  is a closed set. In addition, there is a generalized Jacobi identity defined by

$$\begin{aligned} (-1)^{\mathcal{A}_1 \mathcal{A}_3} [ [ \mathcal{A}_1 , \mathcal{A}_2 ] , \mathcal{A}_3 ] + (-1)^{\mathcal{A}_2 \mathcal{A}_1} [ [ \mathcal{A}_2 , \mathcal{A}_3 ] , \mathcal{A}_1 ] + \\ (-1)^{\mathcal{A}_3 \mathcal{A}_2} [ [ \mathcal{A}_3 , \mathcal{A}_1 ] , \mathcal{A}_2 ] = 0 \quad , \end{aligned} \quad (46)$$

which the generators satisfy.

### 1.6 Superspace Conjugation

Note that if we consider (42) solely for the temporal component of  $x^{\underline{m}}$  (thus breaking Lorentz covariance) we have  $\delta x^{\underline{0}} \propto i (\sigma^{\underline{0}})_{\underline{\mu}\dot{\underline{\mu}}} (\epsilon_0^{\underline{\mu}} \dot{\bar{\theta}}^{\dot{\underline{\mu}}} + \bar{\epsilon}_0^{\dot{\underline{\mu}}} \dot{\theta}^{\underline{\mu}})$ . At first this seems extremely problematical since  $x^{\underline{0}}$  is a real quantity, the factor in the parenthesis is real using standard definitions and  $(\sigma^{\underline{0}})_{\underline{\mu}\dot{\underline{\mu}}}$  is real. Yet this equation suggests that the variation of  $x^{\underline{0}}$  is *purely* imaginary (and nilpotent) even if we use a real spinor basis.

But the variation of  $(ct)$  should be real! We can make this so if the meaning of  $*$  is modified. One way to do this in superspace is to assign the following definition of how complex conjugation is applied

$$\begin{aligned} * &\equiv \mathcal{C} \otimes \mathcal{O}_I \quad , \\ \mathcal{C} &\equiv \text{replace } i \text{ by } -i \text{ and vice - versa} \quad , \\ \mathcal{O}_I &\equiv \text{invert the order of all monomials in all factors} \quad . \end{aligned} \tag{47}$$

To see how this definition of the  $*$  operation solves the problem, consider the brief calculation below. For the purposes of this illustration, we can consider a truly one dimensional world in which spinors are real one-component entities.

$$\begin{aligned} \mathcal{C} \otimes \mathcal{O}_I : [i\frac{1}{2}\epsilon\theta] &= \theta^* \epsilon^* \frac{1}{2}(-i) \\ &= -i\frac{1}{2}\theta\epsilon \\ &= i\frac{1}{2}\epsilon\theta \\ [i\frac{1}{2}\epsilon\theta]^* &= [i\frac{1}{2}\epsilon\theta] \quad . \end{aligned} \tag{48}$$

So under superspace conjugation the variation of the temporal component (and all others) of  $x^{\underline{m}}$  is real.

The definition of the  $*$  operator (which I first introduced in my Ph.D. thesis<sup>2)</sup> above may seem very capricious. It is not. In large numbers of places in the literature on superspace, this is the cause of the appearance of seemingly ‘inexplicable’ signs. Let us continue to study more implications of this rule. Although we had not mentioned it before, the spinor in (16) may be subjected to a reality condition,  $\psi_{(\alpha)} = [\psi_{(\alpha)}]^*$ . However this condition is inconsistent with the basis in which we have given the  $\gamma^5$ -matrix in (18-23). Thus if we wish to use this basis, the spinors in (22,23) must be complex. The same reasoning applies to the spinorial components of superspace. If we use 2-component spinor coordinates for superspace, they must be complex coordinates.

The fact that we use complex Grassmann coordinates is consistent with the definitions that we have introduced for raising and lowering their indices with the  $C$ -tensors since the latter include factors of  $i$  in their definitions. Thus we have (also recall we only use superspace conjugation)

$$\begin{aligned} (\theta^\alpha)^* &= \bar{\theta}^{\dot{\alpha}} \text{ but } (\theta_\alpha)^* = -\bar{\theta}_{\dot{\alpha}} \quad , \\ (\bar{\theta}^{\dot{\alpha}})^* &= \theta^\alpha \text{ but } (\bar{\theta}_{\dot{\alpha}})^* = -\theta_\alpha \quad . \end{aligned} \tag{49}$$

Another consequence follows from

$$\begin{aligned}
(\theta^\alpha \theta_\alpha)^* &= (\theta_\alpha)^* (\theta^\alpha)^* \quad \text{s-conjugation} \\
&= -\bar{\theta}_\alpha \cdot \bar{\theta}^\alpha \\
&= \bar{\theta}^\alpha \bar{\theta}_\alpha \quad \text{Grassmann property} .
\end{aligned} \tag{50}$$

So that the quantity  $\theta^\alpha \theta_\alpha + \bar{\theta}^\alpha \bar{\theta}_\alpha$  is real for us.

Also since  $\partial_\alpha \theta^\beta = \delta_\alpha^\beta$ , it follows that  $(\partial_\alpha)^* = -\bar{\partial}_\alpha$  (and of course  $\bar{\partial}_\alpha \bar{\theta}^\beta = \delta_\alpha^\beta$ , implies  $(\bar{\partial}_\alpha)^* = -\partial_\alpha$ ). This is seen through the calculation below.

$$\begin{aligned}
(\partial_\alpha \theta^\beta)^* &= (\theta^\beta)^* (\bar{\partial}_\alpha)^* \quad \text{s-conjugation} \\
&= \bar{\theta}^\beta (-\bar{\partial}_\alpha) \\
&= \bar{\partial}_\alpha \bar{\theta}^\beta \quad \text{Grassmann property} .
\end{aligned} \tag{51}$$

The general rule is if a superspace tensor is “real” it transforms like a product of superspace monomial coordinates under the action of superspace conjugation. We apply the superspace conjugation rule followed by restoring the order of the indices as they appear on the original tensor. This is simplest demonstrated by some examples.

$$\begin{aligned}
x^{\underline{m}} \sim \theta^\mu \bar{\theta}^\mu ; \quad (\theta^\mu \bar{\theta}^\mu)^* &= (\bar{\theta}^\mu)^* (\theta^\mu)^* \\
&= \theta^\mu \bar{\theta}^\mu \rightarrow (x^{\underline{m}})^* = x^{\underline{m}} ,
\end{aligned} \tag{52}$$

$$\begin{aligned}
x_{\underline{m}} \sim \theta_\mu \bar{\theta}_\mu ; \quad (\theta_\mu \bar{\theta}_\mu)^* &= (\bar{\theta}_\mu)^* (\theta_\mu)^* \\
&= (-\theta_\mu) (-\bar{\theta}_\mu) \rightarrow (x_{\underline{m}})^* = x_{\underline{m}} .
\end{aligned} \tag{53}$$

These examples may be considered too simple, so we finish with a multi-indexed more complicated example,

$$\begin{aligned}
T_{\alpha\beta\gamma\underline{d}}^\epsilon &\sim \theta_\alpha \theta_\beta \bar{\theta}_\gamma x_{\underline{d}} \theta^\epsilon \\
(\theta_\alpha \theta_\beta \bar{\theta}_\gamma x_{\underline{d}} \theta^\epsilon)^* &= (\theta^\epsilon)^* (x_{\underline{d}})^* (\bar{\theta}_\gamma)^* (\theta_\beta)^* (\theta_\alpha)^* \\
&= \bar{\theta}^\epsilon x_{\underline{d}} (-\theta_\gamma) (-\bar{\theta}_\beta) (-\bar{\theta}_\alpha) \\
&= -\bar{\theta}_\alpha \bar{\theta}_\beta \theta_\gamma x_{\underline{d}} \bar{\theta}^\epsilon .
\end{aligned} \tag{54}$$

Thus if  $T$  is a “real” superspace tensor it possesses the property

$$(T_{\alpha\beta\dot{\gamma}\underline{d}}^{\dot{\epsilon}})^* = -T_{\alpha\dot{\beta}\gamma\underline{d}}^{\dot{\epsilon}} \quad . \quad (55)$$

### 1.7 Superfields

In 4D spacetime, fields may be defined as the fundamental dynamical entities with the scalar field  $\phi(x)$  as the simplest example. How do such entities appear in superspace?

Definition: A superfield is a function of  $z^{\underline{m}}$  that is analytic in  $\theta^\mu$  and  $\bar{\theta}^{\dot{\mu}}$ .

A simple example of a superfield is provided by a real scalar superfield  $F(z)$  which may be expanded as

$$\begin{aligned} F(z) = & C(x) + \theta^\alpha \chi_\alpha(x) + \bar{\theta}^{\dot{\alpha}} \bar{\chi}_{\dot{\alpha}}(x) - \theta^2 M(x) - \bar{\theta}^2 \bar{M}(x) \\ & + \theta^\alpha \bar{\theta}^{\dot{\alpha}} A_{\underline{a}}(x) - \bar{\theta}^2 \theta^\alpha \lambda_\alpha(x) - \theta^2 \bar{\theta}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}}(x) + \theta^2 \bar{\theta}^2 d(x) \quad . \end{aligned} \quad (56)$$

The expansion terminates due to the Grassmann nature of the  $\theta$ 's. The “component fields”  $C(x)$ ,  $\chi_\alpha(x)$ ,  $M(x)$ ,  $A_{\underline{a}}(x)$ ,  $\lambda_\alpha(x)$  and  $d(x)$  are ordinary fields. All the bosonic fields are real ones (see later discussion). How do the motions which generate an equivalence class of superspace affect these ordinary fields? The answer to this is found by applying the differential operators which represent the generators to the expansion of the superfield. Thus in terms of the superfield we write,

$$\delta_Q F(z) = i \left( \epsilon^\mu Q_\mu + \bar{\epsilon}^{\dot{\mu}} \bar{Q}_{\dot{\mu}} \right) F(z) \quad . \quad (57)$$

where we have “dropped” the zero subscript on the parameters. In terms of the fields in the expansion, this leads to

$$\begin{aligned} \delta_Q C &= - ( \epsilon^\alpha \chi_\alpha + \bar{\epsilon}^{\dot{\alpha}} \bar{\chi}_{\dot{\alpha}} ) \quad , \\ \delta_Q \chi_\alpha &= \epsilon_\alpha M - \bar{\epsilon}^{\dot{\alpha}} ( i \frac{1}{2} \partial_{\underline{a}} C + A_{\underline{a}} ) \quad , \\ \delta_Q M &= \bar{\epsilon}^{\dot{\alpha}} ( \bar{\lambda}_{\dot{\alpha}} - i \frac{1}{2} \partial_{\underline{a}} \bar{\chi}^{\dot{\alpha}} ) \quad , \\ \delta_Q A_{\underline{a}} &= \epsilon^\beta ( C_{\beta\alpha} \bar{\lambda}_{\dot{\alpha}} + i \frac{1}{2} \partial_{\beta\dot{\alpha}} \chi_\alpha ) + \bar{\epsilon}^{\dot{\beta}} ( C_{\dot{\beta}\alpha} \lambda_\alpha + i \frac{1}{2} \partial_{\alpha\dot{\beta}} \bar{\chi}_{\dot{\alpha}} ) \quad , \\ \delta_Q \lambda_\alpha &= \epsilon^\beta ( C_{\beta\alpha} d + i \frac{1}{2} \partial_{\beta\dot{\alpha}} A^{\dot{\beta}} ) - \bar{\epsilon}^{\dot{\alpha}} \partial_{\underline{a}} \bar{M} \quad , \\ \delta_Q d &= i \frac{1}{2} \left[ \partial_{\underline{a}} ( \epsilon^\alpha \lambda^\alpha + \bar{\epsilon}^{\dot{\alpha}} \bar{\lambda}^{\dot{\alpha}} ) \right] \quad . \end{aligned} \quad (58)$$

This completes the discussion for the translational part of the super Poincaré group, but there are still to be discussed the effects of the pure Lorentz generators (43). To do this consistently, we return to our discussion (6) of the Poincaré group. There we saw the parameter of the pure Lorentz rotations were given by  $(K_0)_{\underline{a}\underline{b}}$ . Similar to the result of (43) we may introduce new parameters  $K_{\alpha\beta}$  and  $K_{\dot{\alpha}\dot{\beta}}$  via

$$(K_0)_{\underline{a}\underline{b}} = \frac{1}{2} [ C_{\dot{\alpha}\dot{\beta}} K_{\alpha\beta} + C_{\alpha\beta} K_{\dot{\alpha}\dot{\beta}} ] \quad . \quad (59)$$

This corresponds to a decomposition of the Lorentz parameters into “right-handed” (or “self-dual”) and “left-handed” (or “anti-self-dual”) parts. In the case of the translations, we added new Grassmann parameters ( $\epsilon$  and  $\bar{\epsilon}$ ) so that the fermionic translations could be implemented. We do something very different for the Lorentz rotations. Even though there exist the supertensor  $L_{\underline{M}\underline{N}}$ , according to (59) not all of its components are independent. Thus, it is unnecessary and redundant to attempt to introduce new parameters for its fermionic components. Instead the motions of the superspace orbital angular momentum generator are derived from

$$\delta_L F(z) = i \frac{1}{2} (K_0)_{\underline{a}\underline{b}} L_{\underline{b}}^{\underline{a}} F(z) = i \frac{1}{2} [ K_{\alpha}^{\beta} L_{\beta}^{\alpha} + K_{\dot{\alpha}}^{\dot{\beta}} L_{\dot{\beta}}^{\dot{\alpha}} ] F(z) \quad . \quad (60)$$

It is an easy exercise to show that this induces at the component level the interesting result that for the components  $C$ ,  $M$  and  $d$  transform as scalars,  $\lambda_{\alpha}$  and  $\chi_{\alpha}$  transform as spinors and  $A_{\underline{a}}$  transforms as a vector.

So the superfield  $F(z)$  contains component fields of different spins even though the superfield itself carries no ‘external’ spinor or vector indices. However, this does *not* preclude us from considering superfields that do. In fact, such superfields do exist and are important in numbers of applications. This leads to the notion that the abstract spin angular momentum operator  $S_{\underline{a}\underline{b}}$  continues to exist in superspace (which it does). It is usually referred to as the “superspin operator” in this context.

The real SF (superfield) is not the simplest, most fundamental 4D supersymmetry representation. This distinction belongs to the chiral SF  $\Phi(z)$ . The chiral superfield is foremost a complex superfield whose expansion takes the form

$$\begin{aligned} \Phi(z) = & A(x) + \theta^{\alpha} \psi_{\alpha}(x) - \theta^2 F(x) + i \frac{1}{2} \theta^{\alpha} \bar{\theta}^{\dot{\alpha}} \partial_{\underline{a}} A(x) \\ & + i \frac{1}{2} \theta^2 \bar{\theta}^{\dot{\alpha}} \partial_{\underline{a}} \psi^{\alpha}(x) + \frac{1}{4} \theta^2 \bar{\theta}^2 \square A(x) \quad . \end{aligned} \quad (61)$$

Here the bosonic fields  $A(x)$  and  $F(x)$  are complex. It should be clear that  $\Phi$  is a type of ‘restricted’ superfield since some of the  $\theta$ -expansion terms which



appear in (56) are missing from the expansion in (61). The changes induced among the fields by the fermionic translation are found from

$$\begin{aligned}\delta_Q \Phi(z) &= i \left( \epsilon^\mu Q_\mu + \bar{\epsilon}^\mu \bar{Q}_\mu \right) \Phi(z) \quad , \\ \rightarrow \quad \delta_Q A &= - \epsilon^\alpha \psi_\alpha \quad , \quad \delta_Q \psi_\alpha = - i \bar{\epsilon}^\alpha \partial_{\underline{a}} A + \epsilon_\alpha F \quad , \\ \delta_Q F &= i \bar{\epsilon}^\alpha \partial_{\underline{a}} \psi^\alpha \quad .\end{aligned}\tag{62}$$

In order to relate the chiral superfield to the general complex superfield, it is convenient to define some new spinorial differential operators,

$$D_\mu \equiv \left[ \partial_\mu + i \frac{1}{2} \bar{\theta}^\mu \partial_{\underline{m}} \right] \quad , \quad \bar{D}_\mu = \left[ \bar{\partial}_\mu + i \frac{1}{2} \theta^\mu \partial_{\underline{m}} \right] \quad , \tag{63}$$

that are called the “supersymmetry covariant derivatives.” It can be seen that these are *not* linearly related to the super translation generators  $Q_\mu$  and  $\bar{Q}_\mu$ . In fact, these possess a very different graded commutator algebra,

$$\begin{aligned}\left[ P_\underline{r} , D_\nu \right] &= 0 \quad , \quad \left[ P_\underline{r} , \bar{D}_\nu \right] = 0 \quad \left[ D_\mu , D_\nu \right] = 0 \quad , \\ \left[ Q_\mu , D_\nu \right] &= 0 \quad , \quad \left[ Q_\mu , \bar{D}_\nu \right] = 0 \quad , \quad \left[ D_\mu , \bar{D}_\nu \right] = i \partial_{\underline{m}} \quad .\end{aligned}\tag{64}$$

We define a “super vector” gradient by

$$D_{\underline{A}} = \left( D_\alpha , \bar{D}_{\dot{\alpha}} , \partial_{\underline{a}} \right) \quad , \tag{65}$$

and this is the basic tool for constructing kinetic energy operators for superfields. The chiral superfield satisfies the differential equation  $\bar{D}_{\dot{\alpha}} \Phi = 0$ . It can also be noted that the  $*$  operation has the following realization on a complex scalar superfield  $H(z)$

$$\begin{aligned}D_\alpha H(z) &= \left[ \partial_\alpha + i \frac{1}{2} \bar{\theta}^\alpha \partial_{\underline{a}} \right] H \quad \rightarrow \\ \left[ D_\alpha H(z) \right]^* &= H^* \left[ \bar{D}_\alpha \right]^* = H^* \left[ [\bar{\partial}_\alpha]^* + [\bar{\partial}_{\underline{a}}]^* (\bar{\theta}^\alpha)^* \frac{1}{2} (-i) \right] \\ &= H^* \left[ - \bar{\partial}_\alpha - i \frac{1}{2} \bar{\partial}_{\underline{a}} (\theta^\alpha) \right] \\ &= - \left[ \bar{\partial}_\alpha + i \frac{1}{2} \theta^\alpha \partial_{\underline{a}} \right] H^* \quad .\end{aligned}\tag{66}$$

Since  $D_\alpha$  has a lower spinor index, we expect  $[\overleftarrow{D}_\alpha]^* = -\overleftarrow{D}_\alpha$  so we obtain,

$$\begin{aligned} H^* \left[ -\overleftarrow{D}_\alpha \right] &= - \left[ \bar{\partial}_\alpha + i\frac{1}{2} \theta^\alpha \partial_\alpha \right] H^* \quad , \quad \text{or} \\ \overleftarrow{D}_\alpha H^* &= \left[ \bar{\partial}_\alpha + i\frac{1}{2} \theta^\alpha \partial_\alpha \right] H^* \quad . \end{aligned} \quad (67)$$

One can verify that all of these operatorial steps make sense by explicitly calculating the rhs of the first line in (66), complex conjugate that resultant and then comparing to the explicit component results obtainable from the rhs of the last line in (67). It is often the case that in other conventions in order to get agreement among these two result, one must insert minus signs which apparently have no explanation.

### 1.8 Superfields; Spin-Statistics & Bose-Fermi Equality

In Minkowski space, bosons (B) are fields with even numbers of spinorial indices

$$B(x) \sim B^{\alpha_1 \dots \alpha_p \dot{\beta}_1 \dots \dot{\beta}_q} \quad , \quad p + q = 2n \quad , \quad (68)$$

and fermions are fields with an odd number of these indices

$$F(x) \sim F^{\alpha_1 \dots \alpha_p \dot{\beta}_1 \dots \dot{\beta}_q} \quad , \quad p + q = 2n + 1 \quad . \quad (69)$$

These two classes of fields obey the statistics rules (classically at least)

$$B_1 B_2 = B_2 B_1 \quad , \quad F_1 F_2 = -F_2 F_1 \quad , \quad B F = F B \quad . \quad (70)$$

If we assume that these also hold for superfields  $B(x) \rightarrow B(z)$  and  $F(x) \rightarrow F(z)$ , then we can derive the fact that a superfield necessarily includes fields of both statistics. Applying this to our prototype superfield  $F$  (56) we find the result

$$\begin{aligned} \{ C(x), M(x), A_\alpha(x), d(x) \} &\in \{ B \} \quad , \\ \{ \lambda_\alpha(x), \chi_\alpha(x) \} &\in \{ F \} \quad . \end{aligned} \quad (71)$$

If we replace  $F \rightarrow F^\alpha$  and demand that  $F_1^\alpha$  and  $F_2^\alpha$  obey fermi statistics then (with an obvious change in notation)

$$\begin{aligned} \{ C^\beta(x), M^\beta(x), A_\alpha^\beta(x), d^\beta(x) \} &\in \{ F \} \quad , \\ \{ \lambda_\alpha^\beta(x), \chi_\alpha^\beta(x) \} &\in \{ B \} \quad . \end{aligned} \quad (72)$$

In particular, we see that that Duffin-Kummer-Petiau fields  $\lambda_\alpha^\beta$  and  $\chi_\alpha^\beta$  fields appear. A similar result occurs if we replace  $F$  according to  $F \rightarrow F^\alpha$ .

The other obvious feature of 4D superfields is that they always introduce multiplets of particles of different spins but where the total number of bosonic degrees of freedom is equal to the total number of fermionic degrees of freedom. The source of this can be made obvious by re-writing our prototype in a slightly different way,

$$\begin{aligned} F(z) = & C(x) - \theta^2 M(x) - \bar{\theta}^2 \overline{M}(x) + \theta^\alpha \bar{\theta}^{\dot{\alpha}} A_{\underline{a}}(x) + \theta^2 \bar{\theta}^2 d(x) \\ & + \theta^\alpha \chi_\alpha(x) + \bar{\theta}^{\dot{\alpha}} \bar{\chi}_{\dot{\alpha}}(x) - \bar{\theta}^2 \theta^\alpha \lambda_\alpha(x) - \theta^2 \bar{\theta}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}}(x) \quad , \end{aligned} \quad (73)$$

and it is seen that the fields are in one-to-one correspondence with the monomials that are constructed from  $\theta$  and  $\bar{\theta}$ . These monomials are in turn isomorphic to a Clifford algebra so there the total number of odd monomials is equal to the total number even monomials. This translates into counting fields as follows,

$$\begin{aligned} N(C) &= 1 && \text{because } C = C^* \quad , \\ N(M) &= 2 && \text{because } M \neq M^* \quad , \\ N(A_{\underline{a}}) &= 4 && \text{because } A_{\underline{a}} = A_{\underline{a}}^* \text{ \& } \underline{a} = 0, \dots, 3 \quad , \\ N(d) &= 1 && \text{because } d = d^* \quad , \\ \rightarrow N_B &= N(C) + N(M) + N(A_{\underline{a}}) + N(d) = 8 \quad . \\ N(\lambda^\alpha) &= 4 && \text{because } \lambda^\alpha \neq [\lambda^\alpha]^* \text{ \& } \alpha = 1, 2 \quad , \\ N(\chi^\alpha) &= 4 && \text{because } \chi^\alpha \neq [\chi^\alpha]^* \text{ \& } \alpha = 1, 2 \quad , \\ \rightarrow N_F &= N(\lambda^\alpha) + N(\chi^\alpha) = 8 \quad . \end{aligned} \quad (74)$$

This same type of procedure can be applied to the chiral scalar superfield with the result that  $N_B = N_F = 4$ , so the chiral multiplet is literally a smaller representation containing fewer independent degrees of freedom than the real scalar superfield.

The equality in the numbers of bosons versus the fermions has far reaching implications for the quantum behavior of supersymmetrical systems. It is the case that the quantum corrections to a particular theory are determined by Feynman diagrams containing “loops.” The fact that fermionic loops have minus signs relative to bosonic loops implies the possibility of some types of cancellations.

### 1.9 Superspace Integration

Berezin was the first to introduce integration over Grassmann numbers. It is useful to review this. Consider a “toy” superfunction dependent on a single

coordinate  $t$  and a single Grassmann coordinate  $\theta$ , then

$$f(z) = A(t) + i\theta\psi(t) \quad (76)$$

(assume  $f(z) = f^*(z)$  for simplicity). Berezin's definition of integration was

$$\int d\theta f(z) = i\psi(t) \quad , \quad (77)$$

but this is the same as

$$\frac{\partial}{\partial\theta}f(z) = i\psi(t) \quad . \quad (78)$$

For Grassmann numbers, integration is the same as differentiation! Using this as a hint we may define the superspace integral by

$$\int d^4\theta = \int d^2\theta d^2\bar{\theta} = \frac{1}{4}\eta^{ab}\frac{\partial}{\partial\theta^\alpha}\frac{\partial}{\partial\theta^\beta}\frac{\partial}{\partial\bar{\theta}^\alpha}\frac{\partial}{\partial\bar{\theta}^\beta} \quad . \quad (79)$$

However, given the forms of  $D_\alpha$  and  $\bar{D}_\alpha$  this definition is equivalent to the following (up to total derivatives)

$$\begin{aligned} \int d^4\theta \mathcal{L} &\equiv \lim_{\theta \rightarrow 0} \lim_{\bar{\theta} \rightarrow 0} \frac{1}{4}\eta^{ab}D_\alpha D_\beta \bar{D}_\alpha \bar{D}_\beta \mathcal{L} \quad , \\ &\equiv D^2 \bar{D}^2 \mathcal{L} \Big| \quad . \end{aligned} \quad (80)$$

Thus we can calculate  $\int d^4\theta F(z)$  for the superfield in (56) to find

$$\int d^4\theta F(z) = d(x) \quad , \quad (81)$$

just like Berezin's result. This is called the “d-component” of a superfield and can be applied to *any* SF to construct an effective supersymmetry invariant,

$$\delta_Q \left( \int d^4\theta \mathcal{L} \right) = \text{total div} \quad . \quad (82)$$

We can say that the action is effectively invariant because this equation shows us that the action (i.e.  $\mathcal{S} = \int d^4x \int d^4\theta \mathcal{L}$ ) only changes by surface terms which are usually ignored in field theories anyway).

For the chiral SF we must be a bit more careful. We note

$$\int d^4\theta \Phi = \int d^4\theta \bar{\Phi} = 0 \quad , \quad (83)$$

(up to total divergences), but

$$\int d^2\theta \Phi \equiv \lim_{\theta \rightarrow 0} D^2\Phi = D^2\Phi| = F(x) \quad , \quad (84)$$

and the super translation variation of this satisfies

$$\delta_Q \left( \int d^2\theta \mathcal{L}_c \right) = \text{total div} \quad . \quad (85)$$

This is called the “ $F$ -component” invariant and it can be used for any chiral superfield. The feature that is special about a chiral SF is that it satisfies a certain differential constraint,  $\overline{D}_\alpha \Phi = 0$ , which makes it unnecessary to integrate over the complete set of superspace coordinates. It is only integrated over a “chiral sub-space.”

So we have three superspace measures with which to build actions

$$\begin{aligned} \int d^6z &\equiv \int d^4x d^2\theta \mathcal{L}_c \quad , \quad \text{if } \overline{D}_\alpha \mathcal{L}_c = 0 \quad , \\ \int d^6\bar{z} &\equiv \int d^4x d^2\bar{\theta} \overline{\mathcal{L}}_c \quad , \quad \text{if } D_\alpha \overline{\mathcal{L}}_c = 0 \quad , \\ \int d^8z &\equiv \int d^4x d^2\theta d^2\bar{\theta} \mathcal{L} \quad , \quad \text{if } \mathcal{L} \text{ is unrestricted} \quad . \end{aligned} \quad (86)$$

### 1.10 Putting It All Together: The Dynamics of the Chiral SF

We can now contemplate the existence of SF Lagrangians

$$\mathcal{L} = \mathcal{L}(\Phi, D_\alpha \Phi, \dots) \quad , \quad (87)$$

that are the analogs of ordinary Lagrangians. These Lagrangians can lead to actions with a definition of superspace integration theory. We will end this discussion by showing how all of the pieces we have introduced in this section are used to write supersymmetrical field theories which turn out to remarkably similar to ordinary relativistic field theories. For the purposes of this illustration we treat the free massive chiral superfield<sup>4</sup>.

As is customary this theory is described by

$$\mathcal{S} = \int d^8z \overline{\Phi} \Phi + \left[ \int d^6z \left( \frac{1}{2} m \Phi^2 \right) + \text{h.c.} \right] \quad . \quad (88)$$

This action involves two functions, the Kähler potential ( $K = \overline{\Phi} \Phi$ ) and the superpotential ( $W = \frac{1}{2} m \Phi^2$ ) In principle we can use explicit  $\theta$ -expansions to

derive the “d” component of the first superfield and the “F” term of the second, but that is not efficient. Instead we note that the component fields of  $\Phi$  may be thought of as Taylor series coefficients of  $\Phi$  which are defined by

$$A(x) \equiv \Phi \Big| , \quad \psi_\alpha \equiv D_\alpha \Phi \Big| , \quad F(x) \equiv D^2 \Phi \Big| . \quad (89)$$

Under this circumstance and recalling that the superspace integrations  $\int d^8 z$  and  $\int d^6 z$  actually correspond to fourth-order and second-order spinorial differential operators, respectively, opens the following route for calculations.

$$\begin{aligned} \int d^6 z \left( \frac{1}{2} m \Phi^2 \right) &= \frac{1}{4} m \int d^4 x D^\alpha D_\alpha \Phi^2 \Big| \\ &= \frac{1}{2} m \int d^4 x D^\alpha [\Phi D_\alpha \Phi] \Big| \\ &= \frac{1}{2} m \int d^4 x [ (D^\alpha \Phi) (D_\alpha \Phi) + \Phi (D^\alpha D_\alpha \Phi) ] \Big| \quad (90) \\ &= \frac{1}{2} m \int d^4 x [ \psi^\alpha \psi_\alpha + A(2F) ] \\ &= \int d^4 x [ \frac{1}{2} m \psi^\alpha \psi_\alpha + m A F ] . \end{aligned}$$

In a similar manner we find

$$\begin{aligned} \int d^8 z \bar{\Phi} \Phi &= \int d^4 x D^2 \bar{D}^2 \bar{\Phi} \Phi \Big| = \int d^4 x \bar{D}^2 D^2 \bar{\Phi} \Phi \Big| \\ &= \int d^4 x \bar{D}^2 [ \bar{\Phi} D^2 \Phi ] \Big| \\ &= \int d^4 x [ (\bar{D}^2 \bar{\Phi}) (D^2 \Phi) + (\bar{D}^\alpha \bar{\Phi}) (\bar{D}_\alpha D^2 \Phi) + \bar{\Phi} ((\bar{D}^2 D^2 \Phi)) ] \Big| . \quad (91) \end{aligned}$$

Now observe

$$\begin{aligned} \bar{D}_\alpha \bar{\Phi} &= 0 \quad \& \quad \text{by definition} \quad D_\alpha \bar{D}_\alpha + \bar{D}_\alpha D_\alpha = i \partial_{\underline{a}} , \\ &\rightarrow \bar{D}_\alpha D_\alpha \Phi = i \partial_{\underline{a}} \Phi . \end{aligned} \quad (92)$$

This result can be used to simplify the differential operators in (91). We basically need to “push” the  $\bar{D}$  operators in the second and third terms until

they act on the factors of  $\Phi$ . This can be obtained systematically as

$$\begin{aligned}
\overline{D}_\alpha \dot{D}^2 \Phi &= \frac{1}{2} \overline{D}_\alpha \dot{D}^\alpha D_\alpha \Phi = -\frac{1}{2} \overline{D}_\alpha \dot{D}_\alpha D^\alpha \Phi \\
&= \frac{1}{2} D_\alpha \overline{D}_\alpha \dot{D}^\alpha \Phi - i \frac{1}{2} \partial_\alpha \dot{D}^\alpha \Phi \\
&= -\frac{1}{2} D^\alpha \overline{D}_\alpha \dot{D}_\alpha \Phi - i \frac{1}{2} \partial_\alpha \dot{D}^\alpha \Phi \\
&= -i \partial_\alpha \dot{D}^\alpha \Phi \quad ,
\end{aligned} \tag{93}$$

where in obtaining this result we have liberally used various points discussed earlier in this presentation. We use the last line of (93) once more to see,

$$\begin{aligned}
\overline{D}^2 D^2 \Phi &= \frac{1}{2} \overline{D}^\alpha \dot{\overline{D}}_\alpha D^2 \Phi = -i \frac{1}{2} \overline{D}^\alpha \partial_\alpha \dot{D}^\alpha \Phi \\
&= -i \frac{1}{2} \partial_\alpha \overline{D}^\alpha \dot{D}^\alpha \Phi \\
&= \frac{1}{2} \partial_\alpha \partial^\alpha \Phi = \square \Phi \quad .
\end{aligned} \tag{94}$$

The final results of (93,94) are now substituted into the last two term of (91) and yield,

$$\int d^8 z \overline{\Phi} \Phi = \int d^4 x [ \overline{F} F - i \overline{\psi}^\alpha \partial_\alpha \psi^\alpha + \overline{A} \square A ] \quad . \tag{95}$$

We have now proven that the action in (88) is equivalent to the following ordinary looking relativistic field theory,

$$\begin{aligned}
\mathcal{S} &= \int d^4 x [ -\frac{1}{2} (\partial^\alpha \overline{A}) (\partial_\alpha A) - i \overline{\psi}^\alpha \partial_\alpha \psi^\alpha + \overline{F} F \\
&\quad + \frac{1}{2} m (\psi^\alpha \psi_\alpha + \overline{\psi}^\alpha \overline{\psi}_\alpha) + m (A F + \overline{A} \overline{F}) ] \quad .
\end{aligned} \tag{96}$$

By re-writing the fermionic translation laws of the fields in terms of four component fermions, we can see that the real part of  $A$  corresponds to a scalar ( $J^P = 0^+$ ) and its imaginary part corresponds to a pseudoscalar ( $J^P = 0^-$ ). In the presence of a nonvanishing mass term the four component spinor corresponds to a Majorana (or real spinor). In the massless limit the spinor may be regarded as either a Majorana or Weyl spinor. The example above shows that 4D superfields are particularly sensitive to the parity assignments of fields. For, example it seems to be impossible to construct a 4D theory which contains two scalar fields and one Weyl or Majorana field.

There is another point about the component actions that we write (but derive from superspace). It is often the case that the simplest definition of

the component fields (similar to (89)) for a supermultiplet leads to unusual normalization compared to “standard” (i.e. non-supersymmetrical definitions). This can always be ‘fixed’ by choosing the definitions of the component fields appropriately. We shall not (as is the usual custom) concern ourselves with this issue except for pointing this out.

By varying this action in the usual way we can derive the following equations of motion,

$$\begin{aligned}\square A + m \bar{F} &= 0 \quad , \\ -i \partial_{\underline{a}} \psi^\alpha + m \bar{\psi}_{\dot{\alpha}} &= 0 \quad , \\ F + m \bar{A} &= 0 \quad .\end{aligned}\tag{97}$$

The last of these is purely algebraic, it may be used to solve for the field  $F$  as  $F = -m \bar{A}$ . For this reason  $F$  is called an “auxiliary field variable” or “auxiliary field.” It can be used in the first equation in (97) to write the equations of motion for  $A(x)$  and  $\psi^\alpha(x)$

$$\begin{aligned}(\square - m^2) A &= 0 \quad , \\ -i \partial_{\underline{a}} \psi^\alpha + m \bar{\psi}_{\dot{\alpha}} &= 0 \quad .\end{aligned}\tag{98}$$

These equations show another characteristic feature of supersymmetric theories; strict supersymmetry implies a degeneracy of the mass spectrum between some bosons and fermions.

We can also use the equation of motion for  $F$  to eliminate it from the action in (96). The resultant action is called the “on-shell” action.

$$\begin{aligned}\mathcal{S}_{on-shell} &= \int d^4x \left[ -\frac{1}{2} (\partial^{\underline{a}} \bar{A}) (\partial_{\underline{a}} A) + m^2 \bar{A} A \right. \\ &\quad \left. - i \bar{\psi}^{\dot{\alpha}} \partial_{\underline{a}} \psi^\alpha + \frac{1}{2} m (\psi^\alpha \psi_\alpha + \bar{\psi}^{\dot{\alpha}} \bar{\psi}_{\dot{\alpha}}) \right] .\end{aligned}\tag{99}$$

We close this lecture by once again noting that an intuitive picture of superspace posits that it is a mathematic structure whose boundary is to be identified with an ordinary bosonic space. This view is very simple and has as its implication that the physics we experience in the universe around us occurs exclusively on this boundary. Later we will discuss integration theory in the case where the bosonic manifold is “curved” and we will see that this view has a simplifying consequence for the generalization of Berezin’s integration theory.



## 2 Second Lecture: Sophomore 4D, N = 1 Superfield Theory

### Introduction

In the second lecture, we will discuss a number of examples of 4D, N = 1 supersymmetrical field theories. These include; (a.) the renormalizable interacting, scalar multiplet, (b.) the gauge-vector multiplet, (c.) Kähler models, (e.) the simplex of super p-forms and (f.) the nonminimal multiplet. The main goal is to show how field theory in 4D, N = 1 superspace is very, very similar to field theory in Minkowski space. Along the way we will learn how to construct Yang-Mills gauge theories and non-linear  $\sigma$ -models.

### Lecture

#### 2.1 The 4D, N = 1 Interacting Chiral Multiplet

We can generalize the superpotential  $W(\Phi)$  of the previous section by considering an action of the form,

$$\mathcal{S} = \int d^8z K(\Phi, \bar{\Phi}) + \left[ \int d^6z W(\Phi) + \text{c.c.} \right] , \quad (100)$$

where  $W(\Phi) = \frac{1}{2}m\Phi^2 + \frac{1}{3!}\lambda\Phi^3$ . Using the same techniques as before (90) but applied to  $W$ , we derive

$$\begin{aligned} \mathcal{S} = \int d^4x \left\{ -\frac{1}{2}(\partial^{\underline{a}}\bar{A})(\partial_{\underline{a}}A) - i\bar{\psi}^{\dot{\alpha}}\partial_{\underline{a}}\psi^{\alpha} + \bar{F}F \right. \\ \left. + [(m + \lambda A)(\frac{1}{2}\psi^{\alpha}\psi_{\alpha} + AF) + \text{c.c.}] \right\} , \end{aligned} \quad (101)$$

and using the new algebraic equation of motion for  $F$  leads to the action

$$\begin{aligned} \mathcal{S} = \int d^4x \left\{ -\frac{1}{2}(\partial^{\underline{a}}\bar{A})(\partial_{\underline{a}}A) - i\bar{\psi}^{\dot{\alpha}}\partial_{\underline{a}}\psi^{\alpha} + \frac{1}{2}m(\psi^{\alpha}\psi_{\alpha} + \bar{\psi}^{\dot{\alpha}}\bar{\psi}_{\dot{\alpha}}) \right. \\ \left. + m^2 A\bar{A} - \frac{1}{2}m\lambda(A\bar{A}^2 + \bar{A}A^2) - \frac{1}{4}\lambda A^2\bar{A}^2 \right. \\ \left. + \frac{1}{2}\lambda(A\psi^{\alpha}\psi_{\alpha} + \bar{A}\bar{\psi}^{\dot{\alpha}}\bar{\psi}_{\dot{\alpha}}) \right\} . \end{aligned} \quad (102)$$

We see a complex spin-0 field

$$A = \frac{1}{\sqrt{2}}[a(x) + i b(x)] \quad \text{scalar , pseudoscalar,} \quad (103)$$

and a Majorana spinor,

$$\psi_{(\alpha)}(x) = \begin{pmatrix} \psi_{\alpha}(x) \\ \bar{\psi}_{\dot{\alpha}}(x) \end{pmatrix} , \quad (104)$$

both with a mass  $m$ . These fields interact via a “ $\varphi^4$ -term” (coupling constant  $\frac{1}{4}\lambda$ ), a “Yukawa-term” (coupling constant  $\frac{1}{2}\lambda$ ) and a “ $\varphi^3$ -term” (coupling constant  $\frac{1}{2}m\lambda$ ). So in addition to imposing the condition  $m_{Spin-0} = m_{Spin-1/2}$  supersymmetry also imposes a set of relationships among the coupling constants of the  $\varphi^4$ -,  $\varphi^3$ - and Yukawa interaction terms.

## 2.2 $U(1)$ Symmetry in $4D$ , $N = 1$ Superspace

Since  $A$  is complex, at first glance it might seem possible to realize the  $U(1)$  symmetry of QED in the form

$$[A]' = e^{ig\alpha} A \quad , \quad [\psi_{\alpha}]' = e^{ig\alpha} \psi_{\alpha} \quad . \quad (105)$$

There are a couple of reasons why this is incorrect. First we note that unless  $m = \lambda = 0$ , these transformations are *not* symmetries of the action above. Next even if we permitted  $m = \lambda = 0$ , it is still not possible to use these transformation to realize a  $U(1)$  symmetry. One reason for this is because the transformation  $[A]' = e^{i\alpha} A$  “rotates” a scalar into a pseudoscalar. Although an axial  $U(1)$  charge (or  $U_A(1)$  charge) has this property, the  $U(1)$  charge of QED does not possess this property. So in order to introduce a rigid  $U(1)$  symmetry we must modify the starting point.

Let us consider the alternate action

$$\mathcal{S} = \int d^8z [\bar{\Phi}_+ \Phi_+ + \bar{\Phi}_- \Phi_-] + \int d^6z [W(\Phi_+, \Phi_-) + \text{c.c.}] \quad , \quad (106)$$

which is obtained by introducing *two* independent chiral scalar superfields  $\Phi_+$  and  $\Phi_-$ . The first two terms in the action clearly possess a rigid  $U(1)$  invariance. For the sake of renormalizability the polynomial superpotential  $W(\Phi_+, \Phi_-)$  must be of order three or less. By a proper choice of this polynomial, this action can be made to be invariant under

$$[\Phi_+]' = e^{ig\alpha} \Phi_+ \quad , \quad [\Phi_-]' = e^{-ig\alpha} \Phi_- \quad . \quad (107)$$

The unique choice of such a polynomial of order three or less is  $W(\Phi_+, \Phi_-) = m \Phi_+ \Phi_-$ . The two scalars  $a_+(x)$  and  $a_-(x)$  may be regarded as the real and

imaginary part of a complex scalar required to realize the U(1) symmetry of QED. Similarly, we can now define a Dirac spinor

$$\psi_{(\alpha)}(x) = \begin{pmatrix} \psi_{\alpha+}(x) \\ \psi_{\alpha-}(x) \end{pmatrix} = \begin{pmatrix} D_{\alpha}\Phi_+ \\ \overline{D}_{\dot{\alpha}}\overline{\Phi}_- \end{pmatrix} , \quad (108)$$

and under the rigid U(1) symmetry this transforms as

$$\left[\psi_{(\alpha)}(x)\right]' = e^{ig\alpha} \psi_{(\alpha)}(x) . \quad (109)$$

We can identify this Dirac spinor as the electron. It is accompanied by two scalar  $(a_+(x), a_-(x))$  and two pseudoscalar  $(b_+(x), b_-(x))$  collectively called “selectrons.” This is another example of the bose-fermi equality of supersymmetrical theories. It is well known that the ordinary Dirac field describes four dynamical fermionic degrees of freedom. The selectrons are the corresponding bosonic degrees of freedom.

We now wish to “gauge” this symmetry by replacing the constant  $\alpha$  by a chiral superfield  $\Lambda$  (i.e.  $\overline{D}_{\dot{\alpha}}\Lambda = 0$ ) so that the local U(1) transformation laws of the matter superfields  $\Phi_+$  and  $\Phi_-$  may be cast into the form

$$\left[\Phi_+\right]' = e^{ig\Lambda}\Phi_+ , \quad \left[\Phi_-\right]' = e^{-ig\Lambda}\Phi_- . \quad (110)$$

For the superpotential term this causes no problems,

$$\left[\Phi_+\Phi_-\right]' = \left[e^{ig\Lambda}\Phi_+e^{-ig\Lambda}\Phi_-\right] = \left[\Phi_+e^{ig\Lambda}e^{-ig\Lambda}\Phi_-\right] = \Phi_+\Phi_- . \quad (111)$$

However for the Kähler potential we see

$$\left[\overline{\Phi}_+\Phi_+ + \overline{\Phi}_-\Phi_-\right]' = \left[\overline{\Phi}_+e^{-ig\bar{\Lambda}}e^{ig\Lambda}\Phi_+ + \overline{\Phi}_-e^{ig\bar{\Lambda}}e^{-ig\Lambda}\Phi_-\right] . \quad (112)$$

This is a major problem since  $\Lambda \neq \bar{\Lambda}$  implies that the Kähler potential terms are *not* gauge invariant. The solution to this problem is to introduce a “photon superfield.” If we recall (56), it is pretty clear how this should be done. There we saw that a real scalar superfield contains a spin-1 field. Via a number of arguments, we can show that the spin-1 field that appears in (56) is not a vector but an axial vector. So as a first step, we need to introduce a pseudoscalar superfield to be denoted by  $V$  (its  $\theta\bar{\theta}$  term is a vector). Further by noting that  $V \sim \theta^{\alpha}\bar{\theta}^{\dot{\alpha}}A_{\underline{a}}$  and  $\Lambda \sim i\frac{1}{2}\theta^{\alpha}\bar{\theta}^{\dot{\alpha}}\partial_{\underline{a}}\alpha(x)$  we see that

$$\left[V\right]' = V - i(\Lambda - \bar{\Lambda}) \rightarrow \left[A_{\underline{a}}(x)\right]' = \left[A_{\underline{a}}(x)\right] + \frac{1}{2}\partial_{\underline{a}}[\alpha(x) + \bar{\alpha}(x)] . \quad (113)$$

The pseudoscalar superfield can be used to modify the Kähler potential term according to,

$$K = \left[ \bar{\Phi}_+ e^{gV} \Phi_+ + \bar{\Phi}_- e^{-gV} \Phi_- \right] , \quad (114)$$

and under the U(1) gauge transformation we now find

$$\begin{aligned} \left[ \bar{\Phi}_+ e^{gV} \Phi_+ + \bar{\Phi}_- e^{-gV} \Phi_- \right]' &= \left[ \bar{\Phi}_+ e^{-ig\bar{\Lambda}} e^{gV'} e^{ig\Lambda} \Phi_+ \right. \\ &\quad \left. + \bar{\Phi}_- e^{ig\bar{\Lambda}} e^{-gV'} e^{-ig\Lambda} \Phi_- \right] \\ &= \left[ \bar{\Phi}_+ e^{gV} \Phi_+ + \bar{\Phi}_- e^{-gV} \Phi_- \right] . \end{aligned} \quad (115)$$

So the matter field terms in a gauge invariant supersymmetric version of QED can be written as<sup>5</sup>

$$\begin{aligned} \mathcal{S}_{QED-matter} &= \int d^8z \left[ \bar{\Phi}_+ e^{gV} \Phi_+ + \bar{\Phi}_- e^{-gV} \Phi_- \right] \\ &\quad + \int d^6z \left[ m \Phi_+ \Phi_- + \text{c.c.} \right] . \end{aligned} \quad (116)$$

Apparently minimal coupling of Yang-Mills symmetries in superspace corresponds to the insertion of the factors involving  $V$ . As we shall later see this action can be shown to contain the usual Dirac action for the four component spinor defined in (108).

### 2.3 Review of Maxwell Theory in Minkowski Space

We now require a generalization of the Maxwell action, More generally it is useful to possess a generalization of the Yang-Mills action. To begin this discussion let us review ordinary 4D Maxwell theory. As is customary, we can introduce a covariant derivative

$$\nabla_{\underline{a}} = \partial_{\underline{a}} - i A_{\underline{a}} t , \quad (117)$$

where  $t$  is the abstract U(1) Lie algebra generator. If a matter field  $\phi(x)$  transforms as

$$\begin{aligned} \left[ \phi \right]' &= \exp[iQ\lambda(x)] \phi(x) \\ &\approx \phi(x) + iQ\lambda(x)\phi(x) \text{ if } \lambda \ll 1 \\ \rightarrow \delta\phi &\equiv \left[ \phi \right]' - \phi = iQ\lambda(x)\phi(x) , \end{aligned} \quad (118)$$

then by definition  $[t, \phi] = Q\phi$ . Although not usually introduced in such discussions,  $t$  allows us to regard  $\nabla_{\underline{a}}$  as a real operator if  $t^* = -t$ . This follows from

$$\begin{aligned} \left( [t, \phi] \right)^* &= Q \left( \phi \right)^* \\ [t^*, \bar{\phi}] &= Q \bar{\phi} \\ [t, \bar{\phi}] &= -Q \bar{\phi} \quad , \end{aligned} \tag{119}$$

and this is what we want since  $\phi(x)$  is associated with a particle possessing charge  $Q$  then the anti-particle should be associated with  $\bar{\phi}(x)$  possessing charge  $-Q$ . Thus,

$$\begin{aligned} \left( \nabla_{\underline{a}} \right)^* &= \left[ \partial_{\underline{a}} - iA_{\underline{a}}t \right]^* \\ &= \partial_{\underline{a}} + iA_{\underline{a}}t^* \\ &= \partial_{\underline{a}} - iA_{\underline{a}}t \quad . \end{aligned} \tag{120}$$

The field strength  $f_{\underline{a}\underline{b}}$  is defined by

$$\begin{aligned} -i f_{\underline{a}\underline{b}} t &\equiv [\nabla_{\underline{a}}, \nabla_{\underline{b}}] = [\partial_{\underline{a}} - iA_{\underline{a}}t, \partial_{\underline{b}} - iA_{\underline{b}}t] \\ &= -i(\partial_{\underline{a}}A_{\underline{b}} - \partial_{\underline{b}}A_{\underline{a}})t - [A_{\underline{a}}t, A_{\underline{b}}t] \\ &= -i(\partial_{\underline{a}}A_{\underline{b}} - \partial_{\underline{b}}A_{\underline{a}})t - A_{\underline{a}}[t, A_{\underline{b}}]t \\ &\quad - A_{\underline{b}}[t, A_{\underline{a}}]t - A_{\underline{a}}A_{\underline{b}}[t, t] \quad . \end{aligned} \tag{121}$$

All of the terms quadratic in the gauge field will vanish if we define

$$[t, A_{\underline{a}}] \equiv 0 \quad , \quad [t, t] \equiv 0 \quad , \quad \rightarrow f_{\underline{a}\underline{b}} = (\partial_{\underline{a}}A_{\underline{b}} - \partial_{\underline{b}}A_{\underline{a}}) \quad . \tag{122}$$

(for nonabelian groups this changes) and the Maxwell action is simply

$$\mathcal{S}_{Maxwell} = -\frac{1}{8} \int d^4x f_{\underline{a}\underline{b}}^a f_{\underline{a}\underline{b}}^a \quad . \tag{123}$$

(The strange-looking factor of 1/8 is a consequence of our definitions in section 1.7.)

#### 2.4 A $U(1)$ Covariant Derivative in $4D$ , $N = 1$ Superspace

In the first lecture, we saw that there exists in superspace a “supergradient operator”  $D_{\underline{A}} \equiv (D_{\alpha}, \bar{D}_{\dot{\alpha}}, \partial_{\underline{a}})$  that generalizes the ordinary gradient  $\partial_{\underline{a}}$ . So it is natural to define a  $4D$ ,  $N = 1$   $U(1)$  covariant operator

$$\nabla_{\underline{A}} \equiv D_{\underline{A}} - i\Gamma_{\underline{A}}t \quad , \tag{124}$$

and use it to define a superspace field strength  $F_{\underline{A}\underline{B}}$ . There is something that needs attention prior to deriving the explicit form of this. We note

$$\begin{aligned} [D_\alpha, D_\beta] &= 0 \quad , \quad [\bar{D}_{\dot{\alpha}}, \bar{D}_{\dot{\beta}}] = 0 \quad , \quad [D_\alpha, \bar{D}_{\dot{\alpha}}] = i\partial_{\underline{a}} \quad , \\ [D_\alpha, \partial_{\underline{b}}] &= 0 \quad , \quad [\bar{D}_{\dot{\alpha}}, \partial_{\underline{b}}] = 0 \quad , \\ [\partial_{\underline{a}}, \partial_{\underline{b}}] &= 0 \quad . \end{aligned} \quad (125)$$

All of these results can be collectively written in the form

$$[D_{\underline{A}}, D_{\underline{B}}] = C_{\underline{A}\underline{B}}^{\underline{C}} D_{\underline{C}} \quad , \quad (126)$$

for some appropriate coefficients  $C_{\underline{A}\underline{B}}^{\underline{C}}$  known as the “object of anholonomy” or more simply as the “anholonomy.” The values of these coefficients are simply read off by comparing this definition with the explicit graded commutators above (125). The appearance of these is important due to the definition of the field strength in superspace. It is defined by

$$[\nabla_{\underline{A}}, \nabla_{\underline{B}}] = C_{\underline{A}\underline{B}}^{\underline{C}} \nabla_{\underline{C}} - i F_{\underline{A}\underline{B}} t \quad , \quad (127)$$

where

$$F_{\underline{A}\underline{B}} t = D_{\underline{A}} \Gamma_{\underline{B}} t - (-)^{AB} D_{\underline{B}} \Gamma_{\underline{A}} t - C_{\underline{A}\underline{B}}^{\underline{C}} \Gamma_{\underline{C}} t \quad . \quad (128)$$

### 2.5 Finding and Solving Covariant Constraints

In the presence of local symmetries, the prior definition of a chiral superfield is not obviously gauge covariant (well it really is by a certain definition) and this suggests that the differential constraint used to define a chiral superfield should be changed according to

$$\bar{\nabla}_{\dot{\alpha}} \Phi = 0 \quad , \quad (129)$$

and since this condition is satisfied then  $\bar{\nabla}_{\dot{\alpha}} \bar{\nabla}_{\dot{\beta}} \Phi = 0$ . If we symmetrize this last result on the  $\dot{\alpha}$  and  $\dot{\beta}$  indices then use the definition of the superspace field strength we find

$$0 = [\bar{\nabla}_{\dot{\alpha}}, \bar{\nabla}_{\dot{\beta}}] \Phi = F_{\dot{\alpha}\dot{\beta}} t \Phi = F_{\dot{\alpha}\dot{\beta}} Q \Phi \quad . \quad (130)$$

This final equation is called an “integrability condition”<sup>6</sup>. It has three solutions;

(A.)  $Q = 0$  (The matter superfield carries no charge.)

(B.)  $F_{\alpha\dot{\beta}} = 0$  (This looks crazy. In ordinary Maxwell theory if we impose the condition  $F_{\underline{a}\underline{b}} = 0$  then its solution is  $A_{\underline{a}} = \partial_{\underline{a}} a$  and there is no photon in the theory.)

(C.)  $\Phi = 0$ . (There is no matter superfield at all.)

Thus, at first it seems as though the notation of a superspace  $U(1)$  covariant derivative has brought us to a bad ending.

However, a pleasant surprise is to be had by studying the second option in more detail. We are able to solve the restriction in option (b.) by following the example of ordinary 4D spacetime.

$$\begin{aligned} F_{\alpha\dot{\beta}} t = 0 &\rightarrow \overline{D}_{\dot{\alpha}} \Gamma_{\dot{\beta}} + \overline{D}_{\dot{\beta}} \Gamma_{\dot{\alpha}} = 0 \quad , \\ &\rightarrow \Gamma_{\dot{\alpha}} t = i (\overline{D}_{\dot{\alpha}} \overline{\Omega}) t = i e^{-\overline{\Omega} t} \overline{D}_{\dot{\alpha}} e^{\overline{\Omega} t} \quad , \\ &\rightarrow \Gamma_{\alpha} t = i (D_{\alpha} \overline{\Omega}) t = i e^{-\Omega t} D_{\alpha} e^{\Omega t} \quad , \end{aligned} \tag{131}$$

where we have obtained the last line by applying the superspace conjugation operator to the penultimate line. Similarly to ordinary Minkowski space, the solution yields a scalar superfield. However, unlike ordinary Minkowski space, the scalar superfield<sup>b</sup>  $\Omega$  is complex. This seemingly minor difference is crucially important as we shall shortly see.

Let us continue by examining the form of  $F_{\alpha\dot{\beta}}$ ,

$$\begin{aligned} F_{\alpha\dot{\beta}} &= D_{\alpha} \Gamma_{\dot{\beta}} + \overline{D}_{\dot{\beta}} \Gamma_{\alpha} - C_{\alpha\dot{\alpha}} \overline{C}_{\dot{\beta}\underline{\beta}} \Gamma_{\underline{\beta}} \\ &= D_{\alpha} \Gamma_{\dot{\beta}} + \overline{D}_{\dot{\beta}} \Gamma_{\alpha} - i \Gamma_{\underline{\beta}} \\ &= i D_{\alpha} \overline{D}_{\dot{\beta}} \overline{\Omega} + i \overline{D}_{\dot{\beta}} D_{\alpha} \Omega - i \Gamma_{\underline{\beta}} \quad . \end{aligned} \tag{132}$$

The interesting point regarding this field strength is that the gauge field which “points” in the Minkowski space direction appears algebraically. This means if we make the definition

$$\Gamma_{\underline{a}} = \overline{D}_{\dot{\alpha}} D_{\alpha} \Omega - D_{\alpha} \overline{D}_{\dot{\alpha}} \overline{\Omega} \quad , \tag{133}$$

---

<sup>b</sup>In 1980 when W. Siegel and I first introduced the term “pre-potential<sup>7</sup>,” it was in the context of quantities of this type that we meant it.

then  $F_{\alpha\beta} = 0$ . So  $\Gamma_{\underline{a}} \neq 0$  even though  $F_{\alpha\beta} = F_{\alpha\dot{\beta}} = F_{\alpha\beta} = 0$ . However, the more important matter is whether this choice for  $\Gamma_{\underline{a}}$  describes a “pure gauge” configuration.

Writing  $\Omega = \frac{1}{2}(V + iW)$  where  $V = V^*$  and  $W = W^*$  we obtain

$$\begin{aligned}\Gamma_{\underline{a}} &= -\frac{1}{2}\left([D_{\alpha}, \overline{D}_{\dot{\alpha}}]V\right) + i\frac{1}{2}\left(\{D_{\alpha}, \overline{D}_{\dot{\alpha}}\}W\right) \\ &= -\frac{1}{2}\left([D_{\alpha}, \overline{D}_{\dot{\alpha}}]V\right) - \frac{1}{2}\partial_{\underline{a}}W \quad ,\end{aligned}\tag{134}$$

and only the second term is seen to be a pure gauge configuration. If the superfield  $V$  here is the same as the one that was used to covariantize the kinetic energy terms for the chiral scalar superfields  $\Phi_+$  and  $\Phi_-$ , we have found the relation between the Yang-Mills prepotential  $V$  and the “usual” connections and field strengths that exist as a basis for ordinary electromagnetism. Without loss of generality, we may set the quantity  $W$  to zero.

With the superspace connections all determined as described above, we can now calculate the remaining components of  $F_{\underline{A}\underline{B}}$ . The next one is  $F_{\alpha\underline{b}}$  where we find,

$$\begin{aligned}F_{\alpha\underline{b}} &= D_{\alpha}\Gamma_{\underline{b}} - \partial_{\underline{b}}\Gamma_{\alpha} \\ &= \frac{1}{2}\left(-D_{\alpha}[D_{\beta}\overline{D}_{\dot{\beta}} - \overline{D}_{\dot{\beta}}D_{\beta}]V - i\partial_{\underline{b}}D_{\alpha}V\right) \\ &= C_{\alpha\beta}D^2\overline{D}_{\dot{\beta}}V \\ &\equiv iC_{\alpha\beta}\overline{W}_{\dot{\beta}} \quad .\end{aligned}\tag{135}$$

The superfield  $\overline{W}_{\dot{\beta}}$  must be anti-chiral due to the identity  $D_{\alpha}D^2 = 0$ . By conjugation  $W_{\alpha}$  is a chiral superfield. After a similar set of calculations we find

$$F_{\underline{a}\underline{b}} = \frac{1}{2}\left[C_{\alpha\beta}(\overline{D}_{\dot{\alpha}}\overline{W}_{\dot{\beta}}) + C_{\dot{\alpha}\dot{\beta}}(D_{\alpha}W_{\beta})\right] \quad .\tag{136}$$

At this point, it is useful to employ a little dimensional analysis.

$$\begin{aligned}[x^{\underline{m}}] &= -1 \quad , \quad [d^4x] = -4 \quad , \quad [\partial_{\underline{m}}] = 1 \quad , \quad [D_{\alpha}] = [\overline{D}_{\dot{\alpha}}] = \frac{1}{2} \quad , \\ [D^2] &= [\overline{D}^2] = 1 \quad , \quad \rightarrow [\int d^2\theta] = [\int d^2\bar{\theta}] = 1 \quad ,\end{aligned}\tag{137}$$

$$\begin{aligned}[\Gamma_{\underline{a}}] &= 1 \quad , \quad [\Gamma_{\alpha}] = [\Gamma_{\dot{\alpha}}] = \frac{1}{2} \quad \rightarrow [F_{\underline{a}\underline{b}}] = 2 \quad , \quad [F_{\alpha\underline{b}}] = \frac{3}{2} \quad , \\ \rightarrow [\int d^4x F^{\underline{a}\underline{b}}F_{\underline{a}\underline{b}}] &= -4 + 2 + 2 = 0 \quad .\end{aligned}\tag{138}$$



This last equation implies that in these units, the ordinary Maxwell action is dimensionless (since the units of  $f$  equal those of  $F$ ). From the non-vanishing components of  $F_{\underline{A}\underline{B}}$  there are only two quadratic action that can be formed

$$[\int d^4x d^2\theta W^\alpha W_\alpha] = -4 + 1 + \frac{3}{2} + \frac{3}{2} = 0 \quad , \quad (139)$$

$$[\int d^4x d^2\theta d^2\bar{\theta} F^{\underline{a}\underline{b}} F_{\underline{a}\underline{b}}] = -4 + 1 + 1 + 2 + 2 = 2 \quad . \quad (140)$$

Only the former of these has the same dimensions as the ordinary Maxwell action and we are led to suggest

$$\mathcal{S}_{Maxwell}^{N=1} = \frac{1}{4} \left[ \int d^4x d^2\theta W^\alpha W_\alpha + \text{h.c.} \right] \quad . \quad (141)$$

We can check whether this is correct by deriving the component fields terms described by this action. In order to do this we must define the component fields through the superfield  $W_\alpha$ . In turn this requires knowledge of the form of all of its spinorial derivatives. There is a systematic way in which such results are derived. This technique is called “solving the superspace Bianchi identities.” It would take us out of the introductory nature of these lectures to discuss this technique. So let’s just use the results that can be derived,

$$\begin{aligned} D_\alpha W_\beta &= \frac{1}{2} C^{\dot{\gamma}\delta} F_{\alpha\dot{\gamma}\beta\delta} - i C_{\alpha\beta} d \quad , \\ D^2 W_\alpha &= -i \partial_{\underline{a}} \bar{W}^{\dot{\alpha}} \quad . \end{aligned} \quad (142)$$

Thus, the components of the 4D,  $N = 1$  Yang-Mills supermultiplet are defined by

$$\begin{aligned} W_\alpha \Big| &\equiv \lambda_\alpha(x) \quad , \\ D_\alpha W_\beta \Big| &\equiv \frac{1}{2} C^{\dot{\gamma}\delta} ( f_{\underline{a}\underline{b}}(x) + \tilde{f}_{\underline{a}\underline{b}}(x) ) - i C_{\alpha\beta} d \Big| \quad , \\ \tilde{f}_{\underline{a}\underline{b}}(x) &\equiv i \frac{1}{2} \epsilon_{\underline{a}\underline{b}}^{\underline{c}\underline{d}} f_{\underline{c}\underline{d}}(x) \quad , \\ d \Big| &\equiv d(x) \quad , \\ D^2 W_\alpha \Big| &\equiv -i \partial_{\underline{a}} \bar{\lambda}^{\dot{\alpha}}(x) \quad . \end{aligned} \quad (143)$$

In these equations, the quantity denoted by  $f_{\underline{a}\underline{b}}$  is exactly the same as that which appears in (122). It is the Maxwell field strength. The fermionic field  $\lambda_\alpha$  is known as the “gaugino.” After we evaluate (141), it will turn out that d

is an auxiliary field. In principle at this stage we could follow a path analogous to that which led from (88) to (90). It is more prudent, however, for us to first discuss the case of Yang-Mills theory, evaluate the component action in this case and then retrieve the Maxwell theory as a special case.

## 2.6 Local Lie Algebraic Theories: Yang-Mills

As is well known, all of the results in (117-123) can be generalized to non-abelian groups. For the field theorist, the most fundamental quantities are fields. The simplest is the scalar field  $\phi(x)$ . If this field carries some representation of a Yang-Mills gauge group, we must add an index to it  $\phi^I(x)$ . The abstract Lie algebraic generator  $t_{\mathcal{A}}$  acts on the scalar field according to

$$t_{\mathcal{A}} \phi^I(x) = (\tilde{t}_{\mathcal{A}})^I_{\mathbf{K}} \phi^K(x) \equiv [t_{\mathcal{A}}, \phi^I(x)] \quad . \quad (144)$$

In writing this equation, we adhere to the convention that  $t_{\mathcal{A}}$  is an operator which obeys a Leibnizian rule when acting on a product of field variables, for example

$$t_{\mathcal{A}}(\phi^I(x) \phi^K(x)) = (t_{\mathcal{A}} \phi^I(x)) \phi^K(x) + \phi^I(x) (t_{\mathcal{A}} \phi^K(x)) \quad . \quad (145)$$

On the other hand, the quantities  $(\tilde{t}_{\mathcal{A}})^I_{\mathbf{K}}$  are a set of constants, i.e. a matrix. The above assumptions have the following consequences,

$$t_{\mathcal{A}} t_{\mathcal{B}} \phi^I(x) = (\tilde{t}_{\mathcal{B}})^I_{\mathbf{K}} t_{\mathcal{A}} \phi^K(x) = (\tilde{t}_{\mathcal{B}})^I_{\mathbf{K}} (\tilde{t}_{\mathcal{A}})^{\mathbf{K}}_{\mathbf{L}} \phi^{\mathbf{L}}(x) \quad . \quad (146)$$

$$\rightarrow [t_{\mathcal{A}}, t_{\mathcal{B}}] \phi^I(x) = - ([\tilde{t}_{\mathcal{A}}, \tilde{t}_{\mathcal{B}}])^I_{\mathbf{K}} \phi^K(x) \quad . \quad (147)$$

This last equation is consistent if

$$[t_{\mathcal{A}}, t_{\mathcal{B}}] = -if_{\mathcal{A}\mathcal{B}}^{\mathcal{C}} t_{\mathcal{C}} \quad , \quad ([\tilde{t}_{\mathcal{A}}, \tilde{t}_{\mathcal{B}}])^I_{\mathbf{K}} = if_{\mathcal{A}\mathcal{B}}^{\mathcal{C}} (\tilde{t}_{\mathcal{C}})^I_{\mathbf{K}} \quad . \quad (148)$$

In many discussions of Yang-Mills theory, the definition of a covariant derivative takes the form (introducing the gauge field  $A_{\underline{a}}^{\mathcal{I}}$ )

$$\nabla_{\underline{a}} \equiv \partial_{\underline{a}} - i A_{\underline{a}}^{\mathcal{I}} \tilde{t}_{\mathcal{I}} = \partial_{\underline{a}} - i A_{\underline{a}} \quad , \quad (149)$$

where  $\tilde{t}_{\mathcal{I}}$  is any matrix representation. The parameter of gauge transformations has the form  $\lambda = \lambda^{\mathcal{A}}(x) \tilde{t}_{\mathcal{A}}$  so that the gauge transformation takes the form

$$(\nabla_{\underline{a}})' = \exp[i\lambda] \nabla_{\underline{a}} \exp[-i\lambda] \rightarrow A_{\underline{a}}' = A_{\underline{a}} - i e^{i\lambda} (\partial_{\underline{a}} e^{-i\lambda}) \quad . \quad (150)$$

The field strength is defined by

$$[\nabla_{\underline{a}}, \nabla_{\underline{b}}] = -i f_{\underline{a}\underline{b}}^{\mathcal{I}} \tilde{t}_{\mathcal{I}} \rightarrow f_{\underline{a}\underline{b}}^{\mathcal{I}} = \partial_{\underline{a}} A_{\underline{b}}^{\mathcal{I}} - \partial_{\underline{b}} A_{\underline{a}}^{\mathcal{I}} - i f_{\mathcal{J}\mathcal{K}}^{\mathcal{I}} A_{\underline{a}}^{\mathcal{J}} A_{\underline{b}}^{\mathcal{K}} . \quad (151)$$

It is interesting to re-calculate  $f_{\underline{a}\underline{b}}^{\mathcal{I}}$  using the abstract Lie algebra generator  $t_{\mathcal{A}}$  in place of  $\tilde{t}_{\mathcal{A}}$ .

$$\begin{aligned} i f_{\underline{a}\underline{b}}^{\mathcal{I}} t_{\mathcal{I}} &= [\partial_{\underline{a}} - i A_{\underline{a}}^{\mathcal{J}} t_{\mathcal{J}}, \partial_{\underline{b}} - i A_{\underline{b}}^{\mathcal{K}} t_{\mathcal{K}}] \\ &= i (\partial_{\underline{a}} A_{\underline{b}}^{\mathcal{I}} - \partial_{\underline{b}} A_{\underline{a}}^{\mathcal{I}}) t_{\mathcal{I}} - [A_{\underline{a}}^{\mathcal{J}} t_{\mathcal{J}}, A_{\underline{b}}^{\mathcal{K}} t_{\mathcal{K}}] \\ &= i (\partial_{\underline{a}} A_{\underline{b}}^{\mathcal{I}} - \partial_{\underline{b}} A_{\underline{a}}^{\mathcal{I}}) t_{\mathcal{I}} - A_{\underline{a}}^{\mathcal{J}} (t_{\mathcal{J}} A_{\underline{b}}^{\mathcal{I}}) t_{\mathcal{I}} \\ &\quad + A_{\underline{b}}^{\mathcal{J}} (t_{\mathcal{J}} A_{\underline{a}}^{\mathcal{I}}) t_{\mathcal{I}} - A_{\underline{a}}^{\mathcal{J}} A_{\underline{b}}^{\mathcal{K}} [t_{\mathcal{J}}, t_{\mathcal{K}}] . \end{aligned} \quad (152)$$

This definition of the field strength will agree with the first one if and only if the gauge field possesses a nonvanishing result under the action of the abstract Lie algebra generator  $t_{\mathcal{J}}$ ,

$$(t_{\mathcal{J}} A_{\underline{b}}^{\mathcal{I}}) = i f_{\mathcal{J}\mathcal{K}}^{\mathcal{I}} A_{\underline{b}}^{\mathcal{K}} . \quad (153)$$

The fact that this condition is satisfied is equivalent to the statement that a nonabelian gauge field is “charged” with regard to its own gauge group.

## 2.7 4D, N = 1 Supersymmetric Yang-Mills Theory

We can now bring all of the experiences of the last few sections to bare on solving the problem of finding the form of 4D, N = 1 supersymmetric Yang-Mills theory. As an ansatz, we directly commandeer the abelian solution,

$$\begin{aligned} \Gamma_{\alpha}^{\mathcal{I}} t_{\mathcal{I}} &\equiv i e^{-\bar{\Omega}^{\mathcal{I}} t_{\mathcal{I}}} \overline{D}_{\alpha} e^{\bar{\Omega}^{\mathcal{I}} t_{\mathcal{I}}} , \\ \rightarrow \Gamma_{\alpha}^{\mathcal{I}} t_{\mathcal{I}} &= i e^{-\Omega^{\mathcal{I}} t_{\mathcal{I}}} D_{\alpha} e^{\Omega^{\mathcal{I}} t_{\mathcal{I}}} , \\ \Gamma_{\underline{a}} &\equiv -i \left[ D_{\alpha} \Gamma_{\alpha} + \overline{D}_{\alpha} \Gamma_{\alpha} - i [\Gamma_{\alpha}, \Gamma_{\alpha}] \right] . \end{aligned} \quad (154)$$

These are to be regarded as definitions of the Yang-Mills connection superfield  $\Gamma_{\underline{A}}$ . Due to these definitions, we find  $F_{\alpha\beta}^{\mathcal{I}} = F_{\alpha\beta}^{\mathcal{I}} = F_{\alpha\beta}^{\mathcal{I}} = 0$ . Using these, the 4D, N = 1 superspace Yang-Mills covariant derivatives satisfy

$$\begin{aligned} [\nabla_{\alpha}, \nabla_{\beta}] &= 0 , \quad [\overline{\nabla}_{\alpha}, \overline{\nabla}_{\beta}] = 0 , \quad [\nabla_{\alpha}, \overline{\nabla}_{\alpha}] = i \nabla_{\underline{a}} , \\ [\nabla_{\alpha}, \nabla_{\underline{b}}] &= C_{\alpha\beta} \overline{W}_{\beta}^{\mathcal{I}} t_{\mathcal{I}} , \quad [\overline{\nabla}_{\alpha}, \nabla_{\underline{b}}] = C_{\alpha\beta} W_{\beta}^{\mathcal{I}} t_{\mathcal{I}} . \end{aligned}$$

$$[\nabla_{\underline{a}}, \nabla_{\underline{b}}] = -i f_{\underline{a}\underline{b}}^{\mathcal{I}} t_{\mathcal{I}} . \quad (155)$$

(The reader is warned that the nonabelian version of  $W_{\alpha}^{\mathcal{I}}$  is defined by this equation *not* the abelian expression in (135).)

With all of these results, we can now calculate the form of the nonabelian version of (141). We essentially follow the same procedure as in (90) (here we suppress the  $\mathcal{I}$ -index for simplicity)

$$\begin{aligned} \int d^6 z \left( \frac{1}{4} W^{\alpha} W_{\alpha} \right) &= \frac{1}{8} \int d^4 x \nabla^{\beta} \nabla_{\beta} W^{\alpha} W_{\alpha} \Big| \\ &= -\frac{1}{4} \int d^4 x \nabla^{\beta} [W^{\alpha} \nabla_{\beta} W_{\alpha}] \Big| \\ &= -\frac{1}{4} \int d^4 x [(\nabla^{\beta} W^{\alpha})(\nabla_{\beta} W_{\alpha}) - W^{\alpha}(\nabla^{\beta} \nabla_{\beta} W_{\alpha})] \Big| . \end{aligned} \quad (156)$$

A feature that may be a bit unexpected is that the superspace measure above was replaced by superspace Yang-Mills covariant derivatives  $d^6 z \rightarrow d^4 x \nabla^2$ . This is valid because the integrand must be a Yang-Mills invariant and thus any apparent dependence on the spinorial connection is illusory.

The definitions of component fields for the nonabelian case are obtained as for the U(1) theory with the appropriate generalizations to equation (143)  $D_{\alpha} \rightarrow \nabla_{\alpha}$ ,  $W_{\alpha} \rightarrow W_{\alpha}^{\mathcal{I}}$ ,  $\lambda_{\alpha} \rightarrow \lambda_{\alpha}^{\mathcal{I}}$ ,  $f_{\underline{a}\underline{b}} \rightarrow f_{\underline{a}\underline{b}}^{\mathcal{I}}$ ,  $d \rightarrow d^{\mathcal{I}}$ , so that

$$\begin{aligned} \nabla_{\alpha} W_{\beta}^{\mathcal{I}} \Big| &\equiv \frac{1}{2} C^{\dot{\kappa}\dot{\lambda}} [f_{\alpha\dot{\kappa}\beta\dot{\lambda}}^{\mathcal{I}} + \tilde{f}_{\alpha\dot{\kappa}\beta\dot{\lambda}}^{\mathcal{I}}] - i C_{\alpha\beta} d^{\mathcal{I}} \\ \nabla^2 W_{\alpha}^{\mathcal{I}} \Big| &\equiv -i \nabla_{\underline{a}} \bar{\lambda}^{\dot{\alpha}\mathcal{I}} . \end{aligned} \quad (157)$$

Therefore, the action quadratic in  $W_{\alpha}$  becomes

$$\begin{aligned} \int d^6 z \left( \frac{1}{4} W^{\alpha} W_{\alpha} \right) &= -\frac{1}{4} \int d^4 x \left[ \frac{1}{4} (f_{\underline{a}\underline{b}}^{\mathcal{I}} f_{\underline{a}\underline{b}}^{\mathcal{I}} + f_{\underline{a}\underline{b}}^{\mathcal{I}} \tilde{f}_{\underline{a}\underline{b}}^{\mathcal{I}}) \right. \\ &\quad \left. + i 2 \lambda^{\alpha\mathcal{I}} \nabla_{\underline{a}} \bar{\lambda}^{\dot{\alpha}\mathcal{I}} - 2 d^{\mathcal{I}} d^{\mathcal{I}} \right] . \end{aligned} \quad (158)$$

So that finally the component form of (141) is takes the form

$$\mathcal{S}_{SYM} = \int d^4 x \left[ -\frac{1}{8} f_{\underline{a}\underline{b}}^{\mathcal{I}} f_{\underline{a}\underline{b}}^{\mathcal{I}} - i \lambda^{\alpha\mathcal{I}} \nabla_{\underline{a}} \bar{\lambda}^{\dot{\alpha}\mathcal{I}} + d^{\mathcal{I}} d^{\mathcal{I}} \right] . \quad (159)$$

## 2.8 Superfield Gauge Transformations and Maurer-Cartan Forms

The main difference in the gauge transformation law of ordinary Yang-Mills theory and the superspace theory is that the latter takes a highly non-linear

form in terms of the gauge superfield  $V = V^{\mathcal{I}} t_{\mathcal{I}}$

$$(e^V)' = e^{i\bar{\Lambda}} e^V e^{-\Lambda} \quad , \quad (160)$$

since in the non-abelian case this equation cannot be simplified we must treat it with some care. The reason for this is that once we write a gauge fixing term, its gauge variation determines the ghost-antighost propagator. For infinitesimal  $\Lambda$  we have

$$\delta_G e^V = i [ \bar{\Lambda} e^V - e^V \Lambda ] \quad . \quad (161)$$

Now we have the obvious identity  $[V, e^V] = 0$ , and applying  $\delta_G$  to both sides of this equation yields

$$\begin{aligned} [\delta_G V, e^V] &= - [V, \delta_G e^V] \rightarrow \\ e^{-\frac{1}{2}V} (\delta_G V) e^{\frac{1}{2}V} - e^{\frac{1}{2}V} (\delta_G V) e^{-\frac{1}{2}V} &= - [V, e^{-\frac{1}{2}V} (\delta_G e^V) e^{-\frac{1}{2}V}] \quad , \end{aligned} \quad (162)$$

after multiplying front and back by  $\exp[-\frac{1}{2}V]$ . We next observe that (161) and (162) can be used to show

$$\begin{aligned} e^{-\frac{1}{2}V} (\delta_G e^V) e^{-\frac{1}{2}V} - e^{\frac{1}{2}V} (\delta_G V) e^{-\frac{1}{2}V} \\ = i [V, e^{-\frac{1}{2}V} \bar{\Lambda} e^{\frac{1}{2}V} - e^{\frac{1}{2}V} \Lambda e^{-\frac{1}{2}V}] \quad . \end{aligned} \quad (163)$$

At this stage it is convenient to introduce a notational device. We can use the symbol  $L_V X \equiv [V, X]$  defined to act on any two matrices  $V$  and  $X$ . Utilizing this we find

$$\begin{aligned} e^{-\frac{1}{2}V} (\delta_G V) e^{\frac{1}{2}V} &= \exp[-\frac{1}{2}L_V] (\delta_G V) \quad , \\ e^{\frac{1}{2}V} (\delta_G V) e^{-\frac{1}{2}V} &= \exp[\frac{1}{2}L_V] (\delta_G V) \quad , \\ e^{-\frac{1}{2}V} \bar{\Lambda} e^{\frac{1}{2}V} &= \exp[-\frac{1}{2}L_V] \bar{\Lambda} \quad , \\ e^{\frac{1}{2}V} \Lambda e^{-\frac{1}{2}V} &= \exp[\frac{1}{2}L_V] \Lambda \quad , \end{aligned} \quad (164)$$

where these equations can be shown to be valid by use of Taylor's theorem with respect to  $V$ . Using these in (163) then yields,

$$\begin{aligned} \sinh(\frac{1}{2}L_V) (\delta_G V) &= -i \frac{1}{2} L_V [ \exp(\frac{1}{2}L_V) \Lambda - \exp(-\frac{1}{2}L_V) \bar{\Lambda} ] \\ &= -i \frac{1}{2} L_V \sinh(\frac{1}{2}L_V) (\Lambda + \bar{\Lambda}) \\ &\quad - i \frac{1}{2} L_V \cosh(\frac{1}{2}L_V) (\Lambda - \bar{\Lambda}) \quad . \end{aligned} \quad (165)$$

As a final step, we formally divide this by  $\sinh(\frac{1}{2}L_V)$

$$\delta_G V = -i \frac{1}{2} L_V [ (\Lambda + \bar{\Lambda}) + \coth(\frac{1}{2}L_V) (\Lambda - \bar{\Lambda}) ] \quad . \quad (166)$$

Yang-Mills theories in superspace depend on the pseudoscalar pre-potential  $V$  superfields which appear “hidden inside” the spinorial superconnections as

$$\Gamma_\alpha = i e^{-\Omega} (D_\alpha e^\Omega) \rightarrow \nabla_\alpha = D_\alpha - i \Gamma_\alpha = e^{-\Omega} D_\alpha e^\Omega \quad . \quad (167)$$

(Recall also  $\Omega = \frac{1}{2}(V + iW)$ .) This final equation is meant to indicate that the factor of  $D_\alpha$  acts on everything that follows it. The operator  $\bar{\nabla}_\alpha$  is just the hermitian conjugate and  $\nabla_{\underline{a}}$  is given by  $\nabla_{\underline{a}} = -i[\nabla_\alpha, \bar{\nabla}_{\dot{\alpha}}]$ . From its definition, it should be clear that  $\Gamma_\alpha$  can be expressed as

$$\Gamma_\alpha = (D_\alpha \Omega^{\mathcal{A}}) L_{\mathcal{A}}^{\mathcal{I}} \tilde{t}_{\mathcal{I}} \quad , \quad (168)$$

for some “matrix” of functions  $L_{\mathcal{A}}^{\mathcal{I}}$ . Let us explore the significance of these objects. The rigid gauge transformation may be written in the form

$$U(\alpha) = \exp[i \alpha^{\mathcal{I}} \tilde{t}_{\mathcal{I}}] \quad , \quad (169)$$

whether in Minkowski space or superspace. If  $\tilde{t}_{\mathcal{I}}$  are the generators of some compact Lie group, then all the elements in the group may be represented by allowing the parameters  $\alpha^{\mathcal{I}}$  to take on a finite range of values. Since the parameters are allowed to vary in this range, it makes sense to consider the quantities

$$L^{\mathcal{J}} \equiv n_0 \text{tr} \left\{ \tilde{t}^{\mathcal{J}} \exp[-i \alpha^{\mathcal{I}} \tilde{t}_{\mathcal{I}}] d \exp[i \alpha^{\mathcal{I}} \tilde{t}_{\mathcal{I}}] \right\} = n_0 \text{tr} \left\{ \tilde{t}^{\mathcal{J}} U^{-1} dU \right\} \quad . \quad (170)$$

From this definition it is clear that under  $L^{\mathcal{J}}$  is invariant under the re-definition

$$U(\alpha) \rightarrow \tilde{U}(\beta) U(\alpha) \quad , \quad (171)$$

as long as  $\beta$  is a constant. However, as a consequence of the chain rule we have

$$dU = d\alpha^{\mathcal{K}} \frac{\partial U}{\partial \alpha^{\mathcal{K}}} \rightarrow L^{\mathcal{J}} = d\alpha^{\mathcal{K}} n_0 \text{tr} \left\{ \tilde{t}^{\mathcal{J}} U^{-1} \frac{\partial U}{\partial \alpha^{\mathcal{K}}} \right\} = d\alpha^{\mathcal{K}} L_{\mathcal{K}}^{\mathcal{J}} \quad . \quad (172)$$

The object denoted by  $L_{\mathcal{K}}^{\mathcal{J}}$  here is exactly the same as that in (168). However, the discussion immediately above makes it clear that this object is intrinsic to the structure of a Lie group. In fact, it is known as the left-invariant Maurer-Cartan form. Upon taking the determinant of this matrix, multiplying by the

volume element constructed from and integrating over the allowed range of  $\alpha^{\mathcal{I}}$ , we can use it to calculate the intrinsic hyper-volume of the Lie group.

Starting from the identity  $[\alpha^{\mathcal{I}} \tilde{t}_{\mathcal{I}}, U] = [\alpha, U] = 0$ , we can repeat part of our derivation of the explicit form of the  $\delta_G V$  variation,

$$\begin{aligned}
[d\alpha, U(\alpha)] &= -[\alpha, dU(\alpha)] \rightarrow \\
&U(-\tfrac{1}{2}\alpha)(d\alpha)U(\tfrac{1}{2}\alpha) - U(\tfrac{1}{2}\alpha)(d\alpha)U(-\tfrac{1}{2}\alpha) \\
&= -[\alpha, U(-\tfrac{1}{2}\alpha)(dU)U(-\tfrac{1}{2}\alpha)] \rightarrow \\
&\sinh(\tfrac{1}{2}L_\alpha)d\alpha = \tfrac{1}{2}L_\alpha U(-\tfrac{1}{2}\alpha)(dU)U(-\tfrac{1}{2}\alpha) \quad ,
\end{aligned} \tag{173}$$

after multiplying front and back by  $U(-\tfrac{1}{2}\alpha)$ . We may perform the formal ‘division’ by  $L_\alpha$  to write

$$U^{-1}dU = U(-\tfrac{1}{2}\alpha)[g_1(\tfrac{1}{2}L_\alpha)d\alpha]U(\tfrac{1}{2}\alpha) \quad , \tag{174}$$

where the function  $g_1(x) = \sinh(x)/x$ . Substituting this into (172) leads to the identification,

$$\begin{aligned}
L_{\mathcal{K}}^{\mathcal{I}}(\alpha) &= in_0 tr \left\{ [exp[\tfrac{1}{2}L_\alpha] \tilde{t}^{\mathcal{I}}] [g_1(\tfrac{1}{2}L_\alpha) \tilde{t}_{\mathcal{K}}] \right\} \\
&= in_0 tr \left\{ \tilde{t}^{\mathcal{I}} [g_2(\tfrac{1}{2}L_\alpha) \tilde{t}_{\mathcal{K}}] \right\} \quad ,
\end{aligned} \tag{175}$$

where  $g_2(x) = e^{-x} \sinh(x)/x$  and the constant  $n_0$  is chosen so that  $L_{\mathcal{K}}^{\mathcal{I}}(0) = \delta_{\mathcal{K}}^{\mathcal{I}}$ .

The invariance of  $L_{\mathcal{K}}^{\mathcal{I}}$  under (171) has a very interesting extension to the supersymmetric case in (168). The spinorial connection there is seen to be invariant under

$$e^\Omega \rightarrow e^{i\bar{\Lambda}} e^\Omega \quad , \quad D_\alpha \bar{\Lambda} = 0 \quad . \tag{176}$$

On the other hand, the superspace Yang-Mills covariant derivative in (155) is seen to transform in the usual covariant way under

$$e^\Omega \rightarrow e^\Omega e^{iK} \quad , \quad K \equiv K^{\mathcal{A}} \tilde{t}_{\mathcal{A}} \quad , \quad K^{\mathcal{A}} = (K^{\mathcal{A}})^* \quad . \tag{177}$$

Thus, we learn that the full set of transformations under which supersymmetric Yang-Mills theory is invariant consists of the transformations

$$(e^\Omega)' = e^{i\bar{\Lambda}} e^\Omega e^{iK} \quad . \tag{178}$$

Superspace Yang-Mills theories (and every other gauge theory in superspace) possesses two *distinct* local symmetry groups. One is parametrized by chiral

superfields (i.e.  $\Lambda^{\mathcal{I}}$  sometimes called the “ $\Lambda$ -gauge group”) and one by real superfields (i.e.  $K^{\mathcal{I}}$  sometimes called the “ $K$ -gauge group”). The transformations with respect to  $K^{\mathcal{I}}$  can be used to “algebraically gauge away” the purely imaginary part of  $\Omega^{\mathcal{I}}$ . Alternately, we may regard the superfields  $V^{\mathcal{I}}$  as being defined by

$$e^V \equiv e^{\Omega} (e^{\Omega})^{\dagger} . \quad (179)$$

### 2.9 The Wess-Zumino Gauge

Let us consider another aspect of these theories and for simplicity only the abelian case<sup>c</sup>. Here the gauge transformation is  $\delta_G V = -i(\Lambda - \bar{\Lambda})$  and components are defined by

$$\begin{aligned} c(x) &\equiv V \Big| , \quad \chi_{\alpha}(x) \equiv D_{\alpha} V \Big| , \quad \bar{\chi}_{\dot{\alpha}}(x) \equiv \bar{D}_{\dot{\alpha}} V \Big| , \\ M(x) &\equiv D^2 V \Big| , \quad \bar{M} \equiv \bar{D}^2 V \Big| , \quad A_{\underline{a}}(x) \equiv -\frac{1}{2} [D_{\alpha}, \bar{D}_{\dot{\alpha}}] V \Big| , \\ \lambda_{\alpha}(x) &\equiv i \bar{D}^2 D_{\alpha} V \Big| , \quad \bar{\lambda}_{\dot{\alpha}}(x) \equiv -i D^2 \bar{D}_{\dot{\alpha}} V \Big| , \quad d(x) \equiv \frac{1}{2} D^{\alpha} \bar{D}^2 D_{\alpha} V \Big| . \end{aligned} \quad (180)$$

The components of the gauge parameter can similarly be defined

$$\Lambda_1(x) \equiv \Lambda \Big| , \quad \Lambda_{\alpha} \equiv D_{\alpha} \Lambda \Big| , \quad \Lambda_2 \equiv D^2 \Lambda \Big| . \quad (181)$$

So that the components of  $V$  change as

$$\begin{aligned} \delta_G c &= -i(\Lambda_1 - \bar{\Lambda}_1) , \quad \delta_G \chi_{\alpha} = \Lambda_{\alpha} , \\ \delta_G M &= -i\Lambda_2 , \quad \delta_G A_{\underline{a}} = \frac{1}{2} \partial_{\underline{a}} (\Lambda_1 + \bar{\Lambda}_1) , \\ \delta_G \lambda_{\alpha} &= 0 , \quad \delta_G d = 0 . \end{aligned} \quad (182)$$

These equation show why at the component level one often finds the statement that the component fields  $(A_{\underline{a}}, \lambda_{\alpha}, d)$  form the 4D,  $N = 1$  gauge vector multiplet. This statement is only true in a special gauge where  $-i(\Lambda_1 - \bar{\Lambda}_1)$  is used to set  $c = 0$ ,  $\Lambda_{\alpha}$  is used to set  $\chi_{\alpha} = 0$  and  $\Lambda_2$  is used to set  $M = 0$ . This uses all of the (the parameter superfield) except  $\frac{1}{2}(\Lambda_1 + \bar{\Lambda}_1)$  which is the ordinary gauge parameter. This special gauge is called “the Wess-Zumino gauge”<sup>5</sup> or “WZ gauge.”

---

<sup>c</sup>The results for the non-abelian case are the same although the proof is more difficult.



Now let us start in this gauge i.e.  $(c = 0, \chi_\alpha = 0, M = 0)$  and note that the WZ-gauge breaks manifest supersymmetry! To see this we observe the supersymmetry variation of  $\chi_\alpha$  is

$$\delta_Q \chi_\alpha = i\epsilon_\alpha M - \bar{\epsilon}^{\dot{\alpha}} \left( \frac{1}{2} \partial_{\underline{a}} c + A_{\underline{a}} \right) . \quad (183)$$

So if we start in the gauge where  $(c, \chi_\alpha, M)$  vanish, after applying  $\delta_Q$  we find a “ $\chi_\alpha$ ” ( $\equiv -\bar{\epsilon}^{\dot{\alpha}} A_{\underline{a}}$ ). To prevent this from happening we can perform a gauge transformation with  $\Lambda_\alpha \equiv i\bar{\epsilon}^{\dot{\alpha}} A_{\underline{a}}$  but this gauge transformation cause “C” and “M” to be non-zero after applying  $\delta_G$ . These can be removed by picking  $\Lambda_2 \equiv -\bar{\epsilon}^{\dot{\alpha}} \bar{\lambda}_{\dot{\alpha}}$ . So the net result is if we start with only  $(A_{\underline{a}}, \lambda_\alpha, d)$  we must apply

$$\delta'_Q = \delta_Q + \delta_G(\tilde{\Lambda}) , \quad (184)$$

where the only components of  $\tilde{\Lambda}$  are the ones described above. Clearly this will lead to a result of the form

$$[ \delta'_Q(\epsilon_1), \delta'_Q(\epsilon_2) ] = \delta_P + \delta_G , \quad (185)$$

i.e. it must contain some dependence on the fields of  $\tilde{\Lambda}$ . This is an example of another statement that one often finds in the literature on supersymmetrical gauge theories. Namely it is often said that the closure of a supersymmetry algebra is dependent on the fields or possibly their equations of motion. This is true only if we work in the WZ gauge. On the other hand, if we always choose to work solely with the component fields  $(A_{\underline{a}}, \lambda_\alpha, d)$  this is equivalent to picking the WZ gauge. This is also why when using components, the closure of supersymmetry algebras almost always includes gauge transformations. The operator  $\delta_G(\tilde{\Lambda})$  is called “the compensating gauge transformation” and it must be introduced to maintain the WZ-gauge.

#### 2.10 4D, $N = 1$ Supersymmetric Yang-Mills-Matter Coupling

One of the most studied 4D theories is quantum electrodynamics (QED). It is of interest to construct the 4D,  $N = 1$  analog of this. Prior to the demonstration of this theory, it is convenient to explicitly introduce “covariantly chiral” superfields. The ordinary definition of a chiral superfield was via the restriction  $\bar{D}_{\dot{\alpha}} \Phi = 0$ . On the other hand, a covariant definition was introduced in (124)  $\bar{\nabla}_{\dot{\alpha}} \Phi_C = 0$  we use the label  $C$  to denote covariant chirality. In writing actions of chiral superfield matter coupled to the Yang-Mills superfields, it is always most convenient to use covariantly chiral superfields. Using the solution to

the constraints on the superspace covariant derivative, we see (working in the partial choice of gauge where  $W = 0$ )

$$\begin{aligned}\nabla_\alpha &= D_\alpha - i\Gamma_\alpha^{\mathcal{I}} t_{\mathcal{I}} = e^{\frac{1}{2}V^{\mathcal{I}} t_{\mathcal{I}}} D_\alpha e^{-\frac{1}{2}V^{\mathcal{I}} t_{\mathcal{I}}} , \\ \bar{\nabla}_{\dot{\alpha}} &= \bar{D}_{\dot{\alpha}} - i\Gamma_{\dot{\alpha}}^{\mathcal{I}} t_{\mathcal{I}} = e^{\frac{1}{2}V^{\mathcal{I}} t_{\mathcal{I}}} \bar{D}_{\dot{\alpha}} e^{-\frac{1}{2}V^{\mathcal{I}} t_{\mathcal{I}}} .\end{aligned}\tag{186}$$

In turn this has the implication

$$\bar{\nabla}_{\dot{\alpha}} \Phi_C = 0 \rightarrow \bar{D}_{\dot{\alpha}} \left( e^{-\frac{1}{2}V^{\mathcal{I}} t_{\mathcal{I}}} \Phi_C \right) = 0 ,\tag{187}$$

which clearly shows the quantity in the parenthesis is an ordinary chiral superfield. Thus we may write

$$\Phi = e^{-\frac{1}{2}V^{\mathcal{I}} t_{\mathcal{I}}} \Phi_C ,\tag{188}$$

and if we use this definition, the explicit factors involving the  $V$  superfield in (116) “disappear.” In deriving results this makes for a major simplification.

As one more step leading up the the derivation of the form of chiral superfields coupled to Yang-Mills superfields, we observe it is an exercise to show that the equations in (155) imply

$$\begin{aligned}\nabla_\alpha \bar{\nabla}^2 &= \bar{\nabla}^2 \nabla_\alpha - i\nabla_{\underline{a}} \bar{\nabla}^{\dot{\alpha}} + iW_\alpha^{\mathcal{C}} t_{\mathcal{C}} , \\ \bar{\nabla}_{\dot{\alpha}} \nabla^2 &= \nabla^2 \bar{\nabla}_{\dot{\alpha}} - i\nabla_{\underline{a}} \nabla^{\alpha} + i\bar{W}_{\dot{\alpha}}^{\mathcal{C}} t_{\mathcal{C}} , \\ \nabla^2 \bar{\nabla}^2 &= \bar{\nabla}^2 \nabla^2 - i\frac{1}{2} \nabla_{\underline{a}} (\nabla^\alpha \bar{\nabla}^{\dot{\alpha}} - \bar{\nabla}^{\dot{\alpha}} \nabla^\alpha) \\ &\quad + i(W^{\alpha\mathcal{C}} \nabla_\alpha - \bar{W}^{\dot{\alpha}\mathcal{C}} \bar{\nabla}_{\dot{\alpha}}) t_{\mathcal{C}} + i\frac{1}{2} (\nabla^\alpha W_\alpha^{\mathcal{C}}) t_{\mathcal{C}} .\end{aligned}\tag{189}$$

If we assume that the matter action  $\mathcal{S}_M$  describes Dirac fermions then it has been almost universally accepted that it must be of the form

$$\begin{aligned}\mathcal{S}_M &= \int d^4x d^2\theta d^2\bar{\theta} \left[ \bar{\Phi}_+ \Phi_+ + \bar{\Phi}_- \Phi_- \right] \\ &\quad + \left[ \int d^4x d^2\theta W(\Phi_+, \Phi_-) + \text{h.c.} \right] .\end{aligned}\tag{190}$$

In writing the action as above, we have made use of covariantly chiral superfields defined by  $\bar{\nabla}_{\dot{\alpha}} \Phi_+ = \bar{\nabla}_{\dot{\alpha}} \Phi_- = 0$ . In terms of components fields this

becomes

$$\begin{aligned}
\mathcal{S}_M = \int d^4x \Big[ & -\frac{1}{2}(\nabla^a \bar{A}_+)(\nabla_a A_+) - \frac{1}{2}(\nabla^a \bar{A}_-)(\nabla_a A_-) - i\bar{\psi}_+ \dot{\nabla}_a \psi_+^\alpha \\
& - i\bar{\psi}_- \dot{\nabla}_a \psi_-^\alpha + \bar{F}_+ F_+ + \bar{F}_- F_- \\
& - [i\lambda^{\alpha\mathcal{C}}[(t_{\mathcal{C}}\psi_{+\alpha})A_+ + (t_{\mathcal{C}}\psi_{-\alpha})A_-] + \text{h.c.}] \\
& - [\frac{1}{2}W_{+,+}(A_+, A_-)\psi_+^\alpha\psi_{+\alpha} + W_{+,+}(A_+, A_-)F_+ + \text{h.c.}] \\
& - [\frac{1}{2}W_{-,-}(A_+, A_-)\psi_-^\alpha\psi_{-\alpha} + W_{-,-}(A_+, A_-)F_- + \text{h.c.}] \\
& - [W_{+,-}(A_+, A_-)\psi_+^\alpha\psi_{-\alpha} + \text{h.c.}] \\
& + d^{\mathcal{C}}[(t_{\mathcal{C}}\bar{A}_+)A_+ + (t_{\mathcal{C}}\bar{A}_-)A_-] \Big] .
\end{aligned} \tag{191}$$

In this expression,  $W_{+,+}$  denotes differentiation of  $W$  with respect to  $A_+$ ,  $W_{-,-}$  denotes differentiation of  $W$  with respect to  $A_-$ , etc. The derivation of this result begins with the expected replacement  $\int d^8z \rightarrow \int d^4x \nabla^2 \bar{\nabla}^2$  which is followed by acting on the Kähler potential term as in (91-95). However, care must be taken with regard to ‘moving’ the derivatives past one another. This is where the commutators in (155) are useful.

Once again there are seen to be auxiliary fields present whose equations of motion are purely algebraic. The algebraic equations can be solved to express the auxiliary fields in terms of the propagating fields. This yields the following on-shell action

$$\begin{aligned}
\mathcal{S}_{on-shell} = \int d^4x \Big[ & -\frac{1}{2}(\nabla^a \bar{A}_+)(\nabla_a A_+) - \frac{1}{2}(\nabla^a \bar{A}_-)(\nabla_a A_-) \\
& - i\bar{\psi}_+ \dot{\nabla}_a \psi_+^\alpha - i\bar{\psi}_- \dot{\nabla}_a \psi_-^\alpha - U(A_+, A_-) \\
& - [i\lambda^{\alpha\mathcal{C}}[(t_{\mathcal{C}}\psi_{+\alpha})A_+ + (t_{\mathcal{C}}\psi_{-\alpha})A_-] + \text{h.c.}] \\
& - [\frac{1}{2}W_{+,+}(A_+, A_-)\psi_+^\alpha\psi_{+\alpha} + \text{h.c.}] \\
& - [\frac{1}{2}W_{-,-}(A_+, A_-)\psi_-^\alpha\psi_{-\alpha} + \text{h.c.}] \\
& - [W_{+,-}(A_+, A_-)\psi_+^\alpha\psi_{-\alpha} + \text{h.c.}] \Big] ,
\end{aligned} \tag{192}$$

where the potential for the spin zero fields takes the form

$$U(A_+, A_-) = |W_{,+}(A_+, A_-)|^2 + |W_{,-}(A_+, A_-)|^2 + \frac{3}{4}|(t_c \bar{A}_+)A_+ + (t_c \bar{A}_-)A_-|^2 \quad (193)$$

For the U(1) theory, after application of the Lie algebra generators this becomes,

$$U(A_+, A_-) = |W_{,+}(A_+, A_-)|^2 + |W_{,-}(A_+, A_-)|^2 + \frac{3}{4}g^2[|\bar{A}_+|^2 - |A_-|^2] \quad (194)$$

where we have introduced the gauge coupling constant  $g$  and used the fact that  $\Phi_-$  carries the opposite charge from that of  $\Phi_+$ . The usual mass of the electron is introduced via the choice  $W = m_e \Phi_+ \Phi_-$ . The two first terms in this potential then describe mass terms for the four spin-0 fields. Following the discussion in section 2.2, it can be seen that this action contains a Dirac field which transforms as

$$[\psi_{(\alpha)}(x)]' = \begin{pmatrix} \exp[ig\Lambda] & 0 \\ 0 & \exp[ig\bar{\Lambda}] \end{pmatrix} \begin{pmatrix} D_\alpha \Phi_+ \\ \bar{D}_{\dot{\alpha}} \bar{\Phi}_- \end{pmatrix} \quad (195)$$

under the local U(1) symmetry of the model. For the gauge group U(1), the supersymmetric Yang-Mills action in (159) has a leading term which is exactly the Maxwell action. The next term describes a massless (free for U(1) since  $tA_{\underline{a}} = 0 \rightarrow tV = 0 \rightarrow t\lambda_\alpha = 0$ ) spinor called the “photino.”

#### 2.11 Other 4D, $N = 1$ Supersymmetric Yang-Mills Terms

In the case of abelian groups or subgroups, the following superfield action is gauge invariant,

$$\mathcal{S}_{F-I} = \int d^4x d^2\theta d^2\bar{\theta} V = \int d^4x d \quad (196)$$

This is known as a Fayet-Iliopoulos term and it can become important in considerations of symmetry breaking.

In our effort to describe supersymmetric QED, we have rushed past something interesting. Notice that in order to write the supersymmetric extension of the Yang-Mills (and Maxwell) action, we had to make use of the real part of (156). It is an obvious question to ask about the imaginary part of this

quantity? The imaginary part of the action quadratic in  $W_\alpha$  becomes

$$\begin{aligned} \int d^6 z [i(\frac{1}{4} W^\alpha W_\alpha) + \text{h.c.}] &= \int d^4 x [ \frac{1}{8} \epsilon^{\underline{abcd}} f_{\underline{ab}}^{\mathcal{A}} f_{\underline{cd}}^{\mathcal{A}} + \nabla_{\underline{a}} (\lambda^\alpha \mathcal{A} \bar{\lambda}^{\dot{\alpha}} \mathcal{A}) ] \\ &\equiv \int d^4 x \partial_{\underline{a}} [ \frac{1}{48} \epsilon^{\underline{abcde}} X_{\underline{abc}}^{CS} + (\lambda^\alpha \mathcal{A} \bar{\lambda}^{\dot{\alpha}} \mathcal{A}) ] . \end{aligned} \quad (197)$$

On the last line above, we have introduced a quantity  $X_{\underline{abc}}^{CS}$  known as the ‘‘Chern-Simons’’ three form. This quantity has topological significance. Its integrated form is known as the ‘‘instanton number.’’ From the way that it appears above, it is clearly a pure surface term. However, it is also clear that in a supersymmetrical theory, as above, even surface terms are often accompanied by fermionic extensions.

One interesting feature of the Chern-Simons form above is that it is a single component of a superspace Chern-Simons form  $X_{\underline{ABC}}^{CS}$  all of whose components are given by

$$\begin{aligned} X_{\alpha\beta\gamma}^{CS} &= \frac{1}{2} [ \Gamma_{(\alpha}^{\mathcal{I}} F_{\beta\gamma)}^{\mathcal{I}} - \frac{1}{3} f_{\mathcal{I}\mathcal{J}\mathcal{K}} \Gamma_{(\alpha}^{\mathcal{I}} \Gamma_{|\beta|}^{\mathcal{J}} \Gamma_{|\gamma)}^{\mathcal{K}} ] , \\ X_{\alpha\beta\dot{\gamma}}^{CS} &= [ \Gamma_{(\alpha}^{\mathcal{I}} F_{\beta)\dot{\gamma}}^{\mathcal{I}} + A_{\dot{\gamma}}^{\mathcal{I}} F_{\alpha\beta}^{\mathcal{I}} - f_{\mathcal{I}\mathcal{J}\mathcal{K}} \Gamma_{\alpha}^{\mathcal{I}} \Gamma_{\beta}^{\mathcal{J}} A_{\dot{\gamma}}^{\mathcal{K}} ] , \\ X_{\alpha\beta\dot{\underline{c}}}^{CS} &= [ \Gamma_{(\alpha}^{\mathcal{I}} F_{\beta)\dot{\underline{c}}}^{\mathcal{I}} + \Gamma_{\dot{\underline{c}}}^{\mathcal{I}} F_{\alpha\beta}^{\mathcal{I}} - f_{\mathcal{I}\mathcal{J}\mathcal{K}} \Gamma_{\alpha}^{\mathcal{I}} \Gamma_{\beta}^{\mathcal{J}} \Gamma_{\dot{\underline{c}}}^{\mathcal{K}} ] , \\ X_{\alpha\dot{\beta}\dot{\underline{c}}}^{CS} &= [ \Gamma_{\alpha}^{\mathcal{I}} F_{\dot{\beta}\dot{\underline{c}}}^{\mathcal{I}} + A_{\dot{\underline{c}}}^{\mathcal{I}} F_{\alpha\dot{\beta}}^{\mathcal{I}} - f_{\mathcal{I}\mathcal{J}\mathcal{K}} \Gamma_{\alpha}^{\mathcal{I}} A_{\dot{\beta}}^{\mathcal{J}} \Gamma_{\dot{\underline{c}}}^{\mathcal{K}} ] , \\ X_{\alpha\dot{\underline{b}}\dot{\underline{c}}}^{CS} &= [ \Gamma_{\alpha}^{\mathcal{I}} F_{\dot{\underline{b}}\dot{\underline{c}}}^{\mathcal{I}} + \Gamma_{\dot{\underline{b}}}^{\mathcal{I}} F_{\dot{\underline{c}}\alpha}^{\mathcal{I}} - f_{\mathcal{I}\mathcal{J}\mathcal{K}} \Gamma_{\alpha}^{\mathcal{I}} \Gamma_{\dot{\underline{b}}}^{\mathcal{J}} \Gamma_{\dot{\underline{c}}}^{\mathcal{K}} ] , \\ X_{\underline{abc}}^{CS} &= \frac{1}{2} [ \Gamma_{[\underline{a}}^{\mathcal{I}} F_{\underline{bc}]}^{\mathcal{I}} - \frac{1}{3} f_{\mathcal{I}\mathcal{J}\mathcal{K}} \Gamma_{[\underline{a}}^{\mathcal{I}} \Gamma_{|\underline{b}|}^{\mathcal{J}} \Gamma_{|\underline{c}]}^{\mathcal{K}} ] . \end{aligned} \quad (198)$$

Interestingly enough, the action of (141) can also be written as

$$\mathcal{S}_{SYM} = \int d^4 x d^2 \theta d^2 \bar{\theta} \eta^{\underline{ab}} X_{\alpha\dot{\underline{a}}\dot{\underline{b}}}^{CS} . \quad (199)$$

The action that we rejected as a description of the 4D,  $N = 1$  Maxwell theory

$$\mathcal{S}(H.D.) = \int d^4 x d^2 \theta d^2 \bar{\theta} F^{\underline{ab}} F_{\underline{ab}} , \quad (200)$$

contains a useful lesson. Although we will not explicitly evaluate this in terms of component fields, the projection techniques may be used by the reader to find its components. Up until this point, we have always encountered some auxiliary fields within superfield actions. This may have led to the false impression that

all superfield actions contain auxiliary fields. This is not the case. In the example of this action it is simple to show that it contains a term of the form  $d\Box d$ , so that the  $d$  field propagates in this action.

This illustrates the point that in general an expression of the form

$$\mathcal{S} = \int d^4x \left[ d^2\theta f_1(W^\alpha) + \text{h.c.} \right] + \int d^4x d^2\theta d^2\bar{\theta} f_2(W^\alpha, \bar{W}^{\dot{\alpha}}, \nabla_{\underline{A}}) \quad , \quad (201)$$

will usually describe the propagation of the gauge field and gauginos but also the propagation of the “auxiliary”  $d$ -fields. In complete generality such an action contains these are dynamical fields. It is known, however, that there does exist a very special class of actions<sup>8</sup> of the form above for which the  $d$ -fields do not propagate. We have referred to these as “auxiliary-free” actions.

### 2.12 The Minimal 4D, $N = 1$ Supersymmetric Nonlinear $\sigma$ -model

In writing the kinetic energy term for the chiral scalar multiplet, we have taken advantage of an insight first obtained by Zumino<sup>9</sup> in calling it the Kähler potential term. An important class of theories arises by considering nonlinear  $\sigma$ -models with actions of the form

$$\mathcal{S}_\sigma = \int d^4x d^2\theta d^2\bar{\theta} K(\Phi^{\mathcal{A}}, \bar{\Phi}^{\mathcal{A}}) \quad , \quad (202)$$

where the Kähler potential  $K \neq \sum \Phi^{\mathcal{A}} \bar{\Phi}^{\mathcal{A}}$  but instead is a general function of  $\Phi^{\mathcal{A}}$  and  $\bar{\Phi}^{\mathcal{A}}$ . At the level of component by use of same technique as in (91-95) we derive

$$\begin{aligned} \mathcal{S}_\sigma = \int d^4x \left[ -\frac{1}{2} g_{\mathcal{A}\bar{\mathcal{B}}} (\mathcal{D}_{\underline{A}} \bar{A}^{\bar{\mathcal{B}}}) (\mathcal{D}_{\underline{A}} A^{\mathcal{A}}) - i g_{\mathcal{A}\bar{\mathcal{B}}} \bar{\psi}^{\dot{\alpha}\bar{\mathcal{B}}} \mathcal{D}_{\underline{A}} \psi^{\alpha\mathcal{A}} \right. \\ \left. + \frac{1}{4} \bar{\psi}^{\dot{\alpha}\mathcal{A}} \bar{\psi}_{\dot{\alpha}}^{\mathcal{B}} \psi^{\alpha\mathcal{C}} \psi_{\alpha}^{\mathcal{D}} \mathcal{R}_{\mathcal{C}\bar{\mathcal{A}}\bar{\mathcal{D}}\bar{\mathcal{B}}} \right] \quad , \end{aligned} \quad (203)$$

after elimination of the auxiliary  $F$ -fields by their equations of motion. In this expression, we have used the notations

$$\begin{aligned} g_{\mathcal{A}\bar{\mathcal{B}}} &\equiv \frac{\partial^2 K}{\partial \Phi^{\mathcal{A}} \partial \bar{\Phi}^{\bar{\mathcal{B}}}} = K_{,\mathcal{A},\bar{\mathcal{B}}} \quad , \\ \mathcal{D}_{\underline{A}} \psi^{\alpha\mathcal{A}} &\equiv \partial_{\underline{A}} \psi^{\alpha\mathcal{A}} - (\partial_{\underline{A}} \bar{A}^{\bar{\mathcal{B}}}) \Gamma_{\mathcal{B}\mathcal{C}}^{\mathcal{A}} \psi^{\alpha\mathcal{C}} \quad , \\ \Gamma_{\mathcal{B}\mathcal{C}}^{\mathcal{A}} &\equiv g^{\mathcal{A}\bar{\mathcal{D}}} K_{,\mathcal{B},\mathcal{C},\bar{\mathcal{D}}} \quad , \\ \mathcal{R}_{\mathcal{A}\bar{\mathcal{C}}\bar{\mathcal{B}}\bar{\mathcal{D}}} &\equiv K_{,\mathcal{A},\mathcal{B},\bar{\mathcal{C}},\bar{\mathcal{D}}} - g^{\mathcal{K}\bar{\mathcal{K}}} K_{,\mathcal{A},\mathcal{B},\bar{\mathcal{K}}} K_{,\bar{\mathcal{C}},\bar{\mathcal{D}},\mathcal{K}} \quad , \end{aligned} \quad (204)$$

where the quantity  $g^{\mathcal{K}\bar{\mathcal{K}}}$  is the matrix inverse to  $g_{\mathcal{K}\bar{\mathcal{K}}}$ . From the form of this component action we see that  $g_{\mathcal{K}\bar{\mathcal{K}}}$  plays the role of a metric in the space in which  $\Phi^{\mathcal{A}}$  and  $\bar{\Phi}^{\mathcal{A}}$  are the coordinates. If  $\mathcal{A} = 1, \dots, p$  then this complex  $p$ -dimensional space is a “Kähler” manifold.

### 2.13 4D, $N = 2$ Kählerian Vector Multiplet Models

As we have seen, particular collections of component fields make up  $N = 1$  supersymmetrical theories. Similarly some particular collections of  $N = 1$  superfields make up  $N = 2$  (and higher) supersymmetrical theories. It has been known for a long time that a pair  $(\Phi, V)$  can be used to provide a realization of an  $N = 2$  model. The action

$$\mathcal{S}_{Maxwell}^{N=2} = \int d^4x d^2\theta d^2\bar{\theta} \bar{\Phi}\Phi + \left[ \int d^4x d^2\theta \frac{1}{4} W^\alpha W_\alpha + \text{h.c.} \right] , \quad (205)$$

consists of the terms we have previously seen involving the propagating fields  $(A, \psi_\alpha, \lambda_\alpha, A_{\underline{a}})$  and auxiliary fields  $(F, d)$  and has a “hidden” second supersymmetry invariance. As the 4D,  $N = 1$  nonlinear  $\sigma$ -model may be regarded as a generalization of the “flat” Kähler potential  $K = \bar{\Phi}\Phi$ , it should be expected that the first term in the action above can be generalized in some way that will maintain the second supersymmetry if the Maxwell action is also modified. There is a special class of Kähler potentials for which this possible. If the  $N = 1$  Kähler potential is of the form

$$K(\Phi, \bar{\Phi}) = \Phi \left( \frac{\partial \bar{H}}{\partial \bar{\Phi}} \right) + \bar{\Phi} \left( \frac{\partial H}{\partial \Phi} \right) , \quad (206)$$

where  $H(\Phi)^{10}$  is an arbitrary function, and if the Maxwell superfield action is modified according to

$$W^\alpha W_\alpha \rightarrow \frac{\partial^2 H}{\partial \Phi \partial \bar{\Phi}} W^\alpha W_\alpha , \quad (207)$$

then the resulting theory<sup>d</sup> retains the extra hidden supersymmetry. Complex manifolds with geometries that satisfy this restricted form of the Kähler potential are now called, “special Kähler geometries.”

Using instantons in 1988, Seiberg<sup>11</sup> argued that the quantum effective action for the fundamental 4D,  $N = 2$  supersymmetric action of (205) has the function  $H(\Phi)$  of the form

$$H(\Phi) = c_0 \Phi^2 \left[ 1 + c_1 \ln \left( \frac{\Phi}{\Lambda} \right) + \dots \right] . \quad (208)$$

---

<sup>d</sup>I first gave these modifications to the  $N = 1$  superfield action above in 1984<sup>10</sup>.

In 1994, Seiberg and Witten<sup>12</sup> presented arguments (for the SU(2) theory) about how duality completely determines all constants ( $c_0, c_1, \dots$ ) and all terms in the ellipsis by use of a parametrical representation of elliptical curves. With the change of notation  $H \rightarrow i\mathcal{F}$ , the KVM models are now often referred to as the “Seiberg-Witten” effective action. It bares the same relation to the  $N = 2$  vector multiplet action in (205) as the leading term of the chiral perturbation theory action for pion physics bears to the underlying QCD action.

#### 2.14 $4D, N = 1$ Supersymmetric P-form Multiplets

The ordinary spin-1 U(1) photon  $A_{\underline{a}}$  is a part of a “simplex” of field theories. To see how the simplex arises, let us note the gauge transformation of the photon takes the form

$$\delta_G A_{\underline{a}} = \partial_{\underline{a}} \lambda(x) \equiv f_{\underline{a}} \quad , \quad (209)$$

and involves a spin-0 field  $\lambda(x)$  which in principle can be used to describe a massless state ( $\mathcal{S} = -\frac{1}{2} \int d^4x f^{\underline{a}} f_{\underline{a}}$ ). The action for the Maxwell field is  $\mathcal{S} = -\frac{1}{8} \int d^4x f^{\underline{a}\underline{b}} f_{\underline{a}\underline{b}}$ . Comparing these two suggests that there might exist another field  $b_{\underline{a}\underline{b}}$  whose gauge transformation has the form of the Maxwell field strength,

$$\delta_G b_{\underline{a}\underline{b}} = \partial_{\underline{a}} \kappa_{\underline{b}}(x) - \partial_{\underline{b}} \kappa_{\underline{a}}(x) \equiv f_{\underline{a}\underline{b}}(\kappa) \quad . \quad (210)$$

It is easy to show that the field strength for the new gauge field (called by various names such as the “axion,” “Kalb-Ramond field,” “second rank anti-symmetric tensor field,” or “skew symmetric tensor field”)  $b_{\underline{a}\underline{b}}$  is given by

$$h_{\underline{a}\underline{b}\underline{c}} = \partial_{\underline{a}} b_{\underline{b}\underline{c}} + \partial_{\underline{b}} b_{\underline{c}\underline{a}} + \partial_{\underline{c}} b_{\underline{a}\underline{b}} \equiv h_{\underline{a}\underline{b}\underline{c}}(b) \quad . \quad (211)$$

This propagation of this field can be described by an action of the form,

$$\mathcal{S}_{Axion} = \int d^4x \left[ -\frac{1}{2 \cdot 3!} h^{\underline{a}\underline{b}\underline{c}} h_{\underline{a}\underline{b}\underline{c}} \right] \quad . \quad (212)$$

However, we can now start this process all over again with a new field  $\mathcal{G}_{\underline{a}\underline{b}\underline{c}}$  and that will work also. Finally after all of that is done, we can repeat the process yet again starting with a field  $\mathcal{K}_{\underline{a}\underline{b}\underline{c}\underline{d}}$ ,

$$\delta_G \mathcal{K}_{\underline{a}\underline{b}\underline{c}\underline{d}} = \partial_{\underline{a}} \ell_{\underline{b}\underline{c}\underline{d}} - \partial_{\underline{b}} \ell_{\underline{c}\underline{d}\underline{a}} + \partial_{\underline{c}} \ell_{\underline{d}\underline{a}\underline{b}} - \partial_{\underline{d}} \ell_{\underline{a}\underline{b}\underline{c}} \equiv L_{\underline{a}\underline{b}\underline{c}\underline{d}}(\ell) \quad . \quad (213)$$

After this point the process stops. The reason is that we have “run out of indices to antisymmetrize.” We call the collection of theories that begins with  $\phi(x)$  and ends with  $\mathcal{K}_{\underline{a}\underline{b}\underline{c}\underline{d}}$  “a simplex.” It is illustrated below



Table 1: Simplex of Component x-space p-forms.

| p | p – form                   | Field Strength   | Gauge Transformation                                       |
|---|----------------------------|------------------|------------------------------------------------------------|
| 0 | $\phi$                     | $f_a$            | $\delta_G \phi = c_0$                                      |
| 1 | $A_a$                      | $f_{a\,b}$       | $\delta_G A_a = \partial_a \lambda$                        |
| 2 | $b_{a\,b}$                 | $h_{a\,b\,c}$    | $\delta_G b_{a\,b} = f_{a\,b}(\kappa)$                     |
| 3 | $\mathcal{G}_{a\,b\,c}$    | $L_{a\,b\,c\,d}$ | $\delta_G \mathcal{G}_{a\,b\,c} = h_{a\,b\,c}(\mu)$        |
| 4 | $\mathcal{K}_{a\,b\,c\,d}$ | 0                | $\delta_G \mathcal{K}_{a\,b\,c\,d} = L_{a\,b\,c\,d}(\ell)$ |

The action for each field in the simplex takes the form

$$\mathcal{S}_p \propto \int d^4x \, (\text{Field Strength})^2 \quad . \quad (214)$$

We can make this table more symmetrical in appearance if we define some additional field strengths via

$$\mathcal{G}_a \equiv \frac{1}{3!} \epsilon_a^{\,bcd} \mathcal{G}_{b\,c\,d} \quad , \quad \mathcal{K} \equiv \frac{1}{4!} \epsilon^{a\,b\,c\,d} \mathcal{K}_{a\,b\,c\,d} \quad . \quad (215)$$

Thus the first two columns may be written as

Table 2: Irreducible x-space p-form simplex.

| p | p – form        |
|---|-----------------|
| 0 | $\phi$          |
| 1 | $A_a$           |
| 2 | $b_{a\,b}$      |
| 3 | $\mathcal{G}_a$ |
| 4 | $\mathcal{K}$   |

The symmetry as we reflect this table vertically around the  $p = 2$  entry is a realization of the “Hodge star map.” In the mathematical language, each of the gauge fields is a differential p-form. The scalar is a “0-form,” the vector is a “1-form,” etc. Each field strength is “the exterior derivative” of the p-form and is itself a (p+1)-form. Each gauge transformation is the exterior derivative of a (p-1)-form. Both the gauge invariance of the field strength and the Bianchi identities that the field strengths satisfy are a consequence of the “Poincaré lemma” for differential forms.

Having observed this structure in Minkowski space, it is natural to inquire whether it is possible to reproduce this in 4D,  $N = 1$  superspace (and higher ones too). The answer turns out to be affirmative<sup>13</sup> at least for the 4D,  $N = 1$  superspace.

Table 3: Irreducible 4D, N = 1 superspace p-form simplex

| p | $\widehat{A}_p$  | $\widehat{dA}_p$                                                                                       | $\delta_G \widehat{A}_p$                                                                                      |
|---|------------------|--------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------|
| 0 | $\Phi$           | $i(\Phi - \bar{\Phi})$                                                                                 | $\delta_G \Phi = c_0$                                                                                         |
| 1 | $V$              | $W_\alpha = i\overline{D}^2 D_\alpha V$                                                                | $\delta_G V = -i(\Lambda - \bar{\Lambda})$                                                                    |
| 2 | $\varphi_\alpha$ | $G = -\frac{1}{2}(D^\alpha \varphi_\alpha + \overline{D}^{\dot{\alpha}} \bar{\varphi}_{\dot{\alpha}})$ | $\delta_G \varphi_\alpha = i\overline{D}^2 D_\alpha \mathcal{X}$                                              |
| 3 | $\widehat{V}$    | $\Pi = \overline{D}^2 \widehat{V}$                                                                     | $\delta_G \widehat{V} = -\frac{1}{2}(D^\alpha L_\alpha + \overline{D}^{\dot{\alpha}} \bar{L}_{\dot{\alpha}})$ |
| 4 | $\widehat{\Phi}$ | 0                                                                                                      | $\delta_G \widehat{\Phi} = \overline{D}^2 \widehat{M}$                                                        |

Of the superfields that propagate physical degrees of freedom,  $\Phi$ ,  $\varphi_\alpha$  and  $\widehat{\Phi}$  are chiral while  $V$  and  $\widehat{V}$  are real unconstrained superfields. Among the superfield parameters  $\Lambda$  and  $L_\alpha$  are chiral while  $\mathcal{X}$  and  $\widehat{M}$  are real unconstrained superfields. It can be seen that the field strength superfields  $W_\alpha$ ,  $G$  and  $\Pi$  satisfy,

$$\overline{D}_\alpha W_\alpha = 0 \ , \ D^2 G = 0 \ , \ G = \overline{G} \ , \ \overline{D}_\alpha \Pi = 0 \ . \quad (216)$$

The field strength  $G$  is of particular interest. As can be seen by comparing the two tables(1, 3), this field strength superfield (in analogy to  $W_\alpha$ ) contains the component level axion field strength. Due the algebraic and differential constraints satisfied by  $G$ , this supermultiplet is called the real linear multiplet. It was first introduced by Siegel<sup>14</sup> in terms of  $G$  as well as its component fields and plays a particularly important role in the theory of supergravity that arises as a limit of superstrings and heterotic strings.

The superfields  $\widehat{A}_p$  in the table immediately above are additional examples of “pre-potentials.” As such we have not seen the analogs of  $\Gamma_{\underline{A}}$  that we know exist for the super 1-form case. It turns out that these analogous quantities do exist. In the case of the Maxwell theory, we have seen how to explicitly construct the quantities  $\Gamma_{\underline{A}}$  starting from  $V$ . We should therefore expect that

$$\begin{aligned} b_{\underline{a}\underline{b}}(x) &\rightarrow B_{\underline{A}\underline{B}}(z) \ , \\ \mathcal{G}_{\underline{a}\underline{b}\underline{c}}(x) &\rightarrow \mathcal{G}_{\underline{A}\underline{B}\underline{C}}(z) \ , \\ \mathcal{K}_{\underline{a}\underline{b}\underline{c}\underline{d}}(x) &\rightarrow \mathcal{K}_{\underline{A}\underline{B}\underline{C}\underline{D}}(z) \ , \end{aligned} \quad (217)$$

where we call the superfields that appear here, “the geometrical super p-forms.” It is left as an exercise to derive these from the corresponding pre-potentials and thus derive the form of the constraints that are satisfied by the field strength

superfields that arise as

$$\begin{aligned} h_{\underline{a}\underline{b}\underline{c}}(x) &\rightarrow H_{\underline{A}\underline{B}\underline{C}}(z) \quad , \\ L_{\underline{a}\underline{b}\underline{c}\underline{d}}(x) &\rightarrow L_{\underline{A}\underline{B}\underline{C}\underline{D}}(z) \quad . \end{aligned} \quad (218)$$

The solutions for the super 3-form field strengths are

$$\begin{aligned} H_{\alpha\dot{\beta}\underline{c}} &= iC_{\alpha\gamma}C_{\dot{\alpha}\dot{\gamma}}G \quad , \\ H_{\alpha\underline{b}\underline{c}} &= iC_{\dot{\beta}\dot{\gamma}}C_{\alpha(\beta}D_{\gamma)}G \quad , \quad H_{\dot{\alpha}\underline{b}\underline{c}} = - (H_{\alpha\underline{b}\underline{c}})^* \quad , \\ H_{\underline{a}\underline{b}\underline{c}} &= \epsilon_{\underline{a}\underline{b}\underline{c}\underline{d}} [ D^{\dot{\delta}} , \overline{D}^{\dot{\delta}} ] G \quad , \end{aligned} \quad (219)$$

with all remaining components vanishing. Similarly for the super 4-form field strengths

$$\begin{aligned} L_{\alpha\beta\underline{c}\underline{d}} &= C_{\dot{\gamma}\dot{\delta}}C_{\alpha(\gamma}C_{\delta)\beta}\overline{\Pi} \quad , \quad L_{\dot{\alpha}\dot{\beta}\underline{c}\underline{d}} = - (L_{\alpha\beta\underline{c}\underline{d}})^* \quad , \\ L_{\alpha\underline{b}\underline{c}\underline{d}} &= - \epsilon_{\underline{a}\underline{b}\underline{c}\underline{d}} \overline{D}^{\dot{\alpha}}\overline{\Pi} \quad , \quad L_{\dot{\alpha}\underline{b}\underline{c}\underline{d}} = - (L_{\alpha\underline{b}\underline{c}\underline{d}})^* \quad , \\ L_{\underline{a}\underline{b}\underline{c}\underline{d}} &= i \epsilon_{\underline{a}\underline{b}\underline{c}\underline{d}} [ D^2\Pi - \overline{D}^2\overline{\Pi} ] \quad , \end{aligned} \quad (220)$$

again with all remaining field strength components vanishing.

#### 2.15 $4D$ , $N = 1$ Supersymmetric Duality and the Complex Linear Multiplet

An example of the concept of duality (there are actually several different such concepts) is provided by our re-examining some of the results in the last section, especially the theory of the massless scalar and that of the axion. In these cases,

$$\begin{aligned} p &= 0 \quad , \quad \mathcal{S}_0 = -\frac{1}{2} \int d^4x \, f^{\underline{a}} f_{\underline{a}} \quad , \quad f_{\underline{a}} = \partial_{\underline{a}}\phi \quad , \\ p &= 2 \quad , \quad \mathcal{S}_2 = \frac{1}{2} \int d^4x \, h^{\underline{a}} h_{\underline{a}} \quad , \quad h_{\underline{a}} = \frac{1}{3!} \epsilon_{\underline{a}}^{\underline{b}\underline{c}\underline{d}} h_{\underline{b}\underline{c}\underline{d}} \quad . \end{aligned} \quad (221)$$

As algebraic consequences we have

$$\begin{aligned} 0 &= \partial_{\underline{a}} f_{\underline{b}} - \partial_{\underline{b}} f_{\underline{a}} \quad , \\ 0 &= \partial^{\underline{a}} h_{\underline{a}} \quad , \end{aligned} \quad (222)$$

and we may think of these as the respective ‘‘Bianchi identities’’ for the two field strengths,  $f_{\underline{a}}$  and  $h_{\underline{a}}$ .

On the other hand, the respective equations of motion can be derived by simple variation of each action

$$\begin{aligned}\frac{\delta S_0}{\delta \phi} &= 0 \rightarrow \partial^a f_{\underline{a}} = 0 \quad , \\ \frac{\delta S_2}{\delta b_{\underline{a}\underline{b}}} &= 0 \rightarrow \partial_{\underline{a}} h_{\underline{b}} - \partial_{\underline{b}} h_{\underline{a}} = 0 \quad .\end{aligned}\tag{223}$$

So it is obvious that the roles of the equation of motion and Bianchi identity become “switched” as we go from one field to the other. We have called this type of duality “Poincaré duality”<sup>e</sup> to distinguish it from other types of “dualities.” The fact that in the presence of the equations of motion (i.e. on-shell) these two systems are exactly the same implies that they both describe a spin-0 degree of freedom.

This example is but one of many that occur within field theories. Given that these dual theories describe the same physical states, it is tempting to regard them as equivalent representations. This however, is false. In this example, it turns out that with regard to properties with respect to conformal symmetries, these theories are very different. So in general given two theories related by Poincaré duality, we must not a priori assume that they are equivalent.

We will now show that Poincaré duality possess an unexpected extension to supersymmetrical theories. First let us observe that there should be an expected extension. This follows from our discussion above taken together with the results of the previous section. There we learned that the supersymmetrical extension of the scalar multiplet is the chiral multiplet with regard to super p-forms. Similarly we saw that there is a supersymmetrical extension of the axion multiplet. Together these two facts imply that there must be a duality that relates the chiral multiplet to the axion multiplet and this is made obvious by the following table

Table 4: Duality of Chiral and Tensor Supermultiplets

| Superfield       | Bianchi identity                                       | Equation of motion                            |
|------------------|--------------------------------------------------------|-----------------------------------------------|
| chiral multiplet | $\overline{D}^2 D_\alpha (\Phi + \overline{\Phi}) = 0$ | $\overline{D}^2 (\Phi + \overline{\Phi}) = 0$ |
| axion multiplet  | $D^2 G = 0$                                            | $\overline{D}^2 D_\alpha G = 0$               |

However, there is yet another type of duality. We can observe this one by

---

<sup>e</sup>This is often called “electromagnetic duality.”

noting the following,

$$\overline{D}_\alpha \Phi = 0 \rightarrow \Phi = \overline{D}^2 \overline{U} \quad , \quad (224)$$

which follows from the algebra of the  $D$ 's. We may regard the complex superfield  $U$  as a “pre-potential” for the chiral scalar multiplet (we will make use of this in the later discussion of quantization). This complex superfield, since it is free of constraints, may be used to calculate the variation of expressions that contain  $\Phi$ . It follows that

$$\mathcal{S}_0 = \int d^4x d^2\theta d^2\bar{\theta} \overline{\Phi} \Phi \quad , \quad \& \quad \frac{\delta \mathcal{S}_0}{\delta \Phi} = 0 \rightarrow D^2 \Phi = 0 \quad , \quad (225)$$

as the equation of motion. So if we regard (224) as the “Bianchi identity” and (225) as yielding the equation of motion, a dual theory should interchange these two and this is certainly not what is contained in the table above.

This other putative dual superfield<sup>15</sup> (which we shall denote by  $\overline{\Sigma}$ ) should satisfy  $D^2 \overline{\Sigma} = 0$  as a constraint (this is called the ‘linear constraint’ as opposed to the chirality constraint). The solution to this constraint is

$$\Sigma = \overline{D}_\alpha U^\alpha \quad . \quad (226)$$

So that we see that there is a simple solution to the constraint. This solution also makes it clear that the  $\Sigma$  is a complex superfield. The next problem to face is what should be the action which describes the dynamics of this multiplet in the simplest possible manner.

The solution to this is provided by looking back at the discussion that surrounded the ordinary axion and scalar. There it can be seen that their actions (221) look almost identical with the exception of the minus sign. This suggests that a similar stratagem may apply to the supersymmetrical theory.

$$\mathcal{S}_{NM} = - \int d^4x d^2\theta d^2\bar{\theta} \overline{\Sigma} \Sigma \quad . \quad (227)$$

Let us also propose the definitions of the component fields for this multiplet as

$$\begin{aligned} B &\equiv \Sigma| \quad , \quad \overline{\zeta}_{\dot{\alpha}} \equiv \overline{D}_{\dot{\alpha}} \Sigma| \quad , \quad \rho_\alpha \equiv D_\alpha \Sigma| \quad , \quad H \equiv D^2 \Sigma| \quad , \\ p_{\underline{a}} &\equiv \overline{D}_{\dot{\alpha}} D_\alpha \Sigma| \quad , \quad \overline{\beta}_{\dot{\alpha}} \equiv \frac{1}{2} D^\alpha \overline{D}_{\dot{\alpha}} D_\alpha \Sigma| \quad . \end{aligned} \quad (228)$$

With this action, we evaluate the component fields as in previous cases to find

$$\begin{aligned} \mathcal{S}_{NM} = \int d^4x \left[ -\frac{1}{2} (\partial^\mu \overline{B})(\partial_\mu B) - i \overline{\zeta}^\alpha \partial_{\underline{a}} \zeta^{\dot{\alpha}} - \overline{H} H \right. \\ \left. + \overline{p}^{\underline{a}} p_{\underline{a}} + \beta^\alpha \rho_\alpha + \overline{\beta}^{\dot{\alpha}} \overline{p}_{\dot{\alpha}} \right] \quad . \end{aligned} \quad (229)$$

On-shell we find the superfield equation of motion  $\overline{D}_\alpha \overline{\Sigma} = 0$  or at the component level

$$\partial^d \partial_{\underline{d}} B = 0 \ , \ -i \partial_{\underline{a}} \overline{\zeta}^\alpha = 0 \ , \ H = 0 \ , \ p_{\underline{a}} = 0 \ , \ \overline{p}_\alpha = 0 \ , \ \beta_\alpha = 0 \ , \quad (230)$$

and off-shell since  $p_{\underline{a}} \neq \overline{p}_{\underline{a}}$ , the action contains 12 bosons and 12 fermions. It is now obvious that  $\Sigma$  also describes a “scalar multiplet” (i.e. one where the propagating component fields have spins no greater than 1/2) we have named this the “nonminimal scalar multiplet.” As indicated by the following table we have indeed accomplished our goal of a second unexpected duality

Table 5: Duality of Chiral and Nonminimal Scalar Multiplets

|               | Constraint                    | Equation of Motion                         |
|---------------|-------------------------------|--------------------------------------------|
| Chiral SF     | $\overline{D} \cdot \Phi = 0$ | $D^2 \Phi = 0$                             |
| Nonminimal SF | $D^2 \Sigma = 0$              | $\overline{D} \cdot \overline{\Sigma} = 0$ |

Presently, there are known two formal distinction that occur using the nonminimal multiplet.

The nonminimal multiplet can be used in conjunction with chiral multiplets to describe a second version of supersymmetric QED. One striking difference in such a construction is that the Dirac spinor contained in such a model is defined by

$$\psi_{(\alpha)}(x) \equiv \left( \begin{array}{c} D_\alpha \Phi \\ \overline{D}_\alpha \Sigma \end{array} \right) \ , \quad (231)$$

which transforms under the U(1) symmetry as

$$\left[ \psi_{(\alpha)}(x) \right]' = \exp[ig\Lambda] \psi_{(\alpha)}(x) \ . \quad (232)$$

The difference between this and (195) is clear.

In general, if one considers an action of the form

$$\mathcal{S}_{C-H.D.} = \int d^4x d^2\theta d^2\bar{\theta} f\left(\Phi, \overline{\Phi}, D_{\underline{A}}\right) \ , \quad (233)$$

it is usually the case that such an expression has the consequence

$$\frac{\delta \mathcal{S}_{C-H.D.}}{\delta F} = 0 \ , \ \rightarrow \text{dynamical equation for } F \ . \quad (234)$$

On the other hand, it has been shown that it is very simple to construct a large class of auxiliary-free actions<sup>16</sup> of the form

$$\mathcal{S}_{CNM-H.D.} = \int d^4x d^2\theta d^2\bar{\theta} f\left(\Phi, \bar{\Phi}, \Sigma, \bar{\Sigma}, D_{\underline{A}}\right) \quad , \quad (235)$$

such that *none* of the auxiliary fields propagate even though the physical fields can possess arbitrarily high order derivatives. We thus have a second example of a 4D, N = 1 supersymmetric system (c.f. comment below (201)). These two sets of observations when taken together show that there are, at least at the level of classical actions, theories with fields of spin less than or equal to one with manifest supersymmetry, higher derivative interaction for propagating fields and *no* propagating auxiliary fields. The significance of this as well as the observation of the preceding paragraph are still to be investigated.

We end this lecture with a slight warning. Although we have met many, many 4D, N = 1 supersymmetrical multiplets. The ones we have seen here are by no means exhaustive. There are many more such multiplets that have appeared only in the research literature. The reasons for the existence of the plethora of 4D, N = 1 superfield representations remains a mystery. Perhaps a more complete theory will rule out the use of many of these other representations.

### 3 Third Lecture: A Non-Supersymmetric Interlude on the Lie Algebraic Origin of Gravitational Theories

#### *Introduction*

Also while I was a graduate student, I acquired a book entitled “Gravitation and Cosmology: Principles and Applications of The General Theory of Relativity” by Steven Weinberg. Among the comments in the preface I read,

*I found that in most textbooks geometrical ideas were given a starring role, so that a student...would come away with the impression that this had something to do with the fact that space-time is a Riemannian manifold.*

*Of course, this was Einstein’s point of view, and his preeminent genius necessarily shapes our understanding of the theory he created. However, I believe that the geometrical approach has driven a wedge between general relativity and the theory of elementary particles.*

At the time this had a profound impact on my thinking about the theory of general relativity. It suggested that there should exist a way to understand theories of gravitation that does not involve geometry. I also believe that this statement stands as a warning that should be recalled in this present epoch of “eka-general relativity” (i.e. strings, superstrings, heterotic strings, D-branes, D-p-branes, M-theory, F-theory and whatever future developments spring forth from the imagination of eka-relativists).

If we are to look elsewhere for the origins of theories of gravitation, the first question becomes, “What will replace Riemannian geometry?” The conclusion I came to is that within the confines of the relativistic field theory formulation of particle physics, the principle of gauge invariance is exquisitely suited to act as a logical foundation upon which to also anchor theories of gravitation. If this view is accepted, then we are logically freed from the confines of ‘geometry.’

The implementation of the a gauge-theory based view of theories of gravitation hinges on being able to identify a Lie algebra which may be gauged. This identification has been a question approached by authors too numerous to list long before I began my efforts in this direction. However, in the late seventies during my collaboration with Warren Siegel<sup>17</sup>, I believe a solution to this question was unearthed that is the simplest of which I am aware and seems to be correct. The development of this particular approach can be found in a number of our papers from that period.

It was the theory of supergravity and the struggle to reach its comprehen-



sive and complete mathematically rigorous formulation in superspace that led to the formalism that I will discuss in this lecture. Since the geometry and pre-geometry of curved Wess-Zumino superspace was not so well studied in those early days of supergravity (and to a large extent even now), we were forced to introduce the usual supergeometrical structures only as devices of convenience. It was the gauge-theory structure that was initially the only clear signpost. We thus unintentionally (and unconsciously) became Feynman's<sup>18</sup> "Venutian field theorists" except that we were not located on the planet Venus but in a place called, "Superspace."

Finally, I believe that the hard won lessons of superspace supergravity have been largely ignored in the effort of the eka-relativists. The complete superfield structure of superspace supergravity has a layer of complication known as the "pre-potential formulation" (upon which I shall barely touch in these lectures) that is totally unexpected. A more complicated theory that contains supergravity must be able to successfully reproduce this structure. This is an updated version of the correspondence principle from quantum mechanics. I have seen little sign of this restriction being fulfilled in many works beyond supergravity with the possible exception of Zwiebach's approach<sup>19</sup> to bosonic string field theory.

## Lecture

### 3.1 Explicit Lie-Algebraic Basis of Gravitation

We want to treat the symmetries of general relativity as *similarly* as possible as are the symmetries treated in Yang-Mills theory. This leads directly to the issue of what operators should play the role of  $t_{\mathcal{I}}$ ? Let us propose that the Lie algebra generators of the gauge group of all 4D gravitational theories includes the follow set of operators

$$\left( -i\partial_{\underline{m}}, \mathcal{M}_{\alpha}{}^{\gamma}, \overline{\mathcal{M}}_{\alpha}{}^{\dot{\gamma}} \right) \quad , \quad (236)$$

acting on fields.

The first of these is recognizable as the momentum generator from elementary first quantized theories, i.e.  $-i\partial_{\underline{m}} = P_{\underline{m}}$ . Accordingly we know that

$$[P_{\underline{m}}, P_{\underline{n}}] = 0 \quad . \quad (237)$$

To understand the significance of  $\mathcal{M}_{\alpha}{}^{\gamma}$  and  $\overline{\mathcal{M}}_{\alpha}{}^{\dot{\gamma}}$  we need to recall a result from our first lecture. There we noted (8) that there exist an infinite number of sets

of matrices whose commutator algebra is isomorphic to that of the “orbital angular momentum operator”  $L_{\underline{m}\underline{n}}$  of elementary first quantized theories. The idea is to treat the sets of matrices  $\tilde{S}_{\underline{m}\underline{n}}$  in exactly the same manner as the sets  $\tilde{t}_{\mathcal{I}}$  in a Yang-Mills theory. This is done by recognizing that  $\mathcal{M}_{\underline{a}\underline{b}}$  is the analog of  $t_{\mathcal{I}}$  in a Yang-Mills theory.

Since  $\mathcal{M}_{\underline{a}\underline{b}}$  has two “vector” indices, we can use it to define two other abstract Lie algebra generators  $\mathcal{M}_{\alpha\beta}$  and  $\overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}$  via the definition

$$\mathcal{M}_{\underline{a}\underline{b}} = \left[ C_{\alpha\dot{\beta}} \mathcal{M}_{\alpha\beta} + C_{\alpha\beta} \overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}} \right] . \quad (238)$$

The generator  $\mathcal{M}_{\underline{a}\underline{b}}$  may be thought of as the “generator of spin angular momentum.” The quantities  $\mathcal{M}_{\alpha\beta}$  and  $\overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}$  may be roughly regarded as the “right-handed spin angular momentum generator” and “left-handed spin angular momentum generator” respectively (c.f. equation (43)). We recall that any theory involving spinors possesses an intrinsic definition of handedness related to the chirality of the spinors. The spin angular momentum commutes with ordinary momentum thus

$$[\mathcal{M}_{\alpha\beta}, P_{\underline{m}}] = 0 , \quad [\overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}, P_{\underline{m}}] = 0 . \quad (239)$$

Since  $[P_{\underline{m}}, x^{\underline{n}}] = -i\delta_{\underline{m}}^{\underline{n}}$ , it follows that

$$[\mathcal{M}_{\alpha\beta}, x^{\underline{m}}] = 0 , \quad [\overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}, x^{\underline{m}}] = 0 . \quad (240)$$

But wait, isn’t  $x^{\underline{m}}$  a vector? After all for a photon  $A_{\underline{a}}(x)$  we can use the definition of  $\mathcal{M}_{\underline{a}\underline{b}}$  and the result of equation (9) to show

$$[\mathcal{M}_{\alpha}^{\beta}, A_{\underline{c}}] = \delta_{\gamma}^{\beta} A_{\alpha\dot{\gamma}} - \frac{1}{2}\delta_{\alpha}^{\beta} A_{\underline{c}} , \quad [\overline{\mathcal{M}}_{\dot{\alpha}}^{\dot{\beta}}, A_{\underline{c}}] = \delta_{\dot{\gamma}}^{\dot{\beta}} A_{\gamma\dot{\alpha}} - \frac{1}{2}\delta_{\dot{\alpha}}^{\dot{\beta}} A_{\underline{c}} . \quad (241)$$

This seems to imply a contradiction!

The way out of this is to recognize that there are two distinct types of “vectors.” We will call vectors like  $P_{\underline{m}}$  “curved vectors” and henceforth will denote such quantities by underlined Latin letters  $\underline{m}, \underline{n}, \dots$  to the end of the alphabet. We will call vectors like  $A_{\underline{a}}$  “flat vectors” and henceforth will denote such quantities by underlined Latin letters  $\underline{a}, \underline{b}, \dots, \underline{\ell}$ . This is sometimes called the “early-late” convention. In our approach *all matter fields possess only flat Lorentz indices*. The realization of  $\mathcal{M}_{\alpha\beta}$  and  $\overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}$  on  $A_{\underline{a}}$  can also be used in two other ways.

We know that given two spinors  $\psi_{\alpha}(x)$  and  $\chi_{\dot{\gamma}}(x)$ , their product  $\psi_{\gamma}\chi_{\dot{\gamma}}$  should transform like  $A_{\underline{c}}$ ,

$$\mathcal{M}_{\alpha}^{\beta}(\psi_{\gamma}\chi_{\dot{\gamma}}) = \delta_{\gamma}^{\beta}(\psi_{\alpha}\chi_{\dot{\gamma}}) - \frac{1}{2}\delta_{\alpha}^{\beta}(\psi_{\gamma}\chi_{\dot{\gamma}}) , \quad (242)$$

or

$$\left( \mathcal{M}_\alpha^\beta \psi_\gamma \right) \chi_{\dot{\gamma}} + \psi_\gamma \left( \mathcal{M}_\alpha^\beta \chi_{\dot{\gamma}} \right) = \delta_\gamma^\beta \left( \psi_\alpha \chi_{\dot{\gamma}} \right) - \frac{1}{2} \delta_\alpha^\beta \left( \psi_\gamma \chi_{\dot{\gamma}} \right) , \quad (243)$$

where on the second line we have used the Leibniz property. The solution we require to this is simply

$$\mathcal{M}_\alpha^\beta \psi_\gamma = \delta_\gamma^\beta \psi_\alpha - \frac{1}{2} \delta_\alpha^\beta \psi_\gamma , \quad \mathcal{M}_\alpha^\beta \chi_{\dot{\gamma}} = 0 . \quad (244)$$

Upon complex conjugation these take the forms

$$\overline{\mathcal{M}}_\alpha^{\dot{\beta}} \bar{\psi}_{\dot{\gamma}} = \delta_{\dot{\gamma}}^{\dot{\beta}} \bar{\psi}_\alpha - \frac{1}{2} \delta_\alpha^{\dot{\beta}} \bar{\psi}_{\dot{\gamma}} , \quad \overline{\mathcal{M}}_\alpha^{\dot{\beta}} \bar{\chi}_{\dot{\gamma}} = 0 . \quad (245)$$

With the results of (224) and (225) in hand, it is easy to derive the commutator algebra of  $\mathcal{M}_{\alpha\beta}$  and  $\overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}$ . These generators are found to satisfy

$$[ \mathcal{M}_\alpha^\beta , \mathcal{M}_\gamma^\delta ] = \delta_\gamma^\beta \mathcal{M}_\alpha^\delta - \delta_\alpha^\delta \mathcal{M}_\gamma^\beta , \quad (246)$$

$$[ \mathcal{M}_\alpha^\beta , \overline{\mathcal{M}}_{\dot{\gamma}}^{\dot{\delta}} ] = 0 , \quad (247)$$

$$[ \overline{\mathcal{M}}_{\dot{\alpha}}^{\dot{\beta}} , \overline{\mathcal{M}}_{\dot{\gamma}}^{\dot{\delta}} ] = \delta_{\dot{\gamma}}^{\dot{\delta}} \overline{\mathcal{M}}_{\dot{\alpha}}^{\dot{\beta}} - \delta_\alpha^{\dot{\delta}} \overline{\mathcal{M}}_{\dot{\gamma}}^{\dot{\beta}} . \quad (248)$$

(Given these and the definition in (238), we can always “re-construct” the commutator algebra of  $\mathcal{M}_{\underline{a}\underline{b}}$ .)

Using equations (246), (247) and (248) we have a complete definition of a Lie algebra for the set of operators in (238). We now wish to “gauge” the algebra in (236) as can be done for any Lie group for a Yang-Mills theory. For this purpose we introduce local parameters for our gauge group;  $\xi^{\underline{m}}(x)$ ,  $\lambda_\alpha^\beta(x)$  and  $\bar{\lambda}_\alpha^{\dot{\beta}}(x)$ . For the infinitesimal parameter of the gauge transformation we define

$$\begin{aligned} K &\equiv i \xi^{\underline{m}} P_{\underline{m}} + \lambda_\alpha^\beta(x) \mathcal{M}_{\beta}^\alpha + \bar{\lambda}_\alpha^{\dot{\beta}}(x) \overline{\mathcal{M}}_{\dot{\beta}}^{\dot{\alpha}} \\ &= \xi^{\underline{m}} \partial_{\underline{m}} + \lambda_\alpha^\beta(x) \mathcal{M}_{\beta}^\alpha + \bar{\lambda}_\alpha^{\dot{\beta}}(x) \overline{\mathcal{M}}_{\dot{\beta}}^{\dot{\alpha}} . \end{aligned} \quad (249)$$

The quantity  $K$  is a local vector in the Lie algebra. The set of all such possible local vectors is closed under commutation. If

$$\begin{aligned} K(\xi_1, \lambda_1, \bar{\lambda}_1) &= \xi_1^{\underline{m}} \partial_{\underline{m}} + \lambda_{1\alpha}^\beta(x) \mathcal{M}_{\beta}^\alpha + \bar{\lambda}_{1\alpha}^{\dot{\beta}}(x) \overline{\mathcal{M}}_{\dot{\beta}}^{\dot{\alpha}} = K_1 , \\ K(\xi_2, \lambda_2, \bar{\lambda}_2) &= \xi_2^{\underline{m}} \partial_{\underline{m}} + \lambda_{2\alpha}^\beta(x) \mathcal{M}_{\beta}^\alpha + \bar{\lambda}_{2\alpha}^{\dot{\beta}}(x) \overline{\mathcal{M}}_{\dot{\beta}}^{\dot{\alpha}} = K_2 , \end{aligned} \quad (250)$$

then there exists (with some restriction regarding the properties of the derivatives of the local parameters)  $\xi_3^{\underline{m}}$ ,  $\lambda_{3\alpha}^\beta$  and  $\bar{\lambda}_{3\alpha}^{\dot{\beta}}$  such that

$$[ K(\xi_1, \lambda_1, \bar{\lambda}_1) , K(\xi_2, \lambda_2, \bar{\lambda}_2) ] = K(\xi_3, \lambda_3, \bar{\lambda}_3) . \quad (251)$$

Furthermore, a Jacobi identity is satisfied

$$[[K_1, K_2], K_3] + [[K_2, K_3], K_1] + [[K_3, K_1], K_2] = 0 \quad . \quad (252)$$

### 3.2 Gravitational Covariant Derivative and Gauge Transformations

This local Lie algebra vector space is *exactly* like the corresponding structure in a Yang-Mills theory. Therefore, we can define gauge transformation acting on scalar, spinors or vector matter fields by

$$\begin{aligned} [\phi(x)]' &\equiv e^K \phi(x) = e^K \phi(x) e^{-K} \quad , \\ [\psi_\alpha(x)]' &\equiv e^K \psi_\alpha(x) \quad , \\ [\chi_\alpha(x)]' &\equiv e^K \chi_\alpha(x) \quad , \\ [A_{\underline{a}}(x)]' &\equiv e^K A_{\underline{a}}(x) \quad . \end{aligned} \quad (253)$$

Since we have chosen to represent the symmetries realized on any field as in (253), this has an interesting consequence. In more conventional treatments of general coordinate transformations, equations such as  $\tilde{x} = f(x)$  are used to describe such symmetries. Not all such transformations can be represented using (253). Only transformation that can be continuously connected to the identity have such representations. In the language of gauge theories, these are “small gauge transformations.” Transformation that do not possess this property (e.g.  $\tilde{x} = 1/x$ ) are in the language of gauge theories known as “large gauge transformations.”

We require a covariant derivative with respect to this gauge group and this can be introduced according to

$$\nabla_{\underline{a}} \equiv e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}} + \frac{1}{2} \omega_{\underline{a}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} \quad , \quad (254)$$

that includes the gauge fields  $e_{\underline{a}}^{\underline{m}}(x)$  and  $\omega_{\underline{a}\underline{c}\underline{d}}(x)$  with Lie-algebraic gauge transformations of the form

$$\left( \nabla_{\underline{a}} \right)' = e^K \nabla_{\underline{a}} e^{-K} \quad , \quad K \equiv K^{\underline{m}} \partial_{\underline{m}} + \frac{1}{2} K_{\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} \quad , \quad (255)$$

in terms of the local parameters  $K^{\underline{m}}(x)$  and  $K^{\underline{c}\underline{d}}(x)$ . (Since  $\omega_{\underline{a}\underline{c}\underline{d}} = -\omega_{\underline{a}\underline{d}\underline{c}}$ , we may also write  $\omega_{\underline{a}\underline{c}\underline{d}} = C_{\underline{\gamma}\underline{\delta}}^{\underline{\gamma}} \omega_{\underline{a}\underline{\gamma}\underline{\delta}} + C_{\underline{\gamma}\underline{\delta}}^{\underline{\delta}} \omega_{\underline{a}\underline{\gamma}\underline{\delta}}$ ).

There are several points to note regarding this definition.

- (a.) Since  $P_{\underline{m}}$  is one of the “generators,” this covariant derivative does not possess a part that is independent of the Lie algebra generators.
- (b.) The operator  $\nabla_{\underline{a}}$  is a “flat vector”. In particular,  

$$[\mathcal{M}_{\underline{a}\underline{b}}, \nabla_{\underline{c}}] = (\eta_{\underline{a}\underline{c}}\delta_{\underline{b}}^{\underline{d}} - \eta_{\underline{b}\underline{c}}\delta_{\underline{a}}^{\underline{d}})\nabla_{\underline{d}} \ ,$$
and this allows us to easily write actions for spinors in the presence of the  $K$ -gauge group (i.e. a minimal coupling procedure is of the form  $\bar{\psi}^{\alpha}\partial_{\underline{a}}\psi^{\alpha} \rightarrow \bar{\psi}^{\alpha}\nabla_{\underline{a}}\psi^{\alpha}$ ).

The gauge field for the translation generator  $P_{\underline{m}}$  is  $e_{\underline{a}}^{\underline{m}}(x)$  and (as we shall see) this corresponds to the field that is usually called the “inverse vierbein” or “inverse tetrad” in geometrical approaches to general relativity. The gauge field for the spin angular momentum generator  $\mathcal{M}_{\underline{a}\underline{b}}$  is  $\omega_{\underline{a}\underline{c}\underline{d}}(x)$  and (once again we shall see) this corresponds to the field that is called “the spin-connection<sup>f</sup>.” Since the  $K$ -gauge transformation law of  $\nabla_{\underline{a}}$  is exact as in a Yang-Mills theory, it follows that the transformation law of a covariant derivative of any matter field also takes the same form as the transformation law of the field itself. For example for a scalar field

$$\left[\nabla_{\underline{a}}\phi(x)\right]' = e^K\nabla_{\underline{a}}\phi(x) \ . \quad (256)$$

If we start with a Lagrangian

$$\mathcal{L} = \mathcal{L}(\phi, \partial_{\underline{a}}\phi) \ , \quad (257)$$

which does not possess the local symmetries generated by the  $K$  operator, it follows that a new Lagrangian obtained by the minimal coupling replacement  $\partial_{\underline{a}} \rightarrow \nabla_{\underline{a}}$  has the property

$$\left[\mathcal{L}(\phi, \nabla_{\underline{a}}\phi)\right]' = e^K\mathcal{L}(\phi, \nabla_{\underline{a}}\phi) = e^{\xi^{\underline{m}}\partial_{\underline{m}}}\mathcal{L}(\phi, \nabla_{\underline{a}}\phi) \ . \quad (258)$$

The transformation law above is exactly the same as for any scalar field.

At this point the analogy with Yang-Mills theory seems to break down. If we look back at Yang-Mills theory then analogously,

$$\begin{aligned} \left[\phi\right]' &= e^{i\lambda}\phi \ , \quad \left[\nabla_{\underline{a}}\right]' = e^{i\lambda}\nabla_{\underline{a}}e^{-i\lambda} \ , \\ \left[\mathcal{L}(\phi, \nabla_{\underline{a}}\phi)\right]' &= \mathcal{L}(\phi, \nabla_{\underline{a}}\phi) \ . \end{aligned} \quad (259)$$

---

<sup>f</sup>We must take caution to note that the spin-connection is *not* the usual Christoffel connection.

The difference can be seen in the operatorial pre-factor that appears in (258) that does not appear in (259). Thus, it at first seems to be impossible to write an action which possesses the local symmetries generated by  $K$ . We will come back this shortly.

Let us turn our attention instead to the infinitesimal gauge field transformation laws implied by (255).

$$\begin{aligned} \left[ \nabla_{\underline{a}} \right]' &= \nabla_{\underline{a}} + [K, \nabla_{\underline{a}}] \rightarrow \\ \delta_K \nabla_{\underline{a}} &\equiv \left[ \nabla_{\underline{a}} \right]' - \nabla_{\underline{a}} = [K, \nabla_{\underline{a}}] . \end{aligned} \quad (260)$$

This final formula looks exactly like its Yang-Mills analog. However, it obscures the details of what is happening to the  $K$ -symmetry gauge field variables. After a straightforward calculation these are found to be

$$\begin{aligned} \delta_K e_{\underline{a}}^{\underline{m}} &= \xi_{\underline{n}} \partial_{\underline{n}} e_{\underline{a}}^{\underline{m}} - e_{\underline{a}}^{\underline{n}} \partial_{\underline{n}} \xi_{\underline{m}} - \lambda_{\underline{a}}^{\underline{b}} e_{\underline{b}}^{\underline{m}} , \\ \delta_K \omega_{\underline{a}\underline{b}}^{\underline{c}} &= \xi_{\underline{n}} \partial_{\underline{n}} \omega_{\underline{a}\underline{b}}^{\underline{c}} - e_{\underline{a}}^{\underline{n}} \partial_{\underline{n}} \lambda_{\underline{b}}^{\underline{c}} - \lambda_{\underline{a}}^{\underline{d}} \omega_{\underline{d}\underline{b}}^{\underline{c}} \\ &\quad - \lambda_{\underline{b}}^{\underline{d}} \omega_{\underline{a}\underline{d}}^{\underline{c}} + \lambda_{\underline{d}}^{\underline{c}} \omega_{\underline{a}\underline{b}}^{\underline{d}} . \end{aligned} \quad (261)$$

These transformation laws give us confidence that we have correctly identified the Lie algebra that lies behind gravitational theories. The gauge algebraic transformation law of (255) unambiguously leads to these results. If we set all the terms proportional to the spin angular momentum generators to zero, then the result above has been derived with no assumptions other than the fact that the exponential of the the operator  $iK^{\underline{m}}(x)P_{\underline{m}}$  must be well defined. If we restrict  $K^{\underline{m}}(x)$  to be a constant, then the basic Lie algebraic structure of gravitational theories appears via the usual translation generator of elementary quantum theory.

The commutator of two of these gravitationally covariant derivatives generate field strengths  $t_{\underline{a}\underline{b}\underline{c}}$  and  $r_{\underline{a}\underline{b}\underline{c}\underline{d}}$

$$\left[ \nabla_{\underline{a}}, \nabla_{\underline{b}} \right] = t_{\underline{a}\underline{b}}^{\underline{c}} \nabla_{\underline{c}} + \frac{1}{2} r_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} . \quad (262)$$

Let us note that it would be just as natural to define the field strengths via

$$\left[ \nabla_{\underline{a}}, \nabla_{\underline{b}} \right] = i \mathcal{F}_{\underline{a}\underline{b}}^{\underline{m}} P_{\underline{m}} + \frac{1}{2} \mathcal{F}_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} . \quad (263)$$

However, we have the following steps.

$$\nabla_{\underline{a}} = i e_{\underline{a}}^{\underline{m}} P_{\underline{m}} + \frac{1}{2} \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \rightarrow e_{\underline{a}}^{\underline{m}} P_{\underline{m}} = -i \left( \nabla_{\underline{a}} - \frac{1}{2} \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \right) . \quad (264)$$

The gauge field  $e_{\underline{a}}^{\underline{m}}(x)$  is a  $4 \times 4$  matrix, so we can define its inverse  $e_{\underline{m}}^{\underline{a}}(x)$  (note that it is the location of the curved versus flat vector indices that distinguishes one from the other) and use it via

$$P_{\underline{m}} = -i e_{\underline{m}}^{\underline{a}} (\nabla_{\underline{a}} - \frac{1}{2} \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}}) \quad . \quad (265)$$

This last result may be substituted into the expression containing  $\mathcal{F}_{\underline{a}\underline{b}}^{\underline{m}}$  and  $\mathcal{F}_{\underline{a}\underline{b}\underline{c}}^{\underline{d}}$  to reach our initial definition.

The field strengths are explicitly given by

$$\begin{aligned} t_{\underline{a}\underline{b}\underline{c}} &= c_{\underline{a}\underline{b}\underline{c}} - \omega_{\underline{a}\underline{b}\underline{c}} + \omega_{\underline{b}\underline{a}\underline{c}} \quad , \\ r_{\underline{a}\underline{b}\underline{c}\underline{d}} &= e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}} \omega_{\underline{b}\underline{d}\underline{c}} - e_{\underline{b}}^{\underline{m}} \partial_{\underline{m}} \omega_{\underline{a}\underline{d}\underline{c}} - c_{\underline{a}\underline{b}}^{\underline{e}} \omega_{\underline{a}\underline{d}\underline{e}} \\ &\quad + \omega_{\underline{a}\underline{c}}^{\underline{e}} \omega_{\underline{b}\underline{e}\underline{d}} - \omega_{\underline{b}\underline{c}}^{\underline{e}} \omega_{\underline{a}\underline{e}\underline{d}} \quad , \end{aligned} \quad (266)$$

with the anholonomy  $(c_{\underline{a}\underline{b}}^{\underline{c}})$  defined by

$$c_{\underline{a}\underline{b}}^{\underline{c}} \equiv \left[ e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}} e_{\underline{b}}^{\underline{n}} - e_{\underline{b}}^{\underline{m}} \partial_{\underline{m}} e_{\underline{a}}^{\underline{n}} \right] e_{\underline{n}}^{\underline{c}} \quad . \quad (267)$$

Due to its definition, the anholonomy satisfies the identity

$$e_{[\underline{a}} c_{\underline{b}\underline{c}]}^{\underline{d}} - c_{[\underline{a}\underline{b}]}^{\underline{k}} c_{\underline{k}\underline{c}]}^{\underline{d}} = 0 \quad , \quad (268)$$

where  $e_{\underline{a}} \equiv e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}}$ .

The two field strengths  $t_{\underline{a}\underline{b}\underline{c}}$  and  $r_{\underline{a}\underline{b}\underline{c}\underline{d}}$  are well known from the study of differential geometry. The foremost is called the torsion tensor and the latter is the Riemann curvature tensor. Recall in the discussion of the supersymmetric Maxwell theory, the fact that a gauge superfield appeared *solely* algebraically in the definition of a field strength meant that it could be used to set the field strength to zero identically. It is natural to ask if this can be repeated?

Before we answer this, let's find out how  $t_{\underline{a}\underline{b}\underline{c}}$  and  $r_{\underline{a}\underline{b}\underline{c}\underline{d}}$  behave under arbitrary variations of the gauge fields  $e_{\underline{a}}^{\underline{m}}$  and  $\omega_{\underline{a}\underline{c}\underline{d}}$ . The results

$$\begin{aligned} \nabla_{\underline{a}} &= e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}} + \frac{1}{2} \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \rightarrow \\ \delta \nabla_{\underline{a}} &= \delta e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}} + \frac{1}{2} \delta \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \\ &= \delta e_{\underline{a}}^{\underline{m}} e_{\underline{m}}^{\underline{b}} e_{\underline{b}}^{\underline{n}} \partial_{\underline{n}} + \frac{1}{2} \delta \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \\ &\equiv h_{\underline{a}}^{\underline{b}} e_{\underline{b}}^{\underline{n}} \partial_{\underline{n}} + \frac{1}{2} \delta \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \\ &= h_{\underline{a}}^{\underline{b}} (\nabla_{\underline{b}} - \frac{1}{2} \omega_{\underline{b}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}}) + \frac{1}{2} \delta \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \\ &\equiv h_{\underline{a}}^{\underline{b}} \nabla_{\underline{b}} + \frac{1}{2} h_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}} \quad , \end{aligned} \quad (269)$$

imply that the variation of the gravitationally covariant derivative is simply expressed in terms of the two quantities  $h_{\underline{a}}^{\underline{b}} = \delta e_{\underline{a}}^{\underline{m}} e_{\underline{m}}^{\underline{b}}$  and  $h_{\underline{a}\underline{d}\underline{e}} = \delta \omega_{\underline{a}\underline{d}\underline{e}} - h_{\underline{a}}^{\underline{b}} \omega_{\underline{b}\underline{d}\underline{e}}$ .

By applying a variation operator starting at

$$\begin{aligned} \delta \left( \left[ \nabla_{\underline{a}}, \nabla_{\underline{b}} \right] \right) &= \delta t_{\underline{a}\underline{b}}^{\underline{c}} \nabla_{\underline{c}} + t_{\underline{a}\underline{b}}^{\underline{c}} \delta \nabla_{\underline{c}} + \delta \frac{1}{2} r_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} \rightarrow \\ \left[ \delta \nabla_{\underline{a}}, \nabla_{\underline{b}} \right] + \left[ \nabla_{\underline{a}}, \delta \nabla_{\underline{b}} \right] &= \delta t_{\underline{a}\underline{b}}^{\underline{c}} \nabla_{\underline{c}} + t_{\underline{a}\underline{b}}^{\underline{c}} \delta \nabla_{\underline{c}} + \delta \frac{1}{2} r_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} , \end{aligned} \quad (270)$$

we can rapidly derive the results

$$\begin{aligned} \delta t_{\underline{a}\underline{b}}^{\underline{d}} &= \nabla_{\underline{a}} h_{\underline{b}}^{\underline{c}} - \nabla_{\underline{b}} h_{\underline{a}}^{\underline{c}} - h_{\underline{a}}^{\underline{c}} t_{\underline{b}\underline{c}}^{\underline{d}} + h_{\underline{b}}^{\underline{c}} t_{\underline{a}\underline{c}}^{\underline{d}} + t_{\underline{a}\underline{b}}^{\underline{c}} h_{\underline{c}}^{\underline{d}} \\ &\quad - h_{\underline{a}\underline{b}}^{\underline{d}} + h_{\underline{b}\underline{a}}^{\underline{d}} , \\ \delta r_{\underline{a}\underline{b}\underline{c}\underline{d}} &= \nabla_{\underline{a}} h_{\underline{b}\underline{c}\underline{d}} - \nabla_{\underline{b}} h_{\underline{a}\underline{c}\underline{d}} - t_{\underline{a}\underline{b}}^{\underline{e}} h_{\underline{e}\underline{c}\underline{d}} + h_{\underline{c}}^{\underline{e}} r_{\underline{a}\underline{b}\underline{e}\underline{d}} \\ &\quad - h_{\underline{a}}^{\underline{e}} r_{\underline{b}\underline{e}\underline{c}\underline{d}} + h_{\underline{b}}^{\underline{e}} r_{\underline{a}\underline{e}\underline{c}\underline{d}} + h_{\underline{d}}^{\underline{e}} r_{\underline{a}\underline{b}\underline{c}\underline{e}} . \end{aligned} \quad (271)$$

We will make use of some of these later.

Let us now return to the issue of the spin-connection. As an assumption, we can impose the condition that the  $t_{\underline{a}\underline{b}}^{\underline{c}}$  should vanish. The solution to this condition is given by choosing

$$\omega_{\underline{a}\underline{b}\underline{c}} = \omega_{\underline{a}\underline{b}\underline{c}}(e) \equiv \frac{1}{2} [ c_{\underline{a}\underline{b}\underline{c}} - c_{\underline{a}\underline{c}\underline{b}} - c_{\underline{b}\underline{c}\underline{a}} ] . \quad (272)$$

So the gauge theory of gravity shares with supersymmetric Maxwell theory the feature that not all of the gauge fields introduced in the covariant derivative need be independent. With the choice  $t_{\underline{a}\underline{b}}^{\underline{c}} = 0$  we find

$$\left[ \nabla_{\underline{a}}, \nabla_{\underline{b}} \right] = \frac{1}{2} r_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} , \quad (273)$$

and the resulting theory is “torsionless.” (In differential geometry this is a “Riemann geometry” as compared to a “Riemann-Cartan geometry” where  $t_{\underline{a}\underline{b}}^{\underline{c}} \neq 0$ .) Since in this expression  $\omega_{\underline{a}\underline{b}\underline{c}}$  is first order in derivatives of the remaining gauge field  $e_{\underline{a}}^{\underline{m}}$ , the quantity  $r_{\underline{a}\underline{b}\underline{c}\underline{d}}$  is second order in derivatives.

One of the curious differences between  $r_{\underline{a}\underline{b}\underline{c}\underline{d}}$  and  $f_{\underline{a}\underline{b}}^{\underline{I}}$  in a Yang-Mills theory is that only in the former case is it true that the “group indices” (i.e. the  $\underline{c}\underline{d}$  indices that were originally contracted with  $\mathcal{M}_{\underline{c}\underline{d}}$  generator) are of the same type as the 4D space-time flat vector indices. This allows the following contraction

$$r_{\underline{a}\underline{c}} = r_{\underline{a}\underline{d}\underline{c}}^{\underline{d}} , \quad (274)$$



which has no analog in Yang-Mills theory. (This contracted quantity is called the Ricci tensor.) Given this quantity it is possible to form one more such contraction

$$r = \eta^{a\bar{c}} r_{\underline{a}\underline{d}\underline{c}}^{\underline{d}} . \quad (275)$$

This object with no remaining free indices is the curvature scalar.

In Yang-Mills theory, there is a result that follows from the Jacobi identity on the Yang-Mills covariant derivative  $\nabla_{\underline{a}} = \partial_{\underline{a}} + iA_{\underline{a}}^{\mathcal{I}} t_{\mathcal{I}}$ ,

$$\begin{aligned} 0 &= [[\nabla_{\underline{a}}, \nabla_{\underline{b}}], \nabla_{\underline{c}}] + [[\nabla_{\underline{b}}, \nabla_{\underline{c}}], \nabla_{\underline{a}}] + [[\nabla_{\underline{c}}, \nabla_{\underline{a}}], \nabla_{\underline{b}}] \\ &\rightarrow 0 = \nabla_{\underline{a}} F_{\underline{b}\underline{c}}^{\mathcal{I}} + \nabla_{\underline{b}} F_{\underline{c}\underline{a}}^{\mathcal{I}} + \nabla_{\underline{c}} F_{\underline{a}\underline{b}}^{\mathcal{I}} . \end{aligned} \quad (276)$$

This equation is also called the Bianchi identity. For the gravitationally covariant derivative  $\nabla_{\underline{a}} = e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}} + \frac{1}{2} \omega_{\underline{a}\underline{d}}^{\underline{e}} \mathcal{M}_{\underline{e}}^{\underline{d}}$  a similar result is obtained,

$$\begin{aligned} 0 &= [[\nabla_{\underline{a}}, \nabla_{\underline{b}}], \nabla_{\underline{c}}] + [[\nabla_{\underline{b}}, \nabla_{\underline{c}}], \nabla_{\underline{a}}] + [[\nabla_{\underline{c}}, \nabla_{\underline{a}}], \nabla_{\underline{b}}] \\ &\rightarrow 0 = \nabla_{\underline{a}} t_{\underline{b}\underline{c}}^{\underline{d}} + \nabla_{\underline{b}} t_{\underline{c}\underline{a}}^{\underline{d}} + \nabla_{\underline{c}} t_{\underline{a}\underline{b}}^{\underline{d}} \\ &\quad - t_{\underline{a}\underline{b}}^{\underline{e}} t_{\underline{e}\underline{c}}^{\underline{d}} - t_{\underline{b}\underline{c}}^{\underline{e}} t_{\underline{e}\underline{a}}^{\underline{d}} - t_{\underline{c}\underline{a}}^{\underline{e}} t_{\underline{e}\underline{b}}^{\underline{d}} \\ &\quad + r_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} + r_{\underline{b}\underline{c}\underline{a}}^{\underline{d}} + r_{\underline{c}\underline{a}\underline{b}}^{\underline{d}} , \\ 0 &= \nabla_{\underline{a}} r_{\underline{b}\underline{c}\underline{k}\underline{\ell}} + \nabla_{\underline{b}} r_{\underline{c}\underline{a}\underline{k}\underline{\ell}} + \nabla_{\underline{c}} r_{\underline{a}\underline{b}\underline{k}\underline{\ell}} \\ &\quad - t_{\underline{a}\underline{b}}^{\underline{e}} r_{\underline{e}\underline{c}\underline{k}\underline{\ell}} - t_{\underline{b}\underline{c}}^{\underline{e}} r_{\underline{e}\underline{a}\underline{k}\underline{\ell}} - t_{\underline{c}\underline{a}}^{\underline{e}} r_{\underline{e}\underline{b}\underline{k}\underline{\ell}} . \end{aligned} \quad (277)$$

Under the restriction that  $t_{\underline{a}\underline{b}}^{\underline{c}} = 0$ , the first of these becomes a purely algebraic restriction on  $r_{\underline{a}\underline{b}\underline{c}\underline{d}}$ . Under this same restriction the second set then takes a form that is identical to that of a Yang-Mills theory. So the similarities between the two types of covariant derivatives are very obvious.

### 3.3 Field Equations and Actions

However, we still have a few items to complete. We may start with the issue of the field equations. In Yang-Mills theory these are known to be

$$\eta^{a\bar{b}} \nabla_{\underline{a}} F_{\underline{b}\underline{c}}^{\mathcal{I}} = 0 . \quad (278)$$

Like many equations of physical significance this is a second order differential equation for the gauge field  $A_{\underline{a}}^{\underline{T}}$ . We have already seen that the condition that the field strength  $t_{\underline{a}\underline{b}}^{\underline{c}} = 0$  implies that the remaining field strength  $r_{\underline{a}\underline{b}\underline{d}\underline{e}}$  is second order in derivatives of the gauge field  $e_{\underline{a}}^{\underline{m}}$ . So if this gauge field also obeys a second order differential equation, this corresponds to something that must be linear in  $r_{\underline{a}\underline{b}\underline{d}\underline{e}}$ . The Yang-Mills equations are derivable from an action principle. It would be nice if this were also true for the field equation for  $e_{\underline{a}}^{\underline{m}}$ . In order to find this, we must go back and solve a problem which we left dangling, “How is  $K$ -gauge invariance to be realized at the level of actions?”

The infinitesimal version of (258) implies

$$\delta_K \mathcal{L}(\phi, \nabla_{\underline{a}} \phi) = \xi^{\underline{m}}(x) \partial_{\underline{m}} \mathcal{L}(\phi, \nabla_{\underline{a}} \phi) \quad , \quad (279)$$

and this is clearly not invariant. To fix this problem we go back and make an observation about the gauge field  $e_{\underline{a}}^{\underline{m}}$ . Since it is a  $4 \times 4$  matrix, its determinant may be calculated

$$e \equiv \det(e_{\underline{a}}^{\underline{m}}) \quad . \quad (280)$$

We also know the gauge variation of  $e_{\underline{a}}^{\underline{m}}$  given in (261) and this can be used to calculate the quantity  $h_{\underline{a}}^{\underline{b}}$  in (269). This takes the form

$$h_{\underline{a}}^{\underline{b}} = \xi^{\underline{n}}(\partial_{\underline{n}} e_{\underline{a}}^{\underline{m}}) e_{\underline{m}}^{\underline{b}} - e_{\underline{a}}^{\underline{n}}(\partial_{\underline{n}} \xi^{\underline{m}}) e_{\underline{m}}^{\underline{b}} - \lambda_{\underline{a}}^{\underline{b}} \quad . \quad (281)$$

Next we use equation (269) applied to directly on  $e$  along with the identity  $\det(M) = \exp[\text{Tr}(\ln M)]$  which is valid for any matrix to derive,

$$\delta_K e = e h_{\underline{a}}^{\underline{a}} = e \xi^{\underline{m}} \partial_{\underline{m}} e - e (\partial_{\underline{m}} \xi^{\underline{m}}) \quad . \quad (282)$$

This then implies

$$\delta_K e^{-1} = (\partial_{\underline{m}} \xi^{\underline{m}} e^{-1}) \quad . \quad (283)$$

This is a most fortunate result! Additionally we then see

$$\begin{aligned} \delta_K (e^{-1} \mathcal{L}) &= (\delta_K e^{-1}) \mathcal{L} + e^{-1} (\delta_K \mathcal{L}) \\ &= [(\partial_{\underline{m}} \xi^{\underline{m}} e^{-1})] \mathcal{L} + e^{-1} \xi^{\underline{m}} \partial_{\underline{m}} \mathcal{L} \\ &= \left[ \partial_{\underline{m}} (\xi^{\underline{m}} e^{-1} \mathcal{L}) \right] \quad . \end{aligned} \quad (284)$$

So if we integrate the quantity  $e^{-1} \mathcal{L}$ , we find

$$\delta_K \left[ \int d^4 x \, e^{-1} \mathcal{L} \right] = \int d^4 x \left[ \delta_K (e^{-1} \mathcal{L}) \right] = \int d^4 x \left[ \partial_{\underline{m}} (\xi^{\underline{m}} e^{-1} \mathcal{L}) \right] \quad . \quad (285)$$

In the middle result, we have used the fact that the operator  $\delta_K$  is defined to act on fields and as such it may be freely “pushed” past the integration measure  $d^4x$ . We thus reach the following understanding of gauge invariance. Although no Lagrangian is invariant under the action of  $\delta_K$ , the integrated product  $e^{-1}\mathcal{L}$  changes only by surface terms under the action of the Lie-algebra based view of the gravitational gauge group. As long as such surface terms are negligible, there is an effective gauge invariance of the theory. (Note also that this is curiously similar to the case of super translation invariance (82,85).)

We may now complete the effort to write the Einstein field equations derivable from an action. The required action (the Einstein-Hilbert action) is simply as

$$\mathcal{S}_{EH} = -\frac{3}{\kappa^2} \int d^4x e^{-1} r(\omega(e), e) \quad , \quad (286)$$

written in the conventions of *Superspace*. We have written it in this form to emphasize that the spin-connection  $\omega$  is the one associated with the vanishing torsion. The variation of this action (using the formula in (271) yields the Einstein field equation

$$\frac{\delta\mathcal{S}_{EH}}{h^{\underline{a}\underline{b}}} = 0 \quad \rightarrow \quad \mathcal{E}_{\underline{a}\underline{b}} \equiv r_{\underline{a}\underline{b}} - \frac{1}{2} \eta_{\underline{a}\underline{b}} r = 0 \quad , \quad (287)$$

as the analog of the Yang-Mills free field equations. (The tensor  $\mathcal{E}_{\underline{a}\underline{b}}$  is often called the “Einstein tensor.”) If we start with a more complicated action constructed from the sum of the Einstein-Hilbert action with other actions  $\mathcal{S}_M$ , then the field equations for  $e_{\underline{a}}^{\underline{m}}$  take the forms

$$r_{\underline{a}\underline{b}} - \frac{1}{2} \eta_{\underline{a}\underline{b}} r = \frac{\kappa^2}{3} \left( \frac{\delta\mathcal{S}_M}{h^{\underline{a}\underline{b}}} \right) \quad . \quad (288)$$

The quantity appearing on the rhs of this equation is the energy-momentum tensor due to the fields in  $\mathcal{S}_M$ . It is also possible to write one additional term that depends solely on the gauge field  $e_{\underline{a}}^{\underline{m}}$  and modifies its equation of motion by terms no more than quadratic in derivatives. This is the famous “cosmological term”

$$\mathcal{S}_{cos} = -\frac{3}{\kappa^2} \int d^4x e^{-1} \quad . \quad (289)$$

### 3.4 Principles for Matter Coupling

The minimal coupling procedure works for matter fields such as scalars or spinors. For example starting with our massless spin-0 or Weyl spinor actions

we are led to

$$\mathcal{S}_{scalar} = \int d^4x \, e^{-1} \left[ -\frac{1}{2}(\nabla^{\underline{a}}\overline{A})(\nabla_{\underline{a}}A) \right] \quad , \quad (290)$$

$$\mathcal{S}_{spinor} = \int d^4x \, e^{-1} \left[ -i\overline{\psi}^{\dot{\alpha}}\nabla_{\underline{a}}\psi^{\alpha} \right] \quad . \quad (291)$$

Note that we can even do some “bizarre” things once we have the gravitationally covariant derivative  $\nabla_{\underline{a}}$  and know how to use  $e^{-1}$  to construct invariant actions. For example, a Schrödinger field subject to a force with potential  $W(x)$ , has the action obtained by the minimal coupling prescription given by

$$\mathcal{S} = \int d^4x \, e^{-1} \left[ i(\overline{\Psi}\nabla_0\Psi - (\nabla_0\overline{\Psi})\Psi) - \frac{1}{2m}(\vec{\nabla}\overline{\Psi})(\vec{\nabla}\Psi) - W\overline{\Psi}\Psi \right] \quad . \quad (292)$$

This action possesses local  $P$ -invariance! However, the local  $SO(1,3)$ -invariance is broken down to local  $SO(3)$ -invariance by the fact that the temporal component of the the gravitational covariant derivative  $(\nabla_0)$  is treated in a way that is not symmetrical with regards to the spatial components  $(\vec{\nabla})$ . Note that the spin-connection is absent entirely from this action if we choose  $\mathcal{M}_{\alpha}^{\beta}\Psi = \overline{\mathcal{M}}_{\alpha}^{\beta}\Psi = 0$ .

The minimal coupling procedure, however, does not apply to matter fields that possess their own gauge invariances not related to the symmetries generated by the  $K$ -operator. This is most clearly seen by considering the case of Yang-Mills symmetries simultaneously present with the  $K$ -symmetries. This case is simplest when treated by an appropriate modification of the gravitationally covariant derivative to include Yang-Mills fields

$$\nabla_{\underline{a}} = e_{\underline{a}}^{\underline{m}}\partial_{\underline{m}} + \frac{1}{2}\omega_{\underline{a}\underline{d}}^{\underline{c}}\mathcal{M}_{\underline{c}}^{\underline{d}} - iA_{\underline{a}}^{\mathcal{I}}t_{\mathcal{I}} \quad . \quad (293)$$

The commutator of this modified derivative defines the Yang-Mills field strength via

$$[\nabla_{\underline{a}}, \nabla_{\underline{b}}] = t_{\underline{a}\underline{b}}^{\underline{c}}\nabla_{\underline{c}} + \frac{1}{2}r_{\underline{a}\underline{b}\underline{c}}^{\underline{d}}\mathcal{M}_{\underline{d}}^{\underline{c}} - if_{\underline{a}\underline{b}}^{\mathcal{I}}t_{\mathcal{I}} \quad , \quad (294)$$

and leads to a modified definition of  $f_{\underline{a}\underline{b}}^{\mathcal{A}}$

$$f_{\underline{a}\underline{b}}^{\mathcal{I}} \equiv e_{\underline{a}}^{\underline{m}}\partial_{\underline{m}}A_{\underline{b}}^{\mathcal{I}} - e_{\underline{b}}^{\underline{m}}\partial_{\underline{m}}A_{\underline{a}}^{\mathcal{I}} - c_{\underline{a}\underline{b}}^{\underline{c}}A_{\underline{c}}^{\mathcal{I}} - if_{\mathcal{A}\mathcal{B}}^{\mathcal{I}}A_{\underline{a}}^{\mathcal{A}}A_{\underline{b}}^{\mathcal{B}} \quad . \quad (295)$$

This result differs from that obtained by naive minimal coupling ( $\partial_{\underline{a}} \rightarrow e_{\underline{a}}^{\underline{m}}\partial_{\underline{m}}$ ) by the term proportional to the anholonomy  $c_{\underline{a}\underline{b}}^{\underline{c}}$ . However, once this definition is used the action for the Yang-Mills field takes its familiar form,

$$\mathcal{S}_{YM} = \int d^4x \, e^{-1} \left[ -\frac{1}{8}f_{\underline{a}\underline{b}}^{\mathcal{I}}f^{\underline{a}\underline{b}\mathcal{I}} \right] \quad . \quad (296)$$

### 3.5 Implicit Lorentz-Invariant and Higher Derivative Gravitation

From the view on the Lie algebraic origins of theories of gravitation, there are some issues that logically can be seen in their proper prospective. One such point that can be seen debated (almost continuously) in the literature is the issue of torsion and whether it should be considered as fundamental. The only role of the torsion is that it determines the form of the spin-connection  $\omega_{\underline{a}\underline{b}\underline{c}}$ . If we use the analogy with supersymmetric field theories, the spin-connection is simply an auxiliary field. To see how irrelevant is this auxiliary field, consider the action given by

$$\mathcal{S}_{C^2} = \int d^4x \, e^{-1} \left[ c_0 c^{\underline{a}\underline{b}\underline{c}} c_{\underline{a}\underline{b}\underline{c}} + c_1 c^{\underline{a}\underline{b}\underline{c}} c_{\underline{c}\underline{a}\underline{b}} + c_2 (c^{\underline{a}\underline{b}\underline{c}})^2 \right] , \quad (297)$$

where  $c_0$ ,  $c_1$  and  $c_2$  are constants. This action is also clearly independent of the spin-connection and is local a  $P$ -invariant. However, like our bizarre example of the Schrödinger field minimally coupled to gravity, it seems as though this action is also not  $\mathcal{M}$ -gauge invariant. In order to verify this, we may perform a local Lorentz gauge transformation upon  $e_{\underline{a}}^{\underline{m}}$  to find

$$\begin{aligned} \delta_{LL} e_{\underline{a}}^{\underline{m}} &= -\lambda_{\underline{a}}^{\underline{b}} e_{\underline{a}}^{\underline{m}} \rightarrow \\ \delta_{LL} \mathcal{S}_{C^2} &= \int d^4x \, e^{-1} \left[ (1 + 4c_0) c^{\underline{b}\underline{a}\underline{d}} (e_{\underline{a}} \lambda_{\underline{b}\underline{d}}) \right. \\ &\quad + (1 - 2c_1) c^{\underline{d}[\underline{a}\underline{b}]} (e_{\underline{a}} \lambda_{\underline{b}\underline{d}}) \\ &\quad + 2(c_2 - 1) c^{\underline{a}\underline{b}\underline{c}} (e_{\underline{d}} \lambda_{\underline{b}\underline{d}}) \\ &\quad \left. - 2((e^{\underline{a}} - c^{\underline{a}\underline{b}\underline{c}}) e^{\underline{c}} \lambda_{\underline{a}\underline{c}}) \right] . \end{aligned} \quad (298)$$

However, we have an identity of the form

$$e^{-1} ((e^{\underline{a}} - c^{\underline{a}\underline{b}\underline{c}}) X_{\underline{a}}) = \partial_{\underline{m}} (e^{-1} X_{\underline{a}} e^{\underline{a}\underline{m}}) , \quad (299)$$

substituting  $X_{\underline{a}} = e^{\underline{c}} \lambda_{\underline{a}\underline{c}}$  and it can be seen that for the special choice of coefficients given by  $-4c_0 = 2c_1 = c_2 = 1$ , the local Lorentz variation of the action is a total divergence. In our Lie-algebraic motivated approach to understanding gravity we always neglect total divergence terms, so effectively for this choice  $\mathcal{S}_{C^2}$  is both a  $P$ -gauge invariant and a  $\mathcal{M}$ -gauge invariant!

A consequence of the result (299) is that the rule for integration-by-parts takes the form

$$\int d^4x \, e^{-1} U^{\underline{a}[X]} e_{\underline{a}} V_{[X]} = - \int d^4x \, e^{-1} [(e_{\underline{a}} U^{\underline{a}[X]}) V_{[X]} - c_{\underline{a}\underline{b}}^{\underline{b}} U^{\underline{a}[X]} V_{[X]}] . \quad (300)$$

In this expression, we have used the notation  $[X]$  to denote any choice of Lorentz vector or spinor indices.

The equation of motion that follows from  $\mathcal{S}_{C^2}$  is clearly of the form of second order derivatives operating on  $e_{\underline{a}}^{\underline{m}}$  and is, in fact, the usual Einstein field equation for free space although it appears in the form

$$\begin{aligned} & e_{(\underline{k}} c_{\underline{l})}^{\underline{c}} - \frac{1}{2} c_{(k\underline{a}}^{\underline{cd}} c_{|\underline{l} \underline{c} \underline{d}} - c_{(\underline{k} | \underline{c}} c_{|\underline{l} \underline{k})}^{\underline{d}} + \frac{1}{4} c_{\underline{c} \underline{d}} c_{(\underline{k}} c^{\underline{c} \underline{d}}_{|\underline{l})} \\ & - 2e^{-1} [(e_{\underline{a}} - c_{\underline{a} \underline{b}}^{\underline{b}})(\eta_{\underline{k} \underline{l}} c_{\underline{d}}^{\underline{a}} - \frac{1}{2} c_{(\underline{k} \underline{l})}^{\underline{a}})] \\ & - \eta_{\underline{k} \underline{l}} \left[ -\frac{1}{4} c^{\underline{a} \underline{b} \underline{c}} c_{\underline{a} \underline{b} \underline{c}} + \frac{1}{2} c^{\underline{a} \underline{b} \underline{c}} c_{\underline{c} \underline{a} \underline{b}} + (c_{\underline{b}}^{\underline{a} \underline{b}})^2 \right] = 0 \quad . \end{aligned} \quad (301)$$

The issue of how (or if)  $\omega_{\underline{a} \underline{b} \underline{c}}$  is specified becomes particularly important when one considers theories (such as the effective gravitational interaction that emerges at the low-energy limit of various string theories<sup>9</sup>) which possess what are commonly referred to as higher curvature terms. A generic term of this form is given by

$$\mathcal{S}_{H.D.} = \int d^4x e^{-1} \left[ f(\nabla_{\underline{a}}, r_{\underline{a} \underline{b} \underline{c} \underline{d}}) \right] \quad , \quad r_{\underline{a} \underline{b} \underline{c} \underline{d}} = r_{\underline{a} \underline{b} \underline{c} \underline{d}}(\omega(e), e) \quad , \quad (302)$$

where  $f$  is a function that is quadratic or higher in the curvature tensor (and its possible contractions) and may depend on of the gravitationally covariant derivative. Since  $\omega_{\underline{a} \underline{b} \underline{c}}(e)$  is solely a function of  $e_{\underline{a}}^{\underline{m}}$ , the only dynamics here are associated with the graviton. In fact, the action above must satisfy (up to a total derivative)

$$\int d^4x e^{-1} \left[ f(\nabla_{\underline{a}}, r_{\underline{a} \underline{b} \underline{c} \underline{d}}) \right] = \int d^4x e^{-1} \left[ g(e_{\underline{a}}, c_{\underline{b} \underline{c} \underline{d}}) \right] \quad , \quad (303)$$

for some function  $g$ . The simplest example of this class of identities is provided by

$$\begin{aligned} f &= \eta^{\underline{a} \underline{c}} \eta^{\underline{b} \underline{d}} r_{\underline{a} \underline{b} \underline{c} \underline{d}}(\omega(e), e) \quad , \\ g &= \left[ -\frac{1}{4} c^{\underline{a} \underline{b} \underline{c}} c_{\underline{a} \underline{b} \underline{c}} + \frac{1}{2} c^{\underline{a} \underline{b} \underline{c}} c_{\underline{c} \underline{a} \underline{b}} + (c_{\underline{b}}^{\underline{a} \underline{b}})^2 \right] \quad . \end{aligned} \quad (304)$$

On the other hand, if we start with the action

$$\mathcal{S}'_{H.D.} = \int d^4x e^{-1} \left[ f(\widehat{\nabla}_{\underline{a}}, \widehat{r}_{\underline{a} \underline{b} \underline{c} \underline{d}}) \right] \quad , \quad \widehat{r}_{\underline{a} \underline{b} \underline{c} \underline{d}} = r_{\underline{a} \underline{b} \underline{c} \underline{d}}(\widehat{\omega}, e) \quad , \quad (305)$$

---

<sup>9</sup>To my knowledge, no rigorous derivations of quantum nonlinear gravitational effective actions have ever been found for any theory other than strings.

where  $\widehat{\omega}_{\underline{a}\underline{b}\underline{c}}$  is left as an unspecified function, then in addition to the dynamics of  $e_{\underline{a}}^{\underline{m}}$ , one must also encounter dynamical equations for  $\widehat{\omega}_{\underline{a}\underline{b}\underline{c}}$ ! Thus, the physics described by  $\mathcal{S}_{H.D.}$  is radically different from that described by  $\mathcal{S}'_{H.D.}$  even though exactly the same function  $f$  appears in the two actions.

As mentioned above, the low-energy purely gravitational field approximation of string-type theories (i.e. where all fields other than  $e_{\underline{a}}^{\underline{m}}$  are set to zero) takes the form<sup>*h*</sup>

$$\mathcal{S}_{stringy-grav.} \propto \mathcal{S}_{C^2} + \mathcal{S}_{H.D.} \quad . \quad (306)$$

It is of interest to note that there exists a restricted form of the higher derivative action that corresponds to a “velocity expansion.” Note that the quantity  $c_{\underline{a}\underline{b}\underline{c}}$  may be thought of as a velocity. One may thus contemplate the special class of higher derivative terms of the form

$$\mathcal{S}_{H.D.}(vel) = \int d^4x \, e^{-1} \sum_{\ell=3}^{\infty} \kappa^{\ell-4} L_{\ell} \mathcal{P}_{\underline{c}_1 \dots \underline{c}_{\ell}}^{(\ell) \underline{a}_1 \dots \underline{a}_{\ell} \underline{b}_1 \dots \underline{b}_{\ell}} \left( \prod_{j=1}^{\ell} c_{\underline{a}_j \underline{b}_j}^{\underline{c}_j} \right) \quad . \quad (307)$$

This expansion is roughly equivalent to the well-known chiral perturbation theory (“soft pion”) representation of meson physics. In this expression,  $L_{\ell}$  are a set of dimensionless numbers and  $\mathcal{P}_{\underline{c}_1 \dots \underline{c}_{\ell}}^{(\ell) \underline{a}_1 \dots \underline{a}_{\ell} \underline{b}_1 \dots \underline{b}_{\ell}}$  correspond to a set of constant tensors chosen in complete generality consistent with the  $K$ -gauge invariance of this action.

We close this section by noting that the quantity  $c_{\underline{a}\underline{b}}^{\underline{c}}$  is the closest gravitational analog to the field strength  $f_{\underline{a}\underline{b}}^{\underline{I}}$  of Yang-Mills theories. Accordingly, we might expect that many of the properties of Yang-Mills theories should have ‘echoes’ for gravitational theories that may be recognized by using this analogy.

---

<sup>*h*</sup>The  $\beta$ -function technique within the confines of bosonic and NSR  $\sigma$ -models is known to provide an explicit way to derive this expansion. The fact that  $\mathcal{S}_{H.D.}$  and not  $\mathcal{S}'_{H.D.}$  appears in string theories is obvious since a spacetime spin connection is ‘removable’ from the  $\sigma$ -model formalism.

## 4 Fourth Lecture: Junior 4D, $N = 1$ Superfield Theory

### *Introduction*

In today's lecture, we will discuss the issue of quantization via functional methods. The discussion will include the preliminary development of superfunctional techniques, definitions of super propagators and supergraphs through the examples of the chiral scalar and "gauge" multiplets. Let me begin by saying something about what I am not going to do.

In going from being an official student to a practitioner (e.g. unofficial student) of QFT, I learned or have learned a number of quantization techniques:

- A. Canonical,
- B. Path Integral,
  - (i) BRST,
  - (ii) Batalin - type,
    - (a.) BV,
    - (b.) BFV.

Although the canonical method is considered the most "rigorous," I know of no completely successful examples of its implementation within the realm of supersymmetrical theories. I suspect that the reason for this is the fact that 4D,  $N = 1$  superspace makes fundamental use of relativistic spinors (i.e. the Grassmann coordinates belong to this representation). The 4-momentum can be written as  $P_{\underline{m}} = (E, \vec{p})$  and in the canonical approach  $E$  or the Hamiltonian is treated differently from the other components. Thus, there is a need to break 4D Lorentz invariance and hence break manifest supersymmetry. Instead, the discussion presented will very closely follow that of *Superspace*.

Accordingly, we will not discuss canonical approaches. In this lecture, we will exclusively consider the path integral approach using at most BRST type techniques. This material can be found in our book *Superspace*. So I won't have anything to say about superfield Batalin-type (BV, BFV) quantization with its attendant use of "the Master Equation" either. I do point out, however, that M. Grisaru et. al. have recently<sup>20</sup> discussed such matters in connection with the quantization of the complex linear multiplet (which was introduced as the dual to the chiral scalar multiplet two lecture ago). The results are most interesting and complicated. The class of Batalin-type quantization techniques are the most advanced known to date. However, in the attempt to quantize the complex linear multiplet, the results derived do not seem easily applicable for the derivation of additional results in interacting theories. This raises the question of the need for as yet unknown advances in the Batalin-type methods.



## Lecture

### 4.1 A Little about Superfunctional Differentiation

We can begin by noting that in ordinary four dimensional quantum field theory treated via functional methods, the Dirac delta function plays a fundamental role in the definition of functional differentiation. Given some function  $J(x)$  (dependent on the 4-vector  $x^{\underline{m}}$ ) we define

$$\frac{\delta J(x)}{\delta J(x')} \equiv \delta^{(4)}(x - x') \quad , \quad (308)$$

and since the operator  $\frac{\delta}{\delta J(x)}$  is essential, we need to have a definition such that given a superfunction  $J(z)$  we may define

$$\frac{\delta J(z)}{\delta J(z')} \equiv \delta^{(4)}(x - x') \delta^{(4)}(\theta - \theta') \quad . \quad (309)$$

But what is the meaning of  $\delta^{(4)}(\theta - \theta')$ ? For a single bosonic coordinate  $y$ , we know that

$$\int_{-\infty}^{\infty} dy f(y) \delta(y - y') = f(y') \quad . \quad (310)$$

So we need something similar in a space with a single bosonic coordinate  $y$  and a single fermionic one  $\zeta$ . A real superfunction  $f(y, \zeta)$  has the form

$$f(y, \zeta) = a(y) + i\zeta\beta(y) \quad . \quad (311)$$

If there is a super Dirac delta function  $\delta(\zeta - \zeta')$  then we expect

$$\int d\zeta f(y, \zeta) \delta(\zeta - \zeta') = f(y, \zeta') \quad . \quad (312)$$

To find an explicit representation for  $\delta(\zeta - \zeta')$  we recall

$$\begin{aligned} \int d\zeta 1 &\equiv \left[ \frac{\partial}{\partial \zeta} + i\frac{1}{2}\zeta \frac{\partial}{\partial y} \right] 1 \Big| = 0 \quad , \\ \int d\zeta \zeta &\equiv \left[ \frac{\partial}{\partial \zeta} + i\frac{1}{2}\zeta \frac{\partial}{\partial y} \right] \zeta \Big| = 1 \quad . \end{aligned} \quad (313)$$

We also note  $f(y, \zeta) (\zeta - \zeta') = \zeta f(y, \zeta') - a(y)\zeta'$ . Thus we have

$$\int d\zeta f(y, \zeta) (\zeta - \zeta') = f(y, \zeta') \quad . \quad (314)$$

so that is it consistent to define  $\delta(\zeta - \zeta') = (\zeta - \zeta')$ . In a superspace with more fermionic dimensions, we can proceed inductively to derive in our case

$$\begin{aligned}\delta^{(4)}(\theta - \theta') &\equiv \delta^{(2)}(\theta - \theta') \delta^{(2)}(\bar{\theta} - \bar{\theta}') \quad , \\ \delta^{(2)}(\theta - \theta') &\equiv \frac{1}{2} (\theta - \theta')^\alpha (\theta - \theta')_\alpha \quad , \\ \delta^{(2)}(\bar{\theta} - \bar{\theta}') &\equiv \frac{1}{2} (\bar{\theta} - \bar{\theta}')^{\dot{\alpha}} (\bar{\theta} - \bar{\theta}')_{\dot{\alpha}} \quad .\end{aligned}\tag{315}$$

So given a general function  $J(z)$ , we define its functional derivative as in (308). Now the super operator  $\frac{\delta}{\delta J(x)}$  can be applied to superspace functionals. For example,

$$\begin{aligned}\mathcal{S} &= \int d^4x d^2\theta d^2\bar{\theta} A(z) J(z) \equiv \int d^8z A(z) J(z) \\ \rightarrow \frac{\delta \mathcal{S}}{\delta J(z')} &= \int d^8z A(z) \delta^{(8)}(z - z') = A(z') \quad , \\ \text{and if } \delta^{(8)}(z - z') &\equiv \delta^{(4)}(x - x') \delta^{(4)}(\theta - \theta') \quad , \\ \text{if } \frac{\delta J(z)}{\delta J(z')} &\equiv \delta^{(8)}(z - z') \quad .\end{aligned}\tag{316}$$

#### 4.2 On Superdeterminants

It sometimes occurs that we require the definition of a super determinant (i.e. to calculate the Jacobian of a change of field variables). For this purpose, it is first useful to define a supermatrix. For purposes of illustration, we return to the one dimensional superspace described by  $y$  and  $\zeta$ . We introduce a supervector and supermatrix

$$z = \begin{pmatrix} y \\ \zeta \end{pmatrix} \quad , \quad M_s = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad ,\tag{317}$$

such that the supermatrix can act on the supervector  $z$ . We see

$$z' = M_s z \quad \rightarrow \quad \begin{pmatrix} y' \\ \zeta' \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} y \\ \zeta \end{pmatrix} .\tag{318}$$

But  $y'$  will not be bosonic if  $B$  is an ordinary (non-fermionic) number. Similarly,  $\zeta'$  will not be fermionic if  $C$  is an ordinary number. Thus  $B$  and  $C$  must

be Grassmann numbers. How does one define a superdeterminant for  $M_s$ ? This problem was solved long ago by Arnowitt, Nath and Zumino<sup>21</sup>. We will give below a definition that leads to their results.

$$sdet(M_s) \equiv K_0 \int dy dy' d\zeta d\zeta' \exp[-\tilde{z}^t M_s z] \quad , \quad \tilde{z}^t \equiv (\tilde{z}, \tilde{\zeta}) \quad . \quad (319)$$

where the constant  $K_0$  is chosen so that  $sdet(I) = 1$  By shifting  $\tilde{y}$  and  $y$  according to  $\tilde{y} \rightarrow \tilde{y} - \tilde{\zeta} C A^{-1}$  and  $y \rightarrow y - A^{-1} B \zeta$  the exponential becomes

$$\exp[-\tilde{z}^t M_s z] = \exp[-\tilde{y} A y - \tilde{\zeta} (D - C A^{-1} B) \zeta] \quad . \quad (320)$$

Instead we can choose to shift the Grassmann coordinates according to  $\zeta \rightarrow$  and  $\tilde{\zeta} \rightarrow$  so that the exponential becomes

$$\exp[-\tilde{z}^t M_s z] = \exp[-\tilde{y} (A - B D^{-1} C) y - \tilde{\zeta} D \zeta] \quad . \quad (321)$$

After performing either of these shifts, the integrals can be done and yield

$$sdet(M_s) = \frac{A}{D - C A^{-1} B} = \frac{A - B D^{-1} C}{D} \quad . \quad (322)$$

If we replace  $A$ ,  $B$ ,  $C$ , and  $D$  by  $2 \times 2$  matrices then  $y$  and  $\zeta$  must become 2-component entities. The same is true for  $\tilde{y}$  and  $\tilde{\zeta}$ . The integrals then yield

$$sdet(M_s) = \frac{\det(A)}{\det[D - C A^{-1} B]} = \frac{\det[A - B D^{-1} C]}{\det(D)} \quad . \quad (323)$$

This is the form of the general answer for a supermatrix. This formula implies that either  $A$  or  $D$  should be nonsingular in order for the superdeterminant to have a well defined meaning.

#### 4.3 Review of the Functional Quantization of the Scalar Field

In order to orient ourselves, let's review ordinary 4D “ $\phi^4$ ” theory.

$$\mathcal{S} = \int d^4x \left[ \frac{1}{2} \phi (\square - m^2) \phi - \frac{\lambda}{4!} \phi^4 \right] \quad . \quad (324)$$

We also add the source term

$$\mathcal{S}_J = \int d^4x J \phi \quad . \quad (325)$$

If we ignore momentarily the interaction term in the Lagrangian

$$\mathcal{S}_{Int} = \int d^4x \left[ -\frac{\lambda}{4!} \phi^4 \right] , \quad (326)$$

and derive the equation of motion for  $\phi$ , we find

$$\begin{aligned} (\square - m^2)\phi &= -J , \\ \phi &= -\left(\frac{J}{\square - m^2}\right) . \end{aligned} \quad (327)$$

and this may be substituted into  $\mathcal{S} - \mathcal{S}_{Int} \equiv W_0(J)$  and we find

$$W_0(J) = \int d^4x \left[ -\frac{1}{2} J \left( \frac{1}{\square - m^2} \right) J \right] . \quad (328)$$

The generator of 1PI Green's function is thus

$$Z(J) = \exp\left\{ \int d^4x \left[ -\frac{\lambda}{4!} \left( \frac{\delta}{\delta J(x)} \right)^4 \right] \right\} \exp\{W_0(J)\} . \quad (329)$$

The propagator is just defined by

$$\begin{aligned} \Delta(x - x') &= \frac{\delta^2 Z(J)}{\delta J(x) \delta J(x')} \Big|_{J=0} \\ &= \frac{1}{\square - m^2} \delta^{(4)}(x - x') + \mathcal{O}(\lambda) . \end{aligned} \quad (330)$$

If we ignore all higher order  $\lambda$  terms this is the bare propagator. In a similar manner the 4-point function is just

$$G^{(4)}(x_1, x_2, x_3, x_4) = \frac{\delta^4 Z(J)}{\delta J(x_1) \delta J(x_2) \delta J(x_3) \delta J(x_4)} \Big|_{J=0} . \quad (331)$$

#### 4.4 Equation of Motion of the Chiral Scalar Superfield

The simplest superfield theory is the free massive chiral multiplet described by the superfield action

$$\mathcal{S}_c = \int d^8z \, \overline{\Phi} \Phi - \left[ \int d^6z \, \frac{1}{2} m \Phi^2 + \text{h.c.} \right] . \quad (332)$$

If we add chiral sources  $J_c$  and  $\overline{J}_c$

$$\mathcal{S}_J = \left[ \int d^6z \, J_c \Phi + \text{h.c.} \right] . \quad (333)$$

we would then like to treat this system just like the scalar “ $\phi^4$ -theory.” First we would like to derive the superfield equation of motion. However, the superfield  $\Phi$  satisfies the differential constraint  $\overline{D}_\alpha \Phi = 0$ . This means that we cannot freely vary the superfield  $\Phi$ . As a simpler example of the problem we have encountered, let us consider an ordinary 4D theory that illustrates the nature of this problem and as well points the way to the solution.

The action for 4D Maxwell theory is

$$\mathcal{S}_{Maxwell} = -\frac{1}{8} \int d^4x f_{\underline{a}\underline{b}} f_{\underline{a}\underline{b}} , \quad (334)$$

and the field strength  $f_{\underline{a}\underline{b}}$  satisfies the differential constraint

$$\partial_{\underline{a}} f_{\underline{b}\underline{c}} + \partial_{\underline{b}} f_{\underline{c}\underline{a}} + \partial_{\underline{c}} f_{\underline{a}\underline{b}} = 0 . \quad (335)$$

If we ignore the constraint on  $f_{\underline{a}\underline{b}}$  and vary the action anyway we find

$$\frac{\delta \mathcal{S}_{Maxwell}}{\delta f_{\underline{b}\underline{c}}} = 0 \rightarrow f_{\underline{b}\underline{c}} = 0 . \quad (336)$$

Of course, the correct way to proceed is to first “solve” the constraint

$$f_{\underline{b}\underline{c}} = \partial_{\underline{b}} A_{\underline{c}} - \partial_{\underline{c}} A_{\underline{b}} , \quad (337)$$

and recognize that a variation of  $f_{\underline{b}\underline{c}}$  takes the form  $\delta f_{\underline{b}\underline{c}} = \partial_{\underline{b}}(\delta A_{\underline{c}}) - \partial_{\underline{c}}(\delta A_{\underline{b}})$  and thus

$$\frac{\delta \mathcal{S}_{Maxwell}}{\delta A_{\underline{b}}} = 0 \rightarrow \partial^{\underline{a}} f_{\underline{a}\underline{b}} = 0 . \quad (338)$$

These observations offer us a notion of how to correctly derive the equation of motion for  $\Phi$ . We first observe

$$\begin{aligned} D_\alpha D_\beta &= -D_\beta D_\alpha \rightarrow D_\alpha D_\beta = -C_{\alpha\beta} D^2 , \\ \rightarrow D_\alpha D_\beta D_\gamma &= -C_{\beta\gamma} D_\alpha D^2 = -\frac{1}{2} C_{\beta\gamma} D_\alpha D^\delta D_\delta \\ &= \frac{1}{2} C_{\beta\gamma} D_\alpha D_\delta D^\delta = -\frac{1}{2} C_{\beta\gamma} D_\delta D_\alpha D^\delta \\ &= \frac{1}{2} C_{\beta\gamma} D_\delta D^\delta D_\alpha = \frac{1}{2} C_{\beta\gamma} D_\delta (\delta_\alpha^\delta D^2) \\ &= \frac{1}{2} C_{\beta\gamma} D_\alpha D^2 = -D_\alpha D_\beta D_\gamma . \\ \rightarrow D_\alpha D_\beta D_\gamma &= -D_\alpha D_\beta D_\gamma \rightarrow \\ D_\alpha D^2 &= 0 \rightarrow \overline{D}_\alpha \overline{D}^2 = 0 . \end{aligned} \quad (339)$$

$$\begin{aligned} \rightarrow D_\alpha D_\beta D_\gamma &= -D_\alpha D_\beta D_\gamma \rightarrow \\ D_\alpha D^2 &= 0 \rightarrow \overline{D}_\alpha \overline{D}^2 = 0 . \end{aligned} \quad (340)$$

So that we see the equation  $\overline{D}_\alpha \Phi = 0$  has a solution  $\Phi = \overline{D}^2 \overline{U}$  where  $U$  is a complex unconstrained (i.e. it is free of all differential constraints) pre-potential superfield. We use  $U$  exactly like  $A_{\underline{a}}$  in Maxwell theory.

$$\begin{aligned} \mathcal{S}_c = & \int d^8 z (\overline{D}^2 \overline{U}) (D^2 U) - \left[ \int d^6 z \frac{1}{2} m (\overline{D}^2 \overline{U})^2 + \text{h.c.} \right] \\ & + \left[ \int d^6 z J_c (\overline{D}^2 \overline{U}) + \text{h.c.} \right] . \end{aligned} \quad (341)$$

If we vary with respect to  $U$  we find

$$\begin{aligned} \delta \mathcal{S}_c = & \int d^8 z (D^2 \delta U) (\overline{D}^2 \overline{U}) - \int d^6 \bar{z} m (D^2 \delta U) (D^2 U) \\ & + \int d^6 \bar{z} \overline{J}_c (D^2 \delta U) . \end{aligned} \quad (342)$$

On the last two integrals, we can use the fact that  $\int d^8 z = \int d^6 \bar{z} D^2$ , thus

$$\delta \mathcal{S}_c = \int d^8 z \left[ (D^2 \delta U) (\overline{D}^2 \overline{U}) - m \delta U (D^2 U) + \overline{J}_c \delta U \right] , \quad (343)$$

$$\rightarrow \delta \mathcal{S}_c = \int d^8 z \left[ (D^2 \delta U) \Phi - m \delta U \overline{\Phi} + \overline{J}_c \delta U \right] . \quad (344)$$

To make further progress we have the identity

$$\begin{aligned} (D^\alpha D_\alpha A) B &= D^\alpha [(D_\alpha A) B] - [(D^\alpha A) (D_\alpha B)] \\ &= A (D^\alpha D_\alpha B) - D^\alpha [(D_\alpha A) B - A (D_\alpha B)] . \end{aligned} \quad (345)$$

Upon choosing  $A = \delta U$  and  $B = \Phi$  this identity then implies

$$\delta \mathcal{S}_c = \int d^8 z \delta U \left[ D^2 \Phi - m \overline{\Phi} + \overline{J}_c \right] . \quad (346)$$

Thus we derive the equation of motion from

$$\frac{\delta \mathcal{S}_c}{\delta U(z')} = 0 \rightarrow D^2 \Phi - m \overline{\Phi} = -\overline{J}_c , \quad (347)$$

and acting on this with  $\overline{D}^2$  on both sides then leads to

$$\begin{aligned} \overline{D}^2 D^2 \Phi - m \overline{D}^2 \overline{\Phi} &= -\overline{D}^2 \overline{J}_c \\ \rightarrow \overline{D}^2 D^2 \Phi - m (m \Phi - J_c) &= -\overline{D}^2 \overline{J}_c \\ \overline{D}^2 D^2 \Phi - m^2 \Phi &= -(m J_c + \overline{D}^2 \overline{J}_c) . \end{aligned} \quad (348)$$

Now we need to do a little D-algebra

$$\begin{aligned}
\overline{D}^2 D^2 &= \frac{1}{4} \overline{D}^\alpha \overline{D}_\alpha \cdot D^\alpha D_\alpha = -\frac{1}{4} \overline{D}^\alpha \overline{D}_\alpha \cdot D_\alpha D^\alpha \\
&= \frac{1}{4} \overline{D}^\alpha D_\alpha \overline{D}_\alpha \cdot D^\alpha - i \frac{1}{4} \partial_{\underline{a}} \overline{D}^\alpha D^\alpha \\
&= -\frac{1}{4} \overline{D}^\alpha D^\alpha \overline{D}_\alpha \cdot D_\alpha + i \frac{1}{4} \partial_{\underline{a}} D^\alpha \overline{D}^\alpha + \frac{1}{4} \partial^a \partial_{\underline{a}} \\
&= \frac{1}{4} \overline{D}^\alpha D^\alpha D_\alpha \overline{D}_\alpha \cdot - i \frac{1}{4} \partial_{\underline{a}} \overline{D}^\alpha D^\alpha + i \frac{1}{4} \partial_{\underline{a}} D^\alpha \overline{D}^\alpha + \frac{1}{4} \partial^a \partial_{\underline{a}} \\
&= \frac{1}{2} \overline{D}^\alpha D^2 \overline{D}_\alpha \cdot + i \frac{1}{2} \partial_{\underline{a}} D^\alpha \overline{D}^\alpha + \square \\
&\rightarrow \overline{D}^2 D^2 \Phi = \square \Phi .
\end{aligned} \tag{349}$$

So the equation of motion for  $\Phi$  in the presence of sources  $J$  and  $\overline{J}$  is

$$(\square - m^2)\Phi = -(mJ_c + \overline{D}^2 \overline{J}_c) . \tag{350}$$

#### 4.5 Functional Quantization of the Chiral Scalar Superfield

At this point for  $\phi^4$ -theory, we simply “inverted” the operator on the lhs by performing a formal division. The justification for this is that we could always choose to work in momentum space via the use of Fourier transformations with respect to the  $x$ -space coordinates. The most obvious generalization is to work in the Fourier transform with respect to  $z$ -space. However, this is not the most convenient way to proceed. Instead the Fourier transform need only be taken with respect to the  $x$ -subspace of superspace. After going to momentum space, the equation of motion is of the form,

$$(p^2 + m^2)\Phi = (mJ_c + \overline{D}^2 \overline{J}_c) , \tag{351}$$

where in this equation we regard  $\Phi = \Phi(\theta, \bar{\theta}, p)$  and  $J_c = J_c(\theta, \bar{\theta}, p)$ . As well the  $D$ -operator is written in  $p$ -space

$$D_\alpha = \partial_\alpha + \frac{1}{2} \bar{\theta}^\alpha p_{\underline{a}} . \tag{352}$$

With this understanding, the equation of motion for  $\Phi$  can be inverted to express  $\Phi$  as a function of  $J_c$  and  $\overline{J}_c$ ,

$$\Phi = - \left[ \frac{mJ_c + \overline{D}^2 \overline{J}_c}{\square - m^2} \right] \quad \text{or} \quad \overline{\Phi} = - \left[ \frac{m\overline{J}_c + D^2 J_c}{\square - m^2} \right] . \tag{353}$$

If we substitute the  $x$ -space version back into the original action we obtain

$$W_0(J_c, \bar{J}_c) = - \int d^8 z \left\{ \bar{J}_c \left[ \frac{1}{\square - m^2} \right] J_c + \frac{1}{2} J_c \left[ \frac{m D^2}{\square (\square - m^2)} \right] J_c \right. \\ \left. + \frac{1}{2} \bar{J}_c \left[ \frac{m \bar{D}^2}{\square (\square - m^2)} \right] \bar{J}_c \right\} . \quad (354)$$

We can define a superfield 1PI generating functional via

$$Z_0(J_c, \bar{J}_c) = \exp[W_0(J_c, \bar{J}_c)] . \quad (355)$$

This is valid, of course, only for the free massive theory. In the case of an interacting theory where we can write

$$\mathcal{L}_{Tot} = \mathcal{L}_0 + \mathcal{L}_{Int}(\Phi, \bar{\Phi}) , \quad (356)$$

the generating functional becomes

$$Z(J_c, \bar{J}_c) = \exp \left[ \int d^8 z \mathcal{L}_{Int} \left( \frac{\delta}{\delta J_c}, \frac{\delta}{\delta \bar{J}} \right) \right] Z_0(J_c, \bar{J}_c) . \quad (357)$$

In order for this definition to make sense, we require a definition of the functional derivative taken with respect to chiral (as opposed to general) superfields. For this purpose we note,

$$\frac{\delta J_c(z)}{\delta J_c(z')} = \bar{D}^2 \delta^{(8)}(z - z') . \quad (358)$$

In a theory only involving chiral superfields we can use this generating function to derive superfield Feynman rules.

(a.) Propagators

$$\begin{aligned} \bar{\Phi} \Phi : & \quad \frac{1}{p^2 + m^2} \delta^{(4)}(\theta - \theta') , \\ \Phi \Phi : & \quad - \frac{m D^2}{p^2(p^2 + m^2)} \delta^{(4)}(\theta - \theta') , \\ \bar{\Phi} \bar{\Phi} : & \quad - \frac{m \bar{D}^2}{p^2(p^2 + m^2)} \delta^{(4)}(\theta - \theta') . \end{aligned} \quad (359)$$

For simplicity's sake only, we can consider the massless limit where the latter two propagators vanish. We may now look at what must be going on in terms of the component fields in the massless limit. If we take  $\langle 0 | \bar{\Phi} \Phi | 0 \rangle$  this must correspond to the propagator  $\langle 0 | \bar{A}(x) A(x') | 0 \rangle$ . If we act on this propagator with  $D_\alpha \bar{D}_\alpha$  we obtain

$$(\bar{D}_\alpha \bar{\Phi})(D_\alpha \Phi) : \quad \frac{D_\alpha \bar{D}_\alpha}{p^2} \delta^{(4)}(\theta - \theta') , \quad (360)$$



and after applying the  $|\cdot\rangle$ -operation, this must correspond to the propagator  $\langle 0|\bar{\psi}_\alpha(x)\psi_\alpha(x')|0\rangle$ . Due to the interacting super Feynman rules, we can replace  $D_\alpha \bar{D}_\alpha \sim ip_{\underline{a}}$  and thus the propagator takes its usual form  $ip_{\underline{a}}/p^2$ . Finally, there is a propagator for the  $F$  field. We can find it by applying  $\bar{D}^2 D^2$  leading to

$$(\bar{D}^2 \bar{\Phi})(D^2 \Phi) : \quad \frac{\bar{D}^2 D^2}{p^2} \delta^{(4)}(\theta - \theta') \quad . \quad (361)$$

Now it might appear that the auxiliary field  $F$  is propagating a massless state (i.e. there seems to be a pole at  $p^2 = 0$ ). But once again the way the super Feynman rules work we can replace  $\bar{D}^2 D^2 \sim -p^2$  so there is no pole at  $p^2 = 0$ .

#### 4.6 Feynman Rules for the Chiral Scalar Superfield

To actually derive both the propagators as well as the rest of the super Feynman rules we require a definition of  $\delta\Phi(z)/\delta\Phi(z')$  where  $\bar{D}_\alpha \Phi = 0$ . For this purpose we simply note the result in (358). The following Feynman rules are obtained in a straight forward way.

- (a.) Vertices are read from the interaction Lagrangian in the usual way for ordinary four dimensional theory, with the additional feature that for each chiral or antichiral line leaving a vertex there is a factor of  $\bar{D}^2$  or  $D^2$  acting on the corresponding purely chiral or antichiral vertices we omit a  $\bar{D}^2$  or  $D^2$  factor from among the ones acting on the propagators.
- (b.) We integrate  $d^2\theta d^2\bar{\theta}$  at each vertex and in momentum space we have loop integrals  $(2\pi)^{-4} \int d^4p$  for each loop, and an overall factor  $(2\pi)^4 \delta^{(4)}(\sum k_{ext})$ .
- (c.) To obtain the effective action  $\Gamma$ , we compute one-particle-irreducible graphs. For each external line with outgoing momentum  $k_i$ , we multiply by a factor  $\int d^4k_i (2\pi)^{-4} \Psi(k_i)$  where  $\Psi$  stands for any of the fields in the effective action. For each external chiral or antichiral line, we have a  $\Phi$  or  $\bar{\Phi}$  factor, but no  $\bar{D}^2$  or  $D^2$  factor
- (d.) There may be symmetry factors associated with certain graphs.

#### 4.7 Review of the Functional Quantization of Yang-Mills Fields

As usual with a gauge theory, the main problem is that gauge-invariance implies that the gauge field propagator cannot be defined. A quick review of

ordinary Yang-Mills theory will suffice to show the problem.

$$\begin{aligned}\mathcal{S}_{YM} &= -\frac{1}{8} \int d^4x \operatorname{tr}[f_{\underline{a}\underline{b}}f_{\underline{a}\underline{b}}] \quad , \\ f_{\underline{a}\underline{b}} &= \partial_{\underline{a}}A_{\underline{b}} - \partial_{\underline{b}}A_{\underline{a}} - i[A_{\underline{a}}, A_{\underline{b}}] \quad .\end{aligned}\tag{362}$$

The infinitesimal gauge transformation is

$$\delta_G A_{\underline{a}} = \partial_{\underline{a}}\omega - i[A_{\underline{a}}, \omega] \quad ,\tag{363}$$

and the purely quadratic part of the action after an integration-by-parts has the form  $A^{\underline{a}}[\eta_{\underline{a}\underline{b}}p^2 - p_{\underline{a}}p_{\underline{b}}]A^{\underline{b}} \equiv A^{\underline{a}}\Delta_{\underline{a}\underline{b}}A^{\underline{b}}$  the object between these two fields is a  $4 \times 4$  matrix

$$\begin{aligned}\Delta_{\underline{a}\underline{b}} &= \begin{pmatrix} |p|^2 + (p_0)^2 & p_0 p_1 & p_0 p_2 & p_0 p_3 \\ p_1 p_0 & |p|^2 - (p_1)^2 & p_1 p_2 & p_1 p_3 \\ p_2 p_0 & p_2 p_1 & |p|^2 - (p_2)^2 & p_2 p_3 \\ p_3 p_0 & p_3 p_1 & p_3 p_2 & |p|^2 - (p_3)^2 \end{pmatrix} \quad , \\ |p|^2 &= - (p_0)^2 + (p_1)^2 + (p_2)^2 + (p_3)^2 \quad .\end{aligned}\tag{364}$$

After a direct calculation we find  $\det(\Delta_{\underline{a}\underline{b}}) = 0$  so it cannot be inverted in order to define a propagator for the gauge field. There is also an indirect way to see that this operator cannot be inverted. It is generally true that if an operator possesses a null space, then that operator has no inverse.

This in turn implies the need for a gauge-fixing term which allows the definition of a propagator for  $A_{\underline{a}}$ . However, this also permits the unphysical (gauge) component of  $A_{\underline{a}}$  propagate. This is fixed by BRST ghost and antighost introduction. These fields cancel out the effects of the unphysical components of  $A_{\underline{a}}$ . So in the end the generator  $Z(J)$  is defined by

$$\begin{aligned}Z(J) &= N' \int [\mathcal{D}A_{\underline{a}}][\mathcal{D}c][\mathcal{D}c'] \exp[\mathcal{S}_{eff} + \mathcal{S}_J] \quad , \\ \mathcal{S}_{eff} &= \mathcal{S}_{YM} - \int d^4x \operatorname{tr}\left[\frac{1}{\alpha}[F(A_{\underline{a}})]^2 + i \int d^4x' c'(x')\left(\frac{\delta F(\delta_G A)}{\delta \omega(x')}\right)c(x')\right] \quad , \\ \mathcal{S}_J &= \int d^4x \operatorname{tr}[A_{\underline{a}}J^{\underline{a}}] \quad ,\end{aligned}\tag{365}$$

where  $F(A_{\underline{a}})$  is a function that is not gauge invariant.

#### 4.8 Functional Quantization of Yang-Mills Superfields

We now wish to repeat this for the 4D,  $N = 1$  supersymmetric Yang-Mills theory. The gauge invariant action for the 4D,  $N = 1$  supersymmetric Yang-Mills theory may be written as

$$\begin{aligned} \mathcal{S}_{YM} &= \frac{1}{2} \left\{ \text{tr} \left[ \int d^6 z W^2 \right] + \text{h.c.} \right\} \\ &= -\frac{1}{4} \left\{ \text{tr} \left[ \int d^8 z (e^{-V} D^\alpha e^V) (\overline{D}^2 (e^{-V} D^\alpha e^V)) \right] + \text{h.c.} \right\} \quad (366) \\ &\approx \frac{1}{2} \text{tr} \left[ \int d^8 z (V D^\alpha \overline{D}^2 D_\alpha V) \right] + 0(V^3) \quad . \end{aligned}$$

To reach the second line we have used the non-abelian definition of  $W_\alpha$  (i.e.  $W_\alpha = [\overline{D}^2 (e^{-V} D^\alpha e^V)]$ ) along with the definition  $\int d^8 z = \int d^6 z \overline{D}^2$ . So the operator between the quadratic  $V$ -terms is given  $0 = D^\alpha \overline{D}^2 D_\alpha$ . Due to the identities,

$$D^\alpha \overline{D}^2 D_\alpha D^2 = 0 \quad , \quad D^\alpha \overline{D}^2 D_\alpha \overline{D}^2 \quad , \quad (367)$$

this operator is seen to possess null-spaces and is not invertible. For the first of these, we already saw a derivation. For the second one we have

$$\begin{aligned} D^\alpha \overline{D}^2 D_\alpha \overline{D}^2 &= \frac{1}{2} D^\alpha \overline{D}^2 D_\alpha \overline{D}^{\dot{\beta}} \overline{D}_{\dot{\beta}} = -\frac{1}{2} D^\alpha \overline{D}^2 D_\alpha \overline{D}_{\dot{\beta}} \overline{D}^{\dot{\beta}} \\ &= \frac{1}{2} D^\alpha \overline{D}^2 \overline{D}_{\dot{\beta}} D_\alpha \overline{D}^{\dot{\beta}} - i \frac{1}{2} \partial_{\alpha\dot{\beta}} D^\alpha \overline{D}^2 \overline{D}^{\dot{\beta}} = 0 \quad , \end{aligned} \quad (368)$$

since  $\overline{D}^2 \overline{D}^{\dot{\beta}} = 0$ .

The gauge variation of the superspace Yang-Mill gauge field is

$$\delta_G V = -i(\Lambda - \overline{\Lambda}) - i \frac{1}{2} L_V (\Lambda + \overline{\Lambda}) + i f(\frac{1}{2} L_V) (\Lambda - \overline{\Lambda}) \quad , \quad (369)$$

where the transcendental function  $f(x)$  is  $f(x) = 1 - x \coth(x)$ . This transformation law may be compared with that of ordinary Yang-Mills theory (150) which when written in term of forms becomes

$$\delta_G A = d\omega - i[A, \omega] = d\omega - i L_A \omega \quad . \quad (370)$$

Written in this form the similarities are obvious. The only major difference between the gauge variations of ordinary Yang-Mills theory and 4D,  $N = 1$  superspace Yang-Mills theory is that the latter involves a transcendental *not* linear function of the  $L$ -operator.

So like ordinary Yang-Mills theory the supersymmetric theory requires a gauge fixing term be introduced. The simplest choice is

$$\mathcal{S}_{GF} = -\frac{1}{\alpha} \text{tr} \left[ \int d^8 z (D^2 V) (\overline{D}^2 V) \right] , \quad (371)$$

and this is the direct analog of  $-\frac{1}{\alpha} \text{tr} \left[ \int d^4 x (\partial^a A_a)^2 \right]$ . Above we have found the explicit form of  $\delta_G V$ , and this can be used to calculate  $\delta_G \mathcal{S}_{GF}$ . After replacing the gauge parameter by chiral ghosts  $c$  and introducing the anti-ghost chiral superfields  $c'$  one finds

$$\mathcal{S}_{FP} = -\frac{1}{2\alpha} \text{tr} \left[ \int d^8 z (c' + \bar{c}') L_V \left[ (c + \bar{c}) + \coth\left(\frac{1}{2} L_V\right) (c - \bar{c}) \right] \right] . \quad (372)$$

where the antighosts  $c'$  and not related to the ghosts  $c$ . This term defines the propagator and interactions for the ghost and antighost superfields. The purely quadratic  $V$ -terms are now

$$-\frac{1}{2} V \left[ D^\alpha \overline{D}^2 D_\alpha - \frac{1}{\alpha} (D^2 \overline{D}^2 + \overline{D}^2 D^2) \right] V . \quad (373)$$

If we pick  $\alpha = 1$  (Fermi-Feynman gauge) and after some more  $D$ -algebra this becomes

$$-\frac{1}{2} V \square V , \quad (374)$$

and if we look at the part of the action that is pure quadratic in ghost-antighost we find

$$\mathcal{S}_{FP}^{(2)} = \int d^8 z \text{tr} \left[ \bar{c}' c - c' \bar{c} \right] . \quad (375)$$

So we conclude that the  $V$  propagator is

$$V V : \quad -\frac{1}{p^2} \delta^{(8)}(z - z') . \quad (376)$$

and since  $c$  and  $c'$  are chiral superfields

$$\begin{aligned} \bar{c}' c : & \quad -\frac{1}{p^2} \delta^{(4)}(\theta - \theta') \\ \bar{c} c' : & \quad -\frac{1}{p^2} \delta^{(4)}(\theta - \theta') \end{aligned} \quad (377)$$

The real magic of using supergraphs only begins at this point. The formal mathematical expressions that we have seen in this lecture can be associated with diagrams precisely like ordinary quantum field theory. In particular the

graphs associated with the structures that we have seen are called “Feynman supergraphs” and a single Feynman supergraph is equivalent to the Feynman graphs of all the component fields contained in the superfield. This is a topic treated in numbers of text books on the subject.

## 5 Fifth Lecture: Senior 4D, N = 1 Superfield Theory

### Introduction

In this final lecture we discuss the theory of 4D, N = 1 supergravity. The presentation includes both aspects of superfield and component descriptions. In terms of superfields, it is shown that 4D, N = 1 supergravity is a natural synthesis of what we have seen as the structure of supersymmetric Yang-Mills theory and ordinary gravity. So a complete structure of constraints as well as their solution, very similar to that of the supersymmetric Yang-Mills case, appears.

### Lecture

#### 5.1 Supergravitation Orientation

In lecture two, the superspace structure of 4D, N = 1 supersymmetric Yang-Mills theory was presented. A superspace connection  $\Gamma_{\underline{A}}^{\mathcal{I}}$  was defined and used to introduce a supergeometrical Yang-Mills covariant derivative,

$$\begin{aligned}\nabla_{\underline{A}} &\equiv D_{\underline{A}} - i\Gamma_{\underline{A}}^{\mathcal{I}} t_{\mathcal{I}} \quad , \\ D_{\underline{A}} &\equiv \left( D_{\alpha}, \overline{D}_{\dot{\alpha}}, \partial_{\underline{a}} \right) \quad , \\ \Gamma_{\underline{A}}^{\mathcal{I}} &\equiv \left( \Gamma_{\alpha}^{\mathcal{I}}, \Gamma_{\dot{\alpha}}^{\mathcal{I}}, \Gamma_{\underline{a}}^{\mathcal{I}} \right) \quad ,\end{aligned}\tag{378}$$

where  $t_{\mathcal{I}}$  denote Lie algebraic generators. This can be used to define field strength superfields  $F_{\underline{A}\underline{B}}^{\mathcal{I}}$ .

$$\left[ \nabla_{\underline{A}}, \nabla_{\underline{B}} \right] = C_{\underline{A}\underline{B}}^{\mathcal{C}} \nabla_{\underline{C}} - i F_{\underline{A}\underline{B}}^{\mathcal{I}} t_{\mathcal{I}} \quad .\tag{379}$$

In lecture three, we saw that the theory of general relativity, although formulated by Einstein in purely geometric terms, has a natural interpretation in terms of concepts that arise from Yang-Mills gauge theory. The most important point of our discussion was that we were able to identify the Lie algebra generators that underlie general relativity

$$t_{\mathcal{I}} \rightarrow \left( P_{\underline{m}}, \mathcal{M}_{\underline{a}\underline{b}} \right) \quad ,\tag{380}$$

where  $P_{\underline{m}}$  is the usual momentum operator from quantum mechanics and (or alternately  $\mathcal{M}_{\alpha\beta}$  and  $\overline{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}$ ) are the abstract Lie algebra generators of spin-angular momentum operators also from quantum mechanics. This particular

suggestion for the Lie-algebraic basis for general relativity was first made in 1978<sup>7</sup>. Once this Lie-algebraic basis for general relativity is accepted, the structure of a covariant derivative follows

$$\nabla_{\underline{a}} \equiv e_{\underline{a}}^{\underline{m}} \partial_{\underline{m}} + \frac{1}{2} \omega_{\underline{a}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} \quad , \quad (381)$$

and its commutator is used to define field strengths

$$\left[ \nabla_{\underline{a}}, \nabla_{\underline{b}} \right] = t_{\underline{a}\underline{b}}^{\underline{c}} \nabla_{\underline{c}} + \frac{1}{2} r_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} \quad , \quad (382)$$

where the first is “the torsion tensor” and the second is “the Riemann curvature tensor.”

In the 4D, N = 1 superspace Yang-Mills theory, the field strength  $F_{\underline{A}\underline{B}}^{\mathcal{I}}$  can be separated into various irreducible Lorentz representations

$$F_{\underline{A}\underline{B}}^{\mathcal{I}} = \begin{pmatrix} F_{\alpha\beta}^{\mathcal{I}} & F_{\alpha\beta}^{\cdot\mathcal{I}} & F_{\alpha\dot{b}}^{\mathcal{I}} \\ F_{\alpha\beta}^{\cdot\mathcal{I}} & F_{\alpha\beta}^{\mathcal{I}} & F_{\alpha\dot{b}}^{\cdot\mathcal{I}} \\ F_{\underline{a}\beta}^{\mathcal{I}} & F_{\underline{a}\beta}^{\cdot\mathcal{I}} & F_{\underline{a}\dot{b}}^{\mathcal{I}} \end{pmatrix} \quad , \quad (383)$$

and a minimal theory exists by imposing algebraic conditions that some of these irreducible Lorentz representations are constrained to vanish:  $F_{\alpha\beta}^{\mathcal{I}} = F_{\alpha\beta}^{\cdot\mathcal{I}} = F_{\alpha\dot{b}}^{\mathcal{I}} = F_{\alpha\dot{b}}^{\cdot\mathcal{I}} = 0$ . The solution to these constraints implies that all of the connection superfields can be defined in terms of a more fundamental object, the pseudo-scalar pre-potential superfield  $V^{\mathcal{I}}$ .

In general relativity, also one of the field strengths is constrained to vanish:  $t_{\underline{a}\underline{b}\underline{c}} = 0$ . The solution to this condition implies that the spin-connection can be defined in terms of the anholonomy. So superspace Yang-Mills theory and general relativity are both gauge theories where some of the a priori non-vanishing field strengths are constrained to vanish.

## 5.2 Superspace Supergravity Covariant Derivatives

From all of this it should be clear that in order to describe a 4D, N = 1 theory that is analogous to general relativity, we need to combine the features of the two distinct theories discussed above. In other words, the Lie algebraic generators of the 4D, N = 1 supersymmetric covariant Yang-Mills derivative need to be replaced by the analog of  $(P_{\underline{m}}, \mathcal{M}_{\underline{a}\underline{b}})$ . From lecture one we know these turn out to be  $(Q_{\mu}, \bar{Q}_{\dot{\mu}}, P_{\underline{m}}, \mathcal{M}_{\underline{a}\underline{b}})$ . However, for convenience sake and without loss of generality, we make the equivalent replacement of  $(D_{\mu}, \bar{D}_{\dot{\mu}}, P_{\underline{m}}, \mathcal{M}_{\underline{a}\underline{b}})$  in the superspace supergravity covariant derivative

$$\nabla_{\underline{A}} \equiv E_{\underline{A}}^{\underline{M}} D_{\underline{M}} + \frac{1}{2} \omega_{\underline{A}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} \quad , \quad (384)$$

where instead of  $(Q_\mu, \bar{Q}_\mu, P_m)$  we use  $(D_\mu, \bar{D}_\mu, \partial_m)$ . In this definition, the quantity  $E_{\underline{A}}^{\underline{M}}(\theta, \bar{\theta}, x)$  is known as the “super vielbein” and  $\omega_{\underline{A}\underline{c}\underline{d}}(\theta, \bar{\theta}, x)$  is known as the “super spin-connection.”

The graded commutator of this operator is used to define the field strengths

$$\left[ \nabla_{\underline{A}}, \nabla_{\underline{B}} \right] = T_{\underline{A}\underline{B}}^{\underline{C}} \nabla_{\underline{C}} + \frac{1}{2} R_{\underline{a}\underline{b}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} \quad , \quad (385)$$

where  $T_{\underline{A}\underline{B}}^{\underline{C}}$  is called “the torsion super-tensor” and  $R_{\underline{a}\underline{b}\underline{c}}^{\underline{d}}$  is called “the Riemann curvature supertensor.” These may be expressed in terms of  $E_{\underline{A}}^{\underline{M}}$ ,  $\omega_{\underline{A}\underline{c}\underline{d}}$  and their derivatives. For the torsion

$$\begin{aligned} T_{\underline{A}\underline{B}}^{\underline{C}} &= C_{\underline{A}\underline{B}}^{\underline{C}} - \omega_{\underline{A}\underline{B}}^{\underline{C}} + (-1)^{\underline{A}\underline{B}} \omega_{\underline{B}\underline{A}}^{\underline{C}} \quad , \\ E_{\underline{A}} &\equiv E_{\underline{A}}^{\underline{M}} D_{\underline{M}} \quad , \quad \left[ E_{\underline{A}}, E_{\underline{B}} \right] = C_{\underline{A}\underline{B}}^{\underline{C}} E_{\underline{C}} \quad , \\ C_{\underline{A}\underline{B}}^{\underline{C}} &= [E_{\underline{A}}^{\underline{M}} (D_{\underline{M}} E_{\underline{B}}^{\underline{N}}) - (-1)^{\underline{A}\underline{B}} E_{\underline{B}}^{\underline{M}} (D_{\underline{M}} E_{\underline{A}}^{\underline{N}})] E_{\underline{N}}^{\underline{C}} \quad , \end{aligned} \quad (386)$$

where  $C_{\underline{A}\underline{B}}^{\underline{C}}$  is the super-anholonomy. In writing the torsion supertensor it appears that  $\omega_{\underline{A}\underline{B}}^{\underline{C}}$  is a general super matrix in its  $\underline{B}$  and  $\underline{C}$  indices. This is not the case. Instead we have

$$\omega_{\underline{A}\underline{B}}^{\underline{C}} = \begin{pmatrix} \omega_{\underline{A}\underline{\beta}}^{\dot{\gamma}} & 0 & 0 \\ 0 & \omega_{\underline{A}\underline{\beta}}^{\dot{\gamma}} & 0 \\ 0 & 0 & \omega_{\underline{A}\underline{b}}^{\underline{c}} \end{pmatrix} \quad , \quad (387)$$

and this is a consequence since we only included  $\mathcal{M}_{\underline{a}\underline{b}}$  (or equivalently  $\mathcal{M}_{\alpha\beta}$  and  $\bar{\mathcal{M}}_{\dot{\alpha}\dot{\beta}}$ ) in the superspace supergravity covariant derivative. (Here we may also write  $\omega_{\underline{A}\underline{b}\underline{c}} = C_{\dot{\beta}\dot{\gamma}}^{\dot{\alpha}} \omega_{\underline{A}\underline{\beta}\dot{\gamma}} + C_{\beta\gamma}^{\dot{\alpha}} \omega_{\underline{A}\underline{\beta}\dot{\gamma}}$  as is our usual custom.) This further implies that for some values of the indices  $\underline{A}$ ,  $\underline{B}$  and  $\underline{C}$ , the torsion is identical to the anholonomy. For example the spin connection superfield does not appear in

$$\begin{aligned} T_{\alpha\beta}^{\dot{\gamma}} &= C_{\alpha\beta}^{\dot{\gamma}} \quad , \quad T_{\alpha\beta}^{\underline{c}} = C_{\alpha\beta}^{\underline{c}} \quad , \\ T_{\alpha\dot{\beta}}^{\underline{c}} &= C_{\alpha\dot{\beta}}^{\underline{c}} \quad , \quad T_{\alpha\underline{b}}^{\dot{\gamma}} = C_{\alpha\underline{b}}^{\dot{\gamma}} \quad , \\ T_{\underline{a}\underline{b}}^{\gamma} &= C_{\underline{a}\underline{b}}^{\gamma} \quad , \end{aligned} \quad (388)$$

and as well for the complex conjugates of these irreducible Lorentz components of the torsion supertensor.

However, for other components of the torsion supertensor, the spin-connection



superfield appears as in

$$\begin{aligned}
T_{\alpha\beta}{}^{\gamma} &= C_{\alpha\beta}{}^{\gamma} + \omega_{\alpha\beta}{}^{\gamma} + \omega_{\beta\alpha}{}^{\gamma} \quad , \\
T_{\alpha\dot{\beta}}{}^{\dot{\gamma}} &= C_{\alpha\dot{\beta}}{}^{\dot{\gamma}} + \omega_{\alpha\dot{\beta}}{}^{\dot{\gamma}} \quad , \\
T_{\alpha\bar{b}}{}^{\gamma} &= C_{\alpha\bar{b}}{}^{\gamma} + \omega_{\bar{b}\alpha}{}^{\gamma} \quad , \\
T_{\alpha\bar{b}}{}^{\underline{c}} &= C_{\alpha\bar{b}}{}^{\underline{c}} - \omega_{\alpha\bar{b}}{}^{\underline{c}} \quad , \\
T_{\underline{a}\bar{b}}{}^{\underline{c}} &= C_{\underline{a}\bar{b}}{}^{\underline{c}} - \omega_{\underline{a}\bar{b}}{}^{\underline{c}} + \omega_{\bar{b}\underline{a}}{}^{\underline{c}} \quad ,
\end{aligned} \tag{389}$$

and in all the complex conjugates of these equations.

### 5.3 Superspace Supergravity Conventional Constraints

In general relativity, the constraint  $t_{\underline{a}\bar{b}\underline{c}} = 0$  allowed the spin connection to be expressed in terms of the anholonomy. A similar stratagem works here where

$$\begin{aligned}
T_{\alpha\beta}{}^{\gamma} = 0 \quad , \quad \rightarrow \quad \omega_{\alpha\beta}{}^{\gamma} &= -\frac{1}{2} [C_{\alpha\beta}{}^{\gamma} - C_{\beta\gamma}{}^{\alpha} + C_{\gamma\alpha}{}^{\beta}] \quad , \\
T_{\alpha\dot{\beta}}{}^{\dot{\gamma}} = 0 \quad , \quad \rightarrow \quad \omega_{\alpha\dot{\beta}}{}^{\dot{\gamma}} &= -\frac{1}{2} [C_{\alpha\dot{\beta}}{}^{\dot{\gamma}} - C_{\dot{\beta}\dot{\gamma}}{}^{\alpha} + C_{\dot{\gamma}\alpha}{}^{\dot{\beta}}] \quad , \\
T_{\alpha(\dot{\beta}\dot{\gamma})} = 0 \quad , \quad \rightarrow \quad \omega_{\alpha(\dot{\beta}\dot{\gamma})} &= -\frac{1}{2} C_{\alpha(\dot{\beta}\dot{\gamma})} \quad , \\
T_{\underline{a}(\beta\gamma)} = 0 \quad , \quad \rightarrow \quad \omega_{\underline{a}(\beta\gamma)} &= C_{\underline{a}(\beta\gamma)} \quad , \\
T_{\underline{a}\bar{b}}{}^{\underline{c}} = 0 \quad , \quad \rightarrow \quad \omega_{\underline{a}\bar{b}}{}^{\underline{c}} &= \frac{1}{2} [C_{\underline{a}\bar{b}}{}^{\underline{c}} - C_{\bar{b}\underline{a}}{}^{\underline{c}} + C_{\underline{c}\bar{a}}{}^{\underline{b}}] \quad .
\end{aligned} \tag{390}$$

Thus,  $\omega_{\underline{a}\bar{b}\underline{c}}$  is completely determined just as in general relativity.

At this stage, the only independent degrees of freedom in  $\nabla_{\underline{A}}$  that remain are those in  $E_{\underline{A}}{}^{\underline{M}}$ . But should all of these be independent? Recall the following analogy

$$\nabla_{\underline{A}}^{(YM)} = D_{\underline{A}} - i\Gamma_{\underline{A}}{}^{\mathcal{I}} t_{\mathcal{I}} \sim E_{\underline{A}} = E_{\underline{A}}{}^{\underline{M}} D_{\underline{M}} \quad . \tag{391}$$

In the Yang-Mills theory, the vectorial connection  $\Gamma_{\underline{a}}{}^{\mathcal{I}}$  was defined in terms of  $\Gamma_{\alpha}{}^{\mathcal{I}}$ ,  $\Gamma_{\dot{\alpha}}{}^{\mathcal{I}}$  and their derivatives. This is motivation to attempt to do the same for supergravity. It turns out that this is possible but in a slightly subtle way.

We note

$$\begin{aligned}
[E_\alpha, E_{\dot{\alpha}}] &= C_{\alpha\dot{\alpha}}^{\underline{\varepsilon}} E_{\underline{\varepsilon}} + C_{\alpha\dot{\alpha}}^{\dot{\gamma}} E_{\dot{\gamma}} + C_{\alpha\dot{\alpha}}^{\dot{\gamma}} E_{\dot{\gamma}} \\
\rightarrow C_{\alpha\dot{\alpha}}^{\underline{\varepsilon}} E_{\underline{\varepsilon}} &= [E_\alpha, E_{\dot{\alpha}}] - C_{\alpha\dot{\alpha}}^{\dot{\gamma}} E_{\dot{\gamma}} - C_{\alpha\dot{\alpha}}^{\dot{\gamma}} E_{\dot{\gamma}} \\
\rightarrow C_{\alpha\dot{\alpha}}^{\underline{\varepsilon}} E_{\underline{\varepsilon}} &= [E_{\underline{A}}^{\underline{M}} (D_{\underline{M}} E_{\underline{B}}^{\underline{N}}) - (-1)^{AB} E_{\underline{B}}^{\underline{M}} (D_{\underline{M}} E_{\underline{A}}^{\underline{N}})] D_{\underline{N}} \\
&\quad - C_{\alpha\dot{\alpha}}^{\dot{\gamma}} E_{\dot{\gamma}} - C_{\alpha\dot{\alpha}}^{\dot{\gamma}} E_{\dot{\gamma}} .
\end{aligned} \tag{392}$$

If  $C_{\alpha\dot{\alpha}}^{\underline{\varepsilon}}$  is a constant, then this equation defines  $E_{\underline{\varepsilon}}$  as a function of  $E_\gamma$ ,  $E_{\dot{\gamma}}$  and their derivatives. The value of this constant can be deduced by considering the “flat limit”  $E_{\underline{\varepsilon}} \rightarrow \partial_{\underline{\varepsilon}}$ ,  $E_\gamma \rightarrow D_\gamma$  and  $E_{\dot{\gamma}} \rightarrow \overline{D}_{\dot{\gamma}}$  which possesses no supergravity fields. This tells us

$$C_{\alpha\dot{\alpha}}^{\underline{\varepsilon}} = i\delta_\alpha^\gamma \delta_{\dot{\alpha}}^{\dot{\gamma}} . \tag{393}$$

The set of constraints in (390) and (393) have been named “conventional constraints”<sup>6</sup>. This name is suggested for a couple of reasons. First, these are conventional in the sense that they define some *a priori* independent gauge fields in terms of others and are thus a ‘convention’ for the definitions of some gauge fields. These constraints are also conventional in the sense that they are the usual type of constraint that was seen in global supersymmetrical Yang-Mills theory and general relativity.

#### 5.4 Superspace Supergravity Representation-Preserving Constraints

Continuing we have the “integrability conditions”<sup>7</sup> coming from chiral matter superfields. In the case of the superspace Yang-Mills theory these conditions arose due the demand that “covariant chiral” superfields should exist even in the presence of gauge symmetries. We may invoke the same argument in the presence of the symmetries of supergravity. In the absence of supergravity, chiral superfields are defined by  $\overline{D}_{\dot{\alpha}}\Phi = 0$ . However, in the presence of supergravity we know that  $\Phi$  can undergo K-gauge transformations

$$(\Phi)' = e^K \Phi , \quad K \equiv K^{\underline{M}} D_{\underline{M}} = K^\mu D_\mu + K^{\dot{\mu}} \overline{D}_{\dot{\mu}} + K^{\underline{m}} \partial_{\underline{m}} . \tag{394}$$

These transformations are precisely the generalization of those that we saw in our discussion of the Lie algebraic formulation of theories of ordinary gravitation.

It is clearly indicated that in order to possess this as a symmetry, the definition of a chiral superfield in the presence of a supergravity should be  $\nabla_{\dot{\alpha}}\Phi = 0$ . (Since  $\Phi$  is a Lorentz scalar, the connection superfields do not appear in this

condition.) We can thus repeat the derivation of the ‘integrability constraints’ but now applied to the superspace supergravity covariant derivative.

$$\begin{aligned}
\nabla_{\dot{\alpha}} \Phi &= 0 \rightarrow \nabla_{\dot{\alpha}} \nabla_{\dot{\beta}} \Phi = 0 \rightarrow [\nabla_{\dot{\alpha}}, \nabla_{\dot{\beta}}] \Phi = 0 \rightarrow \\
T_{\alpha \dot{\alpha}}^{\underline{\varepsilon}} (\nabla_{\underline{\varepsilon}} \Phi) + T_{\alpha \dot{\alpha}}^{\dot{\gamma}} (\nabla_{\dot{\gamma}} \Phi) + T_{\alpha \dot{\alpha}}^{\dot{\gamma}} (\nabla_{\dot{\gamma}} \Phi) &= 0 \rightarrow \\
T_{\alpha \dot{\alpha}}^{\underline{\varepsilon}} (\nabla_{\underline{\varepsilon}} \Phi) + T_{\alpha \dot{\alpha}}^{\dot{\gamma}} (\nabla_{\dot{\gamma}} \Phi) &= 0 \rightarrow \\
C_{\alpha \dot{\alpha}}^{\underline{\varepsilon}} (\nabla_{\underline{\varepsilon}} \Phi) + C_{\alpha \dot{\alpha}}^{\dot{\gamma}} (\nabla_{\dot{\gamma}} \Phi) &= 0 \quad .
\end{aligned} \tag{395}$$

This is just like the Yang-Mills case and has a similar solution

$$C_{\alpha \dot{\alpha}}^{\underline{\varepsilon}} = 0 \quad , \quad C_{\alpha \dot{\alpha}}^{\dot{\gamma}} = 0 \quad . \tag{396}$$

In turn these constraints imply that

$$[E_{\alpha} , E_{\beta}] = C_{\alpha \beta}^{\gamma} E_{\gamma} \quad . \tag{397}$$

This equation means that the differential operators  $E_{\alpha}$  form a closed set under graded commutation. There is a result from differential geometry due to Frobenius which implies that the solution to these constraints is also like supersymmetric Yang-Mills theory

$$E_{\alpha} = \Psi N_{\alpha}^{\mu} e^{\frac{1}{2}U} D_{\mu} e^{-\frac{1}{2}U} \quad , \quad U \equiv U^{\mu} D_{\mu} + \overline{U}^{\dot{\mu}} \overline{D}_{\dot{\mu}} + U^{\underline{m}} \partial_{\underline{m}} \quad . \tag{398}$$

In writing this solution, we have directly gone to a gauge that is analogous to that in (186).

The superfield  $U^{\underline{M}}$ , first suggested by Ogievetsky and Sokatchev<sup>22</sup> who called it the “axial vector superfield,” is to supergravity as  $V^{\mathcal{I}}$  is to supersymmetric Yang-Mills theory. In the expression for  $E_{\alpha}$ ,  $\Psi$  is an arbitrary complex scalar superfield (called the “superfield scale compensator” or “the density-type conformal compensator”) and  $N_{\alpha}^{\mu}$  (called “the superfield Lorentz compensator”) is an arbitrary complex matrix superfield restricted to satisfy the constraint

$$\det(N_{\alpha}^{\mu}) = 1 \quad . \tag{399}$$

(Note that  $N_{\alpha}^{\mu}$  is not a supermatrix.) These two superfields have no analogs within the context of supersymmetric Yang-Mills theory. Although a complete proof is beyond the level of exposition in these lectures, in addition it turns out that  $U^{\mu}$ ,  $\overline{U}^{\dot{\mu}}$  and  $N_{\alpha}^{\mu}$  may all be “gauged away” in an appropriate W-Z gauge. Thus these are extraneous. The constraints in (396) are also called “representation-preserving constraints” since they are required in order for the

chiral superfield representation of rigid superspace to exist in the presence of superspace supergravity.

The result in (398) is very revelatory. The curved Salam-Strathdee superspace (also called a Wess-Zumino superspace) definitely possess a “supergeometry” described by the superspace supergravity covariant derivative  $\nabla_{\underline{A}}$  which is expressed in terms of the vielbein  $E_{\underline{A}}^{\underline{M}}$  and superfield spin-connection  $\omega_{\underline{A}}^{\underline{d}\underline{e}}$ . All of this is the simplest sort of extrapolation of Riemannian geometry to superspace that should be expected. After imposing conventional constraints, all geometrical quantities are expressed solely in terms of  $E_{\alpha}^{\underline{M}}$ . After imposing (396) we enter the domain of “pre-geometry.” This is a realm that, although well understood for 4D,  $N = 1$  superspace, remains essentially not understood for most superspaces and therefore by implication for all theories that contain supergravity. Gaining an understanding of pre-geometry is the most important challenge to superstrings, heterotic strings and all other such constructions. For it is only with the understanding of pre-geometry that the superspace analog of the equivalence principle reveals itself completely.

### 5.5 Superspace Supergravity Conformal-Breaking Constraints

From our discussion in the first lecture we know that a general superfield, such as  $\Psi$  is not the smallest representation of supersymmetry. It would be nice to try to make  $\Psi$  into some type of chiral superfield<sup>23</sup>. This can actually be done if the constraints below are imposed.

$$T_{\alpha\underline{b}}^{\underline{c}} = T_{\alpha\underline{b}}^{\underline{c}} = 0 \quad . \quad (400)$$

The solution to this final constraint is of the form

$$\Psi = \Psi(U^{\underline{m}}, \varphi) \quad , \quad (401)$$

where  $\varphi$  is a chiral superfield in the sense that  $\overline{D}_{\dot{\mu}}\varphi = 0$ . The explicit form of the function  $\Psi(U^{\underline{m}}, \varphi)$  can be found in the last two books of the bibliography.

### 5.6 The WZ Gauge and Supergravity Component Fields

All of the physics of (this version of) supergravity comes solely from  $U^{\underline{m}}$  and  $\varphi$ . As in the Yang-Mills case, there exist a WZ gauge wherein

$$\begin{aligned} U^{\underline{m}}(z) = & \theta^{\alpha}\bar{\theta}^{\dot{\alpha}} e_{\underline{a}}^{\underline{m}}(x) - (i\bar{\theta}^2\theta^{\alpha}\psi_{\alpha}^{\underline{b}}(x) + \text{h.c.}) e_{\underline{b}}^{\underline{m}}(x) + \\ & \theta^2\bar{\theta}^2(A^{\underline{a}}(x) - \frac{1}{3}\epsilon^{\underline{a}\underline{b}\underline{c}\underline{d}}c_{\underline{b}\underline{c}\underline{d}}) e_{\underline{a}}^{\underline{m}}(x) \quad . \end{aligned} \quad (402)$$

and in this same gauge<sup>17</sup>

$$\varphi = e^{-\frac{1}{3}} \{ 1 - \theta^2 [S(x) + iP(x)] \} . \quad (403)$$

Thus, the component fields of (minimal) 4D, N = 1 supergravity are

- $e_a^m(x)$  graviton spin - 2
- $\psi_a^\mu(x)$  gravitino spin - 3/2
- $A_a(x)$  axial-vector auxiliary field
- $S(x)$  scalar auxiliary field spin-0
- $P(x)$  pseudoscalar auxiliary field spin-0

These are the fields that remain in the WZ gauge and are therefore the only component fields that occur in the component-level formulations. A complete set of constraint as imposed in our discussion was first suggested by J. Wess and B. Zumino<sup>24</sup>. In recognition of this they are often called the “Wess-Zumino supergravity constraints.”

### 5.7 Graded Commutator Algebra of the Supergravity Supercovariant Derivative and Supergravity Action

Independently, Warren Siegel and I developed the complete description<sup>17</sup> of  $\nabla_{\underline{A}}$  in terms of  $U^{\underline{m}}$  and  $\varphi$  in the period of 1977-1979. We also used a slightly different set of constraints from Wess and Zumino,

$$T_{\underline{a}\underline{b}}^{\underline{c}} = 0 \rightarrow R_{\alpha\dot{\alpha}\underline{c}\underline{d}} = 0 . \quad (404)$$

This choice only determines  $\omega_{\underline{a}\underline{c}\underline{d}}$  in a slightly different way. In our system, the algebra of the supergravity covariant derivative takes the form

$$\begin{aligned} [\nabla_\alpha, \nabla_\beta] &= -2\bar{R}\mathcal{M}_{\alpha\beta} \quad , \quad [\nabla_\alpha, \bar{\nabla}_{\dot{\alpha}}] = i\nabla_{\underline{a}} \quad , \\ [\nabla_\alpha, \nabla_{\underline{b}}] &= -iC_{\alpha\beta}[\bar{R}\bar{\nabla}_{\dot{\beta}} - G^\gamma_{\dot{\beta}}\nabla_\gamma] - i(\bar{\nabla}_{\dot{\beta}}\bar{R})\mathcal{M}_{\alpha\beta} \\ &\quad + iC_{\alpha\beta}[\bar{W}_{\dot{\beta}\dot{\gamma}}^{\delta}\bar{\mathcal{M}}_{\delta}^{\dot{\gamma}} - (\nabla^\delta G_{\gamma\dot{\beta}})\mathcal{M}_{\delta}^{\gamma}] \quad , \\ [\nabla_{\underline{a}}, \nabla_{\underline{b}}] &= \{ [C_{\dot{\alpha}\dot{\beta}}W_{\alpha\beta}^{\gamma} + \frac{1}{2}C_{\alpha\beta}(\bar{\nabla}_{(\dot{\alpha}}G_{\dot{\beta}}^{\gamma)}) - \frac{1}{2}C_{\dot{\alpha}\dot{\beta}}(\nabla_{(\alpha}R)]\nabla_\gamma \\ &\quad + i\frac{1}{2}C_{\alpha\beta}G^{\gamma(\dot{\alpha}}\nabla_{\dot{\beta}}) - \frac{1}{2}C_{\alpha\beta}(\bar{\nabla}_{(\dot{\alpha}}\nabla_{\dot{\beta}}G_{\delta}^{\gamma)})]\mathcal{M}^{\gamma\delta} \\ &\quad - [C_{\dot{\alpha}\dot{\beta}}W_{\alpha\beta\gamma\delta} - \frac{1}{2}C_{\alpha\beta}(\bar{\nabla}_{(\dot{\alpha}}\nabla_{\dot{\beta}}G_{\delta}^{\gamma)})]\mathcal{M}^{\gamma\delta} \\ &\quad - i\frac{1}{2}C_{\dot{\alpha}\dot{\beta}}C_{\gamma(\alpha}[\frac{1}{4}(\nabla_{\beta)}^{\epsilon}G_{\delta\epsilon} + \nabla_\delta^{\epsilon}G_{\beta\epsilon}]\mathcal{M}^{\gamma\delta} \\ &\quad - \frac{1}{2}C_{\dot{\alpha}\dot{\beta}}C_{\gamma(\alpha}C_{\beta)\delta}[\bar{\nabla}^2\bar{R} + 2R\bar{R}]\mathcal{M}^{\gamma\delta} + \text{h.c.} \} \quad , \end{aligned} \quad (405)$$

the superfields  $W_{\alpha\beta\gamma}$ ,  $G_{\underline{a}}$  and  $R$  are to supergravity as  $W_{\alpha}{}^{\mathcal{I}}$  is to supersymmetric Yang-Mills theory. The superspace Bianchi identities exist in direct analogy with ordinary gravity and supersymmetrical Yang-Mills theory

$$\begin{aligned} 0 = & (-1)^{AC} [[\nabla_{\underline{A}}, \nabla_{\underline{B}}], \nabla_{\underline{C}}] + (-1)^{BA} [[\nabla_{\underline{B}}, \nabla_{\underline{C}}], \nabla_{\underline{A}}] \\ & + (-1)^{CB} [[\nabla_{\underline{C}}, \nabla_{\underline{A}}], \nabla_{\underline{B}}] , \end{aligned} \quad (406)$$

and they imply the following equation for the superfields  $W_{\alpha\beta\gamma}$ ,  $G_{\underline{a}}$  and  $R$

$$\begin{aligned} G_{\alpha\dot{\beta}} &= \overline{G}_{\alpha\dot{\beta}} , \quad \overline{\nabla}_{\dot{\beta}} R = \overline{\nabla}_{\dot{\beta}} W_{\alpha\beta\gamma} = 0 , \\ \nabla^{\alpha} R + \overline{\nabla}_{\dot{\alpha}} G^{\alpha\dot{\alpha}} &= 0 , \quad \nabla^{\beta} W_{\beta\gamma\delta} + i\frac{1}{2} \nabla_{(\gamma\delta} G_{\delta)}^{\dot{\delta}} = 0 , \\ \frac{1}{4!} \nabla_{(\alpha} W_{\beta\gamma\delta)} &= W_{\alpha\beta\gamma\delta} . \end{aligned} \quad (407)$$

In ordinary gravity, the Einstein-Hilbert action takes the form in (286) and in 4D, N=1 supersymmetric Yang-Mills theory the action takes the form in (141). So it is obviously of interest to know what is the form of the 4D, N=1 supergravity action. As first suggested by Siegel<sup>23</sup>, it is

$$\mathcal{S}_{SG} = -\frac{3}{\kappa^2} \int d^4x d^2\theta d^2\bar{\theta} E^{-1}(U^{\underline{m}}, \varphi) , \quad E \equiv sdet(E_{\underline{A}}^{\underline{M}}) . \quad (408)$$

As indicated in the equation above  $E(U^{\underline{m}}, \varphi)$  is a function of  $U^{\underline{m}}$  and  $\varphi$  alone. Variation of this action produces two equations

$$\begin{aligned} e_{\underline{a}}^m \frac{\delta \mathcal{S}_{SG}}{\delta U^{\underline{m}}} &= 0 \quad \rightarrow \quad G_{\underline{a}} = 0 , \\ \frac{\delta \mathcal{S}_{SG}}{\delta \varphi} &= 0 \quad \rightarrow \quad \overline{R} = 0 . \end{aligned} \quad (409)$$

Of course written in terms of superfields, the supergravity action looks (but hides lots) very simple. To get more insight into what's going on, a peek at components is warranted.

### 5.8 Component Supergravity Action

After using some techniques that will be explained later, we find (408) leads to

$$\begin{aligned} \mathcal{S}_{SG} = \frac{1}{\kappa^2} \int d^4x e^{-1} \Big[ & r(\omega(e, \psi, A), e) - i \epsilon^{\underline{a}\underline{b}\underline{c}\underline{d}} \overline{\psi}_{\underline{a}\dot{\beta}} \dot{\psi}_{\underline{c}\underline{d}\beta} \\ & - 3(S^2 + P^2) \Big] . \end{aligned} \quad (410)$$

This result (in a slightly different form) was first derived without its auxiliary field structure by Freedman, Ferrara and van Nieuwenhuizen<sup>25</sup> and later with the auxiliary field by Ferrara and van Nieuwenhuizen<sup>26</sup>. Above the spin-connection is given by

$$\begin{aligned} \omega(e, \psi, A) = & \omega_{\underline{a}\underline{b}\underline{c}}(e) + i \frac{1}{2} ( \psi_{[\underline{b}\alpha} \bar{\psi}_{\underline{c}]} \dot{\bar{\alpha}} + \psi_{[\underline{a}\beta} \bar{\psi}_{\underline{c}]} \dot{\bar{\beta}} - \psi_{[\underline{a}\gamma} \bar{\psi}_{\underline{b}]} \dot{\bar{\gamma}} ) \\ & - \frac{1}{2} \epsilon_{\underline{a}\underline{b}\underline{c}\underline{d}} A^{\underline{d}} , \end{aligned} \quad (411)$$

where  $\omega_{\underline{a}\underline{b}\underline{c}}(e)$  is the spin-connection (272) in ordinary general relativity and the “curl” of the gravitino is explicitly of the form

$$\begin{aligned} \psi_{\underline{a}\underline{b}}^{\gamma} &\equiv e_{[\underline{a}}^{\underline{m}} \left( \partial_{\underline{m}} \psi_{\underline{b}]}^{\gamma} - \omega_{\underline{m}\beta}^{\gamma}(e, \psi, A) \psi_{\underline{b}]}^{\beta} \right) - c_{\underline{a}\underline{b}}^{\underline{d}} \psi_{\underline{d}}^{\gamma} \\ &\equiv e_{[\underline{a}}^{\underline{m}} \left( \mathcal{D}_{\underline{m}}(e, \psi, A) \psi_{\underline{b}]}^{\gamma} \right) - c_{\underline{a}\underline{b}}^{\underline{d}} \psi_{\underline{d}}^{\gamma} \\ &\equiv \left( \mathcal{D}_{[\underline{a}}(e, \psi, A) \psi_{\underline{b}]}^{\gamma} \right) - c_{\underline{a}\underline{b}}^{\underline{d}} \psi_{\underline{d}}^{\gamma} . \end{aligned} \quad (412)$$

We can also define

$$\omega_{\underline{a}\underline{b}\underline{c}}(e, \psi, A) = \omega_{\underline{a}\underline{b}\underline{c}}(e, \psi) - \frac{1}{2} \epsilon_{\underline{a}\underline{b}\underline{c}\underline{d}} A^{\underline{d}} , \quad (413)$$

and use this to re-express the component level action in the form

$$\begin{aligned} \mathcal{S}_{SG} = & \frac{1}{\kappa^2} \int d^4x \, e^{-1} \left[ r(\omega(e, \psi), e) - i \epsilon^{\underline{a}\underline{b}\underline{c}\underline{d}} \bar{\psi}_{\underline{a}} \dot{\bar{\beta}} \psi_{\underline{c}\underline{d}\beta} \right. \\ & \left. - 3 (A^{\underline{a}} A_{\underline{a}} + S^2 + P^2) \right] . \end{aligned} \quad (414)$$

This is the almost universal form in which the supergravity action appears in the literature. Let us end this section by noting that this action has one final form in which it may be written,

$$\begin{aligned} \mathcal{S}_{SG} = & \frac{1}{\kappa^2} \int d^4x \, e^{-1} \left[ -\frac{1}{4} c^{\underline{a}\underline{b}\underline{c}} c_{\underline{a}\underline{b}\underline{c}} + \frac{1}{2} c^{\underline{a}\underline{b}\underline{c}} c_{\underline{c}\underline{a}\underline{b}} + (c^{\underline{a}\underline{b}})^2 \right. \\ & - i \epsilon^{\underline{a}\underline{b}\underline{c}\underline{d}} \bar{\psi}_{\underline{a}} \dot{\bar{\beta}} \hat{\psi}_{\underline{c}\underline{d}\beta} - 3 (A^{\underline{a}} A_{\underline{a}} + S^2 + P^2) \\ & \left. + \psi^4 - \text{terms} \right] . \end{aligned} \quad (415)$$

The gravitino field strength that appears here is defined by

$$\psi_{\underline{c}\underline{d}\beta} = \hat{\psi}_{\underline{c}\underline{d}\beta} + \psi^3 - \text{terms} . \quad (416)$$

The  $\psi^4$  – terms in the action arise from this substitution as well as writing

$$r(\omega(e, \psi), e) = r(\omega(e), e) + \psi^4 - \text{terms} \quad . \quad (417)$$

This form of the action is noteworthy for a couple of reasons. First this is the form of the supergravity action as first noted by Ferrara, Freedman and van Nieuwenhuizen. Secondly, if we recall the component fields in (402) and (403), it is apparent that the variable  $\omega_{\underline{a}\underline{b}\underline{c}}$  does not occur at all among the component fields! In other words the actual evaluation of (408) really leads to the action in (415). The action in (410) is just a very convenient way to re-write the final result.

The most important point about this observation is that any manifestly supersymmetric quantum theory that begins with the action of (408) *can never* lead to a quantum effective action of the form in (305). Instead the pure graviton part of the supersymmetric quantum gravity effective action must take the form of (302).

### 5.9 Local Symmetries of the Component Supergravity Action

The supergravity fields transform under a number of local symmetries. All possess infinitesimal local Lorentz transformation of the form

$$\delta_{L.L.}(\lambda)\mathcal{F} = [ \lambda_\alpha{}^\beta(x)\mathcal{M}_\beta{}^\alpha + \bar{\lambda}_\alpha{}^{\dot{\beta}}(x)\overline{\mathcal{M}}_{\dot{\beta}}{}^{\dot{\alpha}}, \mathcal{F} ] \quad , \quad (418)$$

where  $\mathcal{F} \equiv \{e_{\underline{a}}^{\underline{m}}, \psi_{\underline{a}}^\beta, A_{\underline{a}}, S, P\}$ . Similarly they almost all transform as

$$\delta_{G.C.}(\xi)\mathcal{F} = \xi^{\underline{m}}\partial_{\underline{m}}\mathcal{F} \quad , \quad (419)$$

under an infinitesimal coordinate transformation. The exception is  $e_{\underline{a}}^{\underline{m}}$  for which

$$\delta_{G.C.}(\xi)e_{\underline{a}}^{\underline{n}} = \xi^{\underline{m}}\partial_{\underline{m}}e_{\underline{a}}^{\underline{n}} - e_{\underline{a}}^{\underline{m}}\partial_{\underline{m}}\xi^{\underline{n}} \quad . \quad (420)$$

But the most interesting variations are those under local supersymmetry

$$\begin{aligned} \delta_Q(\epsilon)e_{\underline{a}}^{\underline{m}} &= i(\epsilon^\beta\overline{\psi}_{\underline{a}}^{\dot{\beta}} + \bar{\epsilon}^{\dot{\beta}}\psi_{\underline{a}}^\beta)e_{\underline{b}}^{\underline{m}} \quad , \\ \delta_Q(\epsilon)\psi_{\underline{a}}^\beta &= \mathcal{D}_{\underline{a}}(e, \psi, A)\epsilon^\alpha + i(\epsilon^\beta\overline{\psi}_{\underline{a}}^{\dot{\beta}} + \bar{\epsilon}^{\dot{\beta}}\psi_{\underline{a}}^\beta)\psi_{\underline{b}}^\alpha \\ &\quad + i\bar{\epsilon}_\alpha{}^{\dot{\beta}}\delta_{\dot{\beta}}^\beta(S + iP) - i\epsilon_\alpha A^{\dot{\beta}}{}_{\dot{\alpha}} \quad , \\ \delta_Q(\epsilon)A_{\underline{a}} &= -\frac{1}{2}\left[\epsilon^\gamma\overline{f}_{\gamma\dot{\beta}\alpha}\dot{\beta}_\alpha + \frac{1}{3}\epsilon_\alpha\overline{f}_{\dot{\beta}\alpha}\dot{\beta}_\delta\dot{\delta}^\beta\right] + \text{h.c.} \quad , \\ \delta_Q(\epsilon)(S + iP) &= -\frac{1}{3}\epsilon^\alpha f_{\alpha\dot{\beta}\gamma}\dot{\beta}_\gamma \quad . \end{aligned} \quad (421)$$



In all of these expressions  $\epsilon^\alpha$  and  $\bar{\epsilon}^\alpha$  are functions of  $x$  (i.e. local parameters of supersymmetry).

In the last two lines above, the quantity  $f_{\underline{a}\underline{b}}^\gamma$  is the “supercovariantized curl” of the gravitino (also known as the supercovariantized gravitino field strength) which is explicitly given by

$$\begin{aligned} f_{\underline{a}\underline{b}}^\gamma &= \psi_{\underline{a}\underline{b}}^\gamma + i(\psi_{\underline{a}\beta} A^\gamma_{\dot{\beta}} - \psi_{\underline{b}\alpha} A^\gamma_{\dot{\alpha}}) \\ &\quad - i(S + iP)(\bar{\psi}_{\underline{a}\dot{\beta}} \delta_{\beta}^\gamma - \bar{\psi}_{\underline{b}\dot{\alpha}} \delta_{\alpha}^\gamma) \quad . \end{aligned} \quad (422)$$

The most interesting property of this quantity is that under a local supersymmetry variation, it transforms covariantly (i.e. without terms proportional to spacetime derivatives of  $\epsilon$  or  $\bar{\epsilon}$ ). In a similar manner, the Riemann curvature tensor also possesses a “supercovariantized” version given by

$$\begin{aligned} g_{\underline{a}\underline{b}\underline{c}\underline{d}} &= C_{\dot{\gamma}\dot{\delta}} g_{\underline{a}\underline{b}\gamma\delta} + C_{\gamma\delta} g_{\underline{a}\underline{b}\dot{\gamma}\dot{\delta}} \quad , \\ g_{\underline{a}\underline{b}\gamma\delta} &= r_{\underline{a}\underline{b}\gamma\delta} + (S - iP)\psi_{\underline{a}(\gamma}\psi_{\underline{b}|\delta)} \\ &\quad + i\left\{ \left[ \frac{1}{12}\bar{\psi}_{\underline{a}}\dot{\gamma}f_{(b|\dot{\kappa}|\gamma|\dot{\delta})} + \frac{1}{4}\psi_{\underline{a}\beta}\bar{f}_{(\gamma|\dot{\epsilon}|\delta)}\dot{\epsilon}_{\dot{\beta}} \right. \right. \\ &\quad \left. \left. + \frac{1}{6}\psi_{\underline{a}(\gamma}C_{\delta)\beta}\bar{f}_{\dot{\kappa}\dot{\beta}}\dot{\kappa}_{\dot{\delta}} \right] - (\underline{a} \leftrightarrow \underline{b}) \right\} \quad . \end{aligned} \quad (423)$$

In this expression above, the quantity  $r_{\underline{a}\underline{b}\gamma\delta}$  is defined by

$$r_{\underline{a}\underline{b}\underline{c}\underline{d}} = C_{\dot{\gamma}\dot{\delta}} r_{\underline{a}\underline{b}\gamma\delta} + C_{\gamma\delta} r_{\underline{a}\underline{b}\dot{\gamma}\dot{\delta}} \quad . \quad (424)$$

For this supergravity theory  $r_{\underline{a}\underline{b}\underline{c}\underline{d}}$  is defined as in (266) but where the spin-connection is defined by (411).

If one computes the commutator of two local supersymmetry variations on any field, the answer is

$$\begin{aligned} [\delta_Q(\epsilon_1), \delta_Q(\epsilon_2)] &= \delta_{G.C.}(\xi_{12}) + \delta_{L.L.}(\lambda_{12}) + \delta_Q(\epsilon_{12}) \quad , \\ \xi_{12}^{\underline{m}} &= -i(\epsilon_1^\alpha \bar{\epsilon}_2^\alpha + \bar{\epsilon}_1^\alpha \epsilon_2^\alpha) e_{\underline{a}}^{\underline{m}} \quad , \\ \lambda_{12}^{\underline{b}\underline{c}} &= -i(\epsilon_1^\alpha \bar{\epsilon}_2^\alpha + \bar{\epsilon}_1^\alpha \epsilon_2^\alpha) \omega_{\underline{a}}^{\underline{b}\underline{c}}(e, \psi, A) \\ &\quad + 2[\epsilon_1^{(\beta} \epsilon_2^{\gamma)} C^{\dot{\beta}\dot{\gamma}}(S + iP) + \text{h.c.}] \quad , \\ \epsilon_{12}^\beta &= -i(\epsilon_1^\alpha \bar{\epsilon}_2^\alpha + \bar{\epsilon}_1^\alpha \epsilon_2^\alpha) \psi_{\underline{a}}^\beta \quad . \end{aligned} \quad (425)$$

As can be seen the composition laws for  $\xi_{12}$ ,  $\lambda_{12}$  and  $\epsilon_{12}$  depend on the fields of the supergravity multiplet. For this reason the commutator algebra for the symmetries of supergravity theories are often described as being “field-dependent.”

### 5.10 Supersymmetric Matter Coupling

Coupling matter superfields to the supergravity multiplet works essentially like that described for the component fields coupling of gravity. The key point is to note that local superspace possesses a measure analogous to  $\int d^4x e^{-1}$ . Arnowitt, Nath and Zumino<sup>21</sup> first suggested such integration measures should be written as

$$\int d\mu \equiv \int d^{N_B}x d^{N_F}\theta E^{-1} = \int d^{N_B}x d^{N_F}\theta [sdet(E_{\underline{A}}^{\underline{M}}(\theta, x))]^{-1} , \quad (426)$$

for a superspace of  $N_B$  bosonic coordinate and  $N_F$  fermionic coordinates. For the case of our interest  $N_B = N_F = 4$ . A minimal coupling philosophy suggests that if we start with a supersymmetric action of the form

$$\mathcal{S}_{Matter} = \int d^4x d^2\theta d^2\bar{\theta} \mathcal{L}(\Phi, \bar{\Phi}, D_{\underline{A}}) , \quad (427)$$

its coupling to supergravity may be simply obtained by the replacement

$$\mathcal{S}_{Matter} = \int d^4x d^2\theta d^2\bar{\theta} E^{-1} \mathcal{L}(\Phi, \bar{\Phi}, \nabla_{\underline{A}}) . \quad (428)$$

It is also possible to consider non-minimal coupling by considering the class of actions described by

$$\mathcal{S}_{Non-min.-Matter} = \int d^4x d^2\theta d^2\bar{\theta} E^{-1} \mathcal{L}(\Phi, \bar{\Phi}, \nabla_{\underline{A}}, W_{\alpha\beta\gamma}, G_{\underline{a}}, R) . \quad (429)$$

However, one somewhat surprising feature of superspace supergravity is that it possesses more than one local measure over which integrations may be performed to construct invariant objects. The existence of the “chiral measure” is a local generalization of the “F-terms” invariants that we met in the first lecture. In addition to  $E^{-1}$ , there exists<sup>17</sup> an object  $\mathcal{E}^{-1}$  that permits the definition of a measure

$$\int d\mu_c \equiv \int d^4x d^2\theta \mathcal{E}^{-1} . \quad (430)$$

This measure will lead to an invariant if it is applied to a Lagrangian superfield  $\mathcal{L}_c$  which satisfies the superspace supergravity chirality condition  $\bar{\nabla}_{\dot{\alpha}} \mathcal{L}_c = 0$ . The existence of such a measure is critical so that the superpotential term possesses an extension in the presence of supergravity,

$$\mathcal{S}_{Pot} \equiv \int d^4x d^2\theta \mathcal{E}^{-1} W(\Phi) . \quad (431)$$

Similarly, the chiral measure is important to be able to write the action of supersymmetric Yang-Mills theory in the presence of supergravity,

$$\mathcal{S}_{YM} \equiv \frac{1}{4} \text{Tr} \left[ \int d^4x d^2\theta \mathcal{E}^{-1} W^\alpha W_\alpha + \text{h.c.} \right] . \quad (432)$$

In writing this last equation, we should recognize however, that the superfield  $W_\alpha$  is modified from its rigid definition because in the presence of supergravity a combined supergravity-Yang-Mills supercovariant derivative

$$\nabla_{\underline{A}} = \mathcal{E}_{\underline{A}} + \frac{1}{2} \omega_{\underline{A}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} - i \Gamma_{\underline{A}}^{\mathcal{T}} t_{\mathcal{T}} , \quad (433)$$

must be used to define it.

### 5.11 Theory of Local Superspace Integrations and Component Results

In the case of rigid supersymmetry, we were efficiently able to derive component results from superspace ones via an equation of the form

$$\int d^4x d^2\theta d^2\bar{\theta} \mathcal{L} \equiv \frac{1}{2} \left\{ \int d^4x [D^2 \bar{D}^2 \mathcal{L}] + \text{h.c.} \right\} , \quad (434)$$

where

$$D_\alpha \equiv \partial_\alpha + i \frac{1}{2} \bar{\theta}^{\dot{\alpha}} \partial_{\underline{a}} , \quad \bar{D}_{\dot{\alpha}} \equiv \bar{\partial}_{\dot{\alpha}} + i \frac{1}{2} \theta^\alpha \partial_{\underline{a}} . \quad (435)$$

In the presence of supergravity, it is important to maintain this efficiency and level of completeness in our understanding of the theory. This suggests that there should exist for a general superspace an operator  $\mathcal{D}^{N_F}$  such that

$$\int d^{N_B+N_F} z \mathcal{E}^{-1} \mathcal{L} = \int d^{N_B} z e^{-1} [ \mathcal{D}^{N_F} \mathcal{L} ] , \quad (436)$$

independent of the superfield  $\mathcal{L}$  that appears in this equation and where

$$e^{-1} \equiv [\det(e_{\underline{a}}^{\underline{m}}(x))]^{-1} , \quad \mathcal{D}^{N_F} \mathcal{L} \equiv \lim_{\theta \rightarrow 0} (\mathcal{D}^{N_F} \mathcal{L}) . \quad (437)$$

Also this operator should be such that it can be written as

$$\int d^{N_B} z e^{-1} [ \mathcal{D}^{N_F} \mathcal{L} ] = \int d^{N_B} z e^{-1} \left[ \sum_{i=0}^{N_F} c_{(N_F-i)} (\nabla \cdot \dots \nabla)^{N_F-i} \mathcal{L} \right] , \quad (438)$$

in terms of some field-dependent coefficients  $c_{(N_F-i)}$  and powers of the spinorial superspace supergravity covariant derivative  $\nabla_\alpha$ . It is a remarkable fact that

although the operator  $\mathcal{D}^{N_F}$  has been known on a case-by-case basis for many examples, until quite recently<sup>27</sup> there existed no systematic way to construct these field dependent coefficients.

It is now known that there are two such methods to find these coefficients by the use of; (a.) the super differential calculus of super p-forms and (b.) a normal-coordinate expansion of the  $\theta$  coordinate of superspace. Both of these methods lead to the same result which we now describe. The critical point is the form of an operator  $\mathcal{D}^2$  that takes the form

$$\mathcal{D}^2 \equiv \nabla^2 + i\bar{\psi}_{\underline{a}}^{\dot{\alpha}} \nabla_{\dot{\alpha}} + 3\bar{R} + \frac{1}{2} C^{\alpha\beta} \bar{\psi}_{\underline{a}}^{(\dot{\alpha}} \bar{\psi}_{\underline{b}}^{\dot{\beta})} \quad . \quad (439)$$

Using this we define

$$\begin{aligned} \int d\mu \mathcal{L} &\equiv \int d^4x e^{-1} \left[ \mathcal{D}^4 \mathcal{L} \right] \\ &\equiv \frac{1}{2} \left\{ \int d^4x e^{-1} \left[ \mathcal{D}^2 (\bar{\nabla}^2 + R) \mathcal{L} \right] + \text{h.c.} \right\} \quad , \end{aligned} \quad (440)$$

where the operator  $\mathcal{D}^4$  is defined by

$$\mathcal{D}^4 = \frac{1}{2} \left[ \mathcal{D}^2 (\bar{\nabla}^2 + R) + \bar{\mathcal{D}}^2 (\nabla^2 + \bar{R}) \right] \quad . \quad (441)$$

Similarly, for the case of the chiral measure we define

$$\int d\mu_c \mathcal{L}_c = \int d^4x e^{-1} \left[ \mathcal{D}^2 \mathcal{L}_c \right] \quad . \quad (442)$$

By using the formulae above (that we have named “density projectors”) the problem of finding the explicit component forms of any local superspace action can be reduced to the problems of a series of differentiations, etc.

In the case of supersymmetric Yang-Mills theory, we saw how the component fields also emerged from the field strength supertensors associated with the theory. A similar occurrence takes place for supergravity. In particular looking back at the form of the superspace supergravity commutator algebra, three supertensors are apparent;  $W_{\alpha\beta\gamma}$ ,  $G_{\underline{a}}$  and  $R$ . It is of interest to know the correspondence of these to the component fields of supergravity. After some long calculations starting from the solution to constraints, one can verify

$$G_{\underline{a}}| = A_{\underline{a}} \quad , \quad R| = (S + iP) \quad . \quad (443)$$

In a similar manner it can be shown that

$$T_{\underline{a}\underline{b}}^{\gamma}| = f_{\underline{a}\underline{b}}^{\gamma} \quad , \quad R_{\underline{a}\underline{b}\gamma}^{\delta}| = g_{\underline{a}\underline{b}\gamma}^{\delta} \quad . \quad (444)$$

The lowest component of the superfield  $W_{\alpha\beta\gamma}$  just corresponds to

$$W_{\alpha\beta\gamma}| = \frac{1}{12} C^{\dot{\alpha}\dot{\beta}} f_{(\alpha|\dot{\alpha}|\beta|\dot{\beta}|\gamma)} \quad . \quad (445)$$

Finally, to be able to evaluate most actions completely in terms of component fields, it is useful to know the identities,

$$\begin{aligned} \nabla_{\alpha} R &= -\frac{1}{3} T_{\alpha\dot{\alpha}\gamma}^{\dot{\alpha}\gamma} \quad , \\ \nabla_{\alpha} G_{\underline{b}} &= -\frac{1}{2} \left[ T_{\alpha\dot{\alpha}\beta}^{\dot{\alpha}\beta} \dot{\gamma}_{\dot{\gamma}} - \frac{1}{3} C_{\alpha\beta} T_{\gamma\dot{\beta}}^{\gamma\dot{\delta}} \dot{\delta}_{\dot{\delta}} \right] \quad . \end{aligned} \quad (446)$$

Now we can explain the derivation of (410) from first principles. Since we know that the action for old minimal supergravity is just given by the integral of the volume element of the full superspace, it follows from (440) that

$$\int d\mu \, 1 = \frac{1}{2} \left\{ \int d^4x \, e^{-1} \left[ \mathcal{D}^2 R \right] + \text{h.c.} \right\} \quad . \quad (447)$$

It remains to simply evaluate the differential operator  $\mathcal{D}^2$  acting on  $R$ . The first derivative expression is given just above. For, the second derivative, once again using the technique known as “solving the supergravity Bianchi identities” the appropriate expression can be derived. After completing this, we arrive at the action in (440).

We would be remiss if we did not note that the superspace geometry and its corresponding supergravity theory discussed here is but one of a number of such theories. It is called “old minimal supergravity” to distinguish it from other possibilities. This is an off-shell, irreducible local supersymmetry representation. All other off-shell, irreducible versions of supergravity only differ from this one by the set of constraints on their supergeometries and the auxiliary component fields that appear.

### 5.12 4D, $N = 1$ Supergeometry from Heterotic and Superstrings

Presently many physicists regard the most fundamental of theories as constructs that go well beyond the theory of supergravity. However, a feature most common to many extensions beyond supergravity is that it is recovered in a certain limit<sup>i</sup>. Although it is popular now to discuss a most fundamental theory in terms of “M-theory,” I will be a bit more conservative and limit my considerations to superstrings and heterotic strings.

---

<sup>i</sup>This is another modern version of the correspondence principle.

These are mathematical constructions that are vastly more complicated than is the construction of supergravity. However, it is my personal view that for even these theories, their ultimate formulations are presently a mystery. The solution of this mystery is the central problem of “covariant superstring (and heterotic string) field theory.” For although there has been progress in understanding superstring and heterotic string theory, the simple notion of a ‘stringy’ field strength is at present unknown. My experience with the ultimate formulation of supergravity suggests that there still remains to be given “geometrical” formulations of superstring and heterotic string theory. I expect this belief to ultimately be realized at some point in the future. Then it is natural to expect that the supergeometry of 4D,  $N = 1$  supergravity will be contained in the eka-geometry of some more fundamental theory just the non-Riemannian geometry of gravity is contained in the superspace geometry of Wess-Zumino superspace.

Rather than wait until these putative geometrical formulations of superstring and heterotic strings are found, it is possible to attempt to find the superspace supergravity limit of these by looking for “special versions” of 4D,  $N = 1$  supergravity directly. Although there is a large literature on this topic, in 1988<sup>28</sup> a special version we now called  $\beta FFC$  supergravity was proposed to be this limit. Although additional time will likely pass before a rigorous derivation proves that this is the unique 4D,  $N = 1$  supergravity theory associated with eka-supergravity, strong evidence for this has started to appear recently<sup>29</sup>. So we will end our discussion of supergravity with an eye toward the future.

A most distinctive feature of this theory is that it contains an additional local symmetry (a  $U(1)$  gauge symmetry) for which there does *not* appear a fundamental gauge field. In our book “Superspace” we call such a phenomenon a “fake” gauge symmetry. Thus, the starting point of the  $\beta FFC$  theory is a superspace supercovariant derivative of the form

$$\nabla_{\underline{A}} = E_{\underline{A}} + \omega_{\underline{A}\underline{c}}^{\underline{d}} \mathcal{M}_{\underline{d}}^{\underline{c}} - i \Gamma_{\underline{A}} \mathcal{Y} \quad . \quad (448)$$

This looks very similar to the form of the supergravity-Yang-Mills covariant derivative in (378). It possesses a crucial difference. For an internal symmetry of the Yang-Mills theory we have

$$[t_{\mathcal{I}}, \nabla_{\underline{A}}] = 0 \quad , \quad (449)$$

whereas for the generator  $\mathcal{Y}$  instead we define

$$\begin{aligned} [\mathcal{Y}, \nabla_\alpha] &= -\frac{1}{2} \nabla_\alpha \quad , \\ [\mathcal{Y}, \bar{\nabla}_\alpha] &= \frac{1}{2} \bar{\nabla}_\alpha \quad , \\ [\mathcal{Y}, \nabla_{\underline{a}}] &= 0 \quad . \end{aligned} \tag{450}$$

So in a sense this generator has more in common with  $\mathcal{M}_{\underline{a}\underline{b}}$  for which it is also true that  $[\mathcal{M}_{\underline{a}\underline{b}}, \nabla_{\underline{A}}] \neq 0$ . The complete set of generators that do *not* commute with  $\bar{\nabla}_{\underline{A}}$  is called the “holonomy group.” Thus the holonomy group of  $\beta FFC$  supergravity is  $SL(2, \mathbb{C}) \otimes U(1)$ . The appearance of this  $U(1)$  factor is highly significant. Every known 4D,  $N = 1$  truncation of heterotic or superstring theory possesses such a  $U(1)$  symmetry on its world sheet.

The commutator algebra of  $\beta FFC$  supergravity is considerably different from that of the minimal theory and is given by

$$\begin{aligned} [\nabla_\alpha, \nabla_\beta] &= 0 \quad , \\ [\nabla_\alpha, \bar{\nabla}_{\dot{\alpha}}] &= i \nabla_{\underline{a}} + H_{\beta\dot{\alpha}} \mathcal{M}_\alpha{}^\beta - H_{\alpha\dot{\beta}} \bar{\mathcal{M}}_{\dot{\alpha}}{}^{\dot{\beta}} + H_{\underline{a}} \mathcal{Y} \quad , \\ [\nabla_\alpha, \nabla_{\underline{b}}] &= i(\nabla_\beta H_{\gamma\dot{\beta}}) [\mathcal{M}_\alpha{}^\gamma + \delta_\alpha{}^\gamma \mathcal{Y}] \\ &\quad + i [C_{\alpha\beta} \bar{W}_{\dot{\beta}\dot{\gamma}}{}^{\dot{\delta}} - \frac{1}{3} \delta_{\dot{\beta}}{}^{\dot{\delta}} (2 \nabla_\alpha H_{\beta\dot{\gamma}} + \nabla_\beta H_{\alpha\dot{\gamma}})] \bar{\mathcal{M}}_{\dot{\delta}}{}^{\dot{\gamma}} \quad , \\ [\nabla_{\underline{a}}, \nabla_{\underline{b}}] &= \left\{ \frac{1}{2} C_{\alpha\beta} [i H^\gamma{}_{(\dot{\alpha}} \nabla_{\gamma\dot{\beta})} - (\nabla^\gamma \bar{\nabla}_{(\dot{\alpha}} H_{\gamma|\dot{\beta})}) \mathcal{Y}] \right. \\ &\quad + [C_{\dot{\alpha}\dot{\beta}} (W_{\alpha\beta}{}^\gamma - \frac{1}{6} (\bar{\nabla}^{\dot{\gamma}} H_{(\alpha\dot{\gamma}}) \delta_{\beta}{}^{\gamma)}) - \frac{1}{2} C_{\alpha\beta} (\bar{\nabla}_{(\dot{\alpha}} H^\gamma{}_{\dot{\beta})})] \nabla_\gamma \\ &\quad - C_{\dot{\alpha}\dot{\beta}} [W_{\alpha\beta\gamma\delta} + i \frac{1}{4} C_{\gamma(\alpha} (\nabla_{|\beta}{}^{\dot{\epsilon}} H_{\delta\dot{\epsilon}})] \\ &\quad + \frac{1}{12} C_{\gamma(\alpha} C_{\beta)\delta} (\nabla^\epsilon \bar{\nabla}^{\dot{\epsilon}} H_{\epsilon\dot{\epsilon}})] \mathcal{M}^{\gamma\delta} \\ &\quad \left. + \frac{1}{2} C_{\alpha\beta} [\nabla_\gamma \bar{\nabla}_{(\dot{\alpha}} H_{\delta\dot{\beta})}] \mathcal{M}^{\gamma\delta} + \text{h.c.} \right\} \quad , \end{aligned} \tag{451}$$

As we have described the theory above, we see that only the components of the torsion supertensor ( $T_{\alpha\dot{\underline{b}}}$  and  $T_{\underline{a}\underline{b}\underline{c}}$ ) are non-zero.

The irreducible Bianchi identities associated with this theory are given by,

$$\begin{aligned} \nabla^{\underline{a}} H_{\underline{a}} &= 0 \quad , \quad \bar{\nabla}_{\dot{\beta}} W_{\alpha\beta\gamma} = 0 \quad , \quad \nabla^\beta \nabla_\beta H_{\underline{a}} = 0 \quad , \\ \nabla^\alpha W_{\alpha\beta\gamma} &= -\frac{1}{6} \nabla_{(\beta} \bar{\nabla}^{\dot{\gamma}} H_{\gamma)\dot{\gamma}} - \frac{1}{2} \bar{\nabla}^{\dot{\gamma}} \nabla_{(\beta} H_{\gamma)\dot{\gamma}} \quad . \end{aligned} \tag{452}$$

As in our previous discussion, the effects of the constraints above are to determine now both  $\omega_{\underline{A}\underline{a}\underline{b}}$  and  $\Gamma_{\underline{A}}$  in terms of other quantities.

In addition to the supergravity covariant derivative, there is also the super 2-form  $B_{\underline{A}\underline{B}}$  whose super exterior derivative  $H_{\underline{A}\underline{B}\underline{C}}$  satisfies the conditions,

$$H_{\alpha\beta\gamma} = H_{\alpha\beta\dot{\gamma}} = H_{\alpha\beta\gamma} = H_{\alpha\beta\underline{c}} = H_{\alpha\dot{\beta}\underline{c}} - i\frac{1}{2}C_{\alpha\gamma}C_{\dot{\beta}\dot{\gamma}} = 0 \quad ,$$

$$H_{\alpha\underline{bc}} = 0 \quad , \quad H_{\underline{abc}} = i\frac{1}{4}[C_{\beta\gamma}C_{\dot{\alpha}(\dot{\beta}}H_{\alpha\dot{\gamma})} - C_{\dot{\beta}\dot{\gamma}}C_{\alpha(\beta}H_{\gamma)\dot{\alpha}}] \quad . \quad (453)$$

There is an interesting similarity of the components of the super 3-form field strength and the components of the torsion supertensor. Namely, the only nonvanishing components of the latter are  $H_{\alpha\dot{\beta}\underline{c}}$  and  $H_{\underline{abc}}$ .

There is one more supertensor required to give a complete description of 4D,  $N = 1$   $\beta$ FFC supergravity. We denote this additional tensor by  $L$ . The spinorial derivatives of this superfield are particularly important since

$$\begin{aligned} \nabla_{\alpha}L &= -\frac{4}{3}\chi_{\alpha} \quad , \quad \nabla_{\alpha}\chi_{\beta} = C_{\alpha\beta}\overline{R} \quad , \\ \nabla_{\alpha}\bar{\chi}_{\dot{\alpha}} &= \frac{1}{4}H_{\alpha\dot{\alpha}} + \frac{1}{2}G_{\alpha\dot{\alpha}} - i\frac{3}{8}\nabla_{\underline{a}}L \quad . \end{aligned} \quad (454)$$

Thus, the three basic field strength superfields of 4D,  $N = 1$   $\beta$ FFC supergravity are  $W_{\alpha\beta\gamma}$ ,  $H_{\underline{a}}$  and  $L$ . However, note the total absence of  $L$  in (451) and (453)

It is appropriate here to make some comments about how the component fields arise in  $\beta$ FFC supergravity in contrast to their appearance in old minimal supergravity. At lowest order in  $\theta$ , the superfields  $W_{\alpha\beta\gamma}$  and  $H_{\underline{a}}$  correspond to the supercovariantized “curls” of the gravitino and axion component fields. The lowest component of  $W_{\alpha\beta\gamma\delta}$  corresponds to the supercovariantized Weyl tensor (as usual the remaining components of the Riemann curvature are at higher orders in the superfields). The lowest order components of  $L$  and  $\chi_{\alpha}$  are the dilaton and dilatino, respectively. Finally the lowest order components of  $G_{\underline{a}}$  and  $R$  correspond to the usual auxiliary fields of old minimal supergravity.

Given the structure of  $\beta$ FFC supergeometry, its action is given by

$$\mathcal{S}_{\beta FFC SG} = -\frac{3}{\kappa^2} \int d^4x d^2\theta d^2\bar{\theta} E^{-1} e^{f(L)} \quad , \quad (455)$$

where  $f$  is any function of  $L$ . Even though the component fields of  $\beta$ FFC supergravity correspond to those of minimal supergravity ( $(e_{\underline{a}}^{\underline{m}}(x), \psi_{\underline{a}}^{\beta}(x), A_{\underline{a}}(x), S(x), P(x))$  plus a super 2-form multiplet  $(\varphi(x), b_{\underline{a}\underline{b}}(x), \chi_{\beta}(x))$ ) it is a remarkable fact that for  $\beta$ FFC supergravity,

$$-\frac{3}{\kappa^2} \int d^4x d^2\theta d^2\bar{\theta} E^{-1} = 0 \quad . \quad (456)$$



We emphasize this result holds for the  $\beta FFC$  supervielbein that satisfies (451) not the one that satisfies (405). We believe the fact that the  $\beta FFC$  supergravity action cannot be written without the use of  $L$  (the field strength of the tensor multiplet) is a good indication that the two multiplets are intimately related. That such a relation should be present is well representative of heterotic and superstring theory where it is known that both multiplets describe the “stringy” supergravitational sector of the theories.

Within the confines of  $\beta FFC$  supergravity coupling to matter fields is distinctly different than in the old minimal theory. For example, a  $\beta FFC$  SG-chiral multiplet coupled system has the form

$$\begin{aligned} \mathcal{S} = & -\frac{3}{\kappa^2} \int d^4x d^2\theta d^2\bar{\theta} E^{-1} \left[ e^{f(L)} + K(\Phi, \bar{\Phi}) \right] \\ & + \left\{ \int d^4x d^2\theta \mathcal{E}^{-1} W(\Phi) + \text{h.c.} \right\} , \end{aligned} \quad (457)$$

and when the coupling of the spin-0 fields is evaluated, it can be shown to possess no “improvement terms” of the form  $A\bar{A}\eta^{\underline{a}\underline{c}}r_{\underline{a}\underline{b}\underline{c}}^{\underline{b}}$ . Evaluation of this superfield action using the old minimal supergeometry does to “improvement terms.” Finally, the chiral volume of  $\beta FFC$  supergravity is proportional to the mass-like term for the gravitino,

$$\int d^4x d^2\theta \mathcal{E}^{-1} = \frac{1}{2} \int d^4x C^{\alpha\beta} \bar{\psi}_{\underline{a}}^{(\dot{\alpha}} \bar{\psi}_{\underline{b}}^{\dot{\beta})} . \quad (458)$$

## *Epilogue*

We have come to the end of this series of lectures. I have a great desire to say, “Free at last, free at last,...,” but these have only begun to brush the surface of the current developments in modern relativistic theories of fundamental objects. I have enjoyed lecturing to an enthusiastic set of aspiring physicists. Beyond the simply mastery of numbers of topics that were touched upon in the lectures there looms superstring and heterotic string field theory and even more fearsome ‘beasts.’

The role of superstrings and heterotic strings is to ‘control’ many aspects of the theories we have discussed. For example, super and heterotic strings are very restrictive and arbitrary choices for gauge groups (i.e.  $V^T$ ), numbers and representations of matter (i.e.  $\Phi^I$  and perhaps even  $\Sigma^I$ ) and ‘stringy’ supergravity (i.e.  $\beta FFC$  SG) are not in general consistent. Thus, 4D,  $N = 1$  super and heterotic string theory only permit special choices. In the realm of quantum theories, 4D,  $N = 1$  superstrings and heterotic strings are the only known models where nonlinear expressions like (201), (233), (235), (307), (302) and (429) may derived to arbitrary order in any of the fields. So the long dreamt about reconciliation between quantum theory and gravitation has apparently been accomplished. From this point of view, it remains to simply discover seemingly the answer to a few things.

Even beyond strings, more recently, there has been suggested the existence of ‘M-theory’ and relatives. However, I believe that in our rush to move even further from the known physics of fundamental particles there have been left unsolved major problems even in the realm of string theory. The principal among these is that we do not possess a fundamental understanding of ‘pre-geometry’ beyond the realm of supergravity. As we have seen in these lectures, Salam-Strathdee superspace provides a comprehensive concept upon which to investigate supersymmetrical theories. It is a fact that for even the simplest superstring and heterotic string theories, there does not exist the analog of Salam-Strathdee superspace. While many physicists have regarded this as of not much import and have continued to increase our understanding of ever more imaginative mathematical constructions, it is my opinion that until we possess a truly geometrical understanding of strings we will not have achieved the same comprehensive level of understanding of these theories as that which is Einstein’s legacy embodied by the “equivalence principle.” This lack in our understanding will continue to offer a deep challenge to the theoretical understanding of fundamental physics.

## Acknowledgments

I wish to acknowledge the support for my research program under grants # PHY-96-43219 from the U.S. National Science Foundation and # CRG-93-0789 from NATO.

## Bibliography

For the student wishing to continue to study supersymmetry there are by now many textbooks available. At the introductory level, I highly recommend

- [1.] J. Wess and J. Bagger, “Supersymmetry and Supergravity,” Princeton Univ. Press, Princeton, NJ (1983).
- [2.] P. West, “Introduction to Supersymmetry and Supergravity,” World Scientific, Singapore (1986).

At the comprehensive level, there are only two books that treat the subject in real depth.

- [3.] S.J.Gates, Jr., M.T.Grisaru, M. Roček and W. Siegel, “Superspace or One Thousand and One Lessons in Supersymmetry,” Benjamin Cummings (Addison-Wesley), Reading, MA (1983).
- [4.] I. Buchbinder and S. Kuzenko, “Ideas and Methods of Supersymmetry and Supergravity or A Walk Through Superspace,” IOP Publishing, Bristol and Philadelphia, (1995).

The first of these is out of print, so you may have to look in your library or write to the publisher. These latter two books nicely compliment each other covering slightly different matters. If you master these things, you will be an expert.

## References

1. A. Salam and J. Strathdee, *Phys. Rev. D* **11** (1975), 1521; *Nucl. Phys. B* **B86** (1975), 142; S.Ferrara, J. Wess and B. Zumino, *Phys. Lett. B* **51** (1974), 239.
2. S.J.Gates, Jr., “Symmetry Principles in Selected Problems in Field Theory,” Ph.D. Thesis, MIT (1977), p. 51-52, unpublished.

3. F.A.Berezin, *The Method of Second Quantization*, Academic Press, New York, (1966).
4. Yu. A. Gol'fand and E. P. Likhtman, JETP Lett. **13** (1971) 323; J. Wess and B. Zumino, *Nucl. Phys. B* **70** (1974), 39.
5. J. Wess and B. Zumino, *Phys. Lett. B* **49** (1974), 52.
6. S.J.Gates, Jr., K. S. Stelle and P. C. West, *Nucl. Phys. B* **B169** (1980), 347.
7. S.J.Gates, Jr. and W.Siegel, *Nucl. Phys. B* **164** (1980), 484.
8. A. Karlhede, U. Lindstrom, M. Roček and G. Theodoridis, *Nucl. Phys. B* **294** (1987), 498.
9. B.Zumino, *Phys. Lett. B* **87** (1979), 205.
10. S.J.Gates, Jr., *Nucl. Phys. B* **238** (1984), 349.
11. N. Seiberg, *Phys. Lett. B* **206** (1988), 75.
12. N. Seiberg and E. Witten, *Nucl. Phys. B* **426** (1994), 19; ERRATUM-*ibid. Nucl. Phys. B* **430** (1994), 485.
13. S.J.Gates, Jr., *Nucl. Phys. B* **184** (1981), 390.
14. W.Siegel, *Phys. Lett. B* **85** (1979), 333.
15. S.J.Gates, Jr. and W.Siegel, *Nucl. Phys. B* **187** (1981), 389.
16. S. J. Gates, Jr., *Phys. Lett. B* **366** (1996), 132,, *idem.*, "A New Proposed Description of the 4D,  $N = 1$  Supersymmetric Effective Action for Scalar Multiplets" to appear in the proceedings of the Second International Sakharov Conference on Physics, May 20-25, Moscow, Russia; *idem.* Univ. of Maryland Preprint, UMDEPP 96-99 (hep-th/9508153), *idem.*, *Nucl. Phys. B* **485** (1997), 145; S.J.Gates, Jr., M.T.Grisaru, M.Knutt-Wehlau, M. Roček and O. Soloviev, *Phys. Lett. B* **396** (1997), 167; "N = 1 Supersymmetry and the Phenomenological Pion Physics Effective Action" in the proceedings of the Supersymmetry '97 meeting, Proceedings Supplement, Nucl.Phys.B. (North-Holland) pp. 171-181.
17. S.J.Gates, Jr. and W.Siegel, *Nucl. Phys. B* **147** (1979), 77; *Nucl. Phys. B* **163** (1980), 519.
18. R.P.Feynman, F.B.Moringo and W.G.Wagner, "Feynman Lectures on Gravitation" ed. B.Hatrfield, Addison-Wesley, Reading, MA (1995).
19. B.Zwiebach, *Nucl. Phys. B* **336** (1990), 185; *Phys. Lett. B* **241** (1990), 343; *Nucl. Phys. B* **390** (1993), 33.
20. M. Grisaru, A. Van Proeyen, D. Zanon, *Nucl. Phys. B* **502** (1997), 345.
21. R.Arnowitz, B.Zumino and P.Nath, *Phys. Lett. B* **56** (1975), 81.
22. V.Ogievetsky and E. Sokatchev, *Nucl. Phys. B* **124** (1977), 309; S.ferrara and B.Zumino, *Nucl. Phys. B* **134** (1978), 301.
23. W.Siegel, "Supergravity Superfields Without a Supermetric," Harvard Univ. preprint HUTP-77/A068 (1977) unpublished; "The Superfield

- Supergravity Action,” Harvard Univ. preprint HUTP-77/A080 (1977) unpublished.
24. J. Wess and B. Zumino, *Phys. Lett. B* **66** (1977), 361; *Phys. Lett. B* **74** (1978), 1.
  25. D.Z.Freedman, S.Ferrara and P.van Nieuwenhuizen, *Phys. Rev. D* **13** (1976), 3214
  26. S.Ferrara and P.van Nieuwenhuizen, *Phys. Lett. B* **74** (1978), 333
  27. S.J. Gates, Jr., “Ectoplasm Has No Topology: The Prelude,” invited talk at the *International Seminar on Supersymmetries and Quantum Symmetries*, Dubna, Russia, (22-26 July 1997), to appear in proceedings, Univ. of MD preprint UMDEPP 98-13, hep-th/9709104; S.J.Gates, Jr., M.T.Grisaru, M.E.Knutt-Wehlau, and W. Siegel, “Component Actions from Curved Superspace: Normal Coordinates and Ectoplasm,” Bradeis Univ. preprint BRX-TH-421, McGill Univ. preprint McGill/97-31, Univ. of Maryland preprint 98-47, SUNY, Stony Brook preprint ITP-SB-97-65, hep-th/9711151.
  28. S.J.Gates, Jr., P.Majumdar, R.Oerter, and A.E.M. van de Ven, *Phys. Lett. B* **214** (1988), 26.
  29. J.de Boer and K.Skenderis, *Nucl. Phys. B* **481** (1996), 129.